AP (19.1) (13) 1Φ.15. 24
26

The two major systemic veins → & the large coronary vein aka the Coronary sinus that drains the heart myocardium empty into the right atrium,

Superior Vena Cava → drains blood from regions superior to the diaphragm

→ i.e to the head, neck, upper limbs & (thoracic region)

Superior Vena Cava → empties into the superior & posterior portions of the right atrium.

Inferior Vena Cava → drains blood from areas inferior to the diaphragm

The lower limbs - however, it too empties into the posterior portion of the atria, but inferior to the opening of the superior vena cava

Immediately superior & slightly medial to the opening of the inferior vena cava on the posterior surface of the atrium is the opening of the coronary sinus.

It is a thin-walled vessel drains most of the coronary veins systemic blood from the heart.

The major of the internal heart structures discussed in this & subsequent Bulk of the right atrium is smooth, the depression of the fossa ovalis is medial, & anterior surface demonstrates prominent, & ridges of muscle called pectinate muscles

The right auricle is also has pectinate muscles but → the left atrium does not have pectinate muscles except in the auricle

The atria receive venous blood nearly on cont basis this prevents venous flow from stopping while the ventricles are contracting

Just Remember → the tricuspid Valve → the opening between the atrium & ventricle is guarded by the tricuspid valve