while the ventricular filling occurs while the atria are relaxed, they do demonstrate a contractile phase & activity pump ~~just~~ blood into the ventricle's just prior to ventricular contraction.

Right Ventricle → The right ventricle receives blood from the right atrium through the ~~tricusp~~ tricuspid valve.

Chordae tendineae → tendinous cords

↳ each Flap of the valve is attached to strong strands of connective tissue. (heart Strings)

There are several chordae tendineae associated with each ~~other~~ of the flaps They are composed of approx 8φ % collagenous fibers, with the remainder consisting of elastic fibers & endothelium

The chordae tendineae connect each of the flaps to a **papillary muscle** that extends from the inferior ventricular surface.

There are 3 papilary muscles in the right ventricle, called the anterior, posterior, & septal muscles. Which correspond to 3 sections of the valves.

When the myocardium of the ventricle contracts → pressure → within the ventricular chamber rises

Blood moves → high pressure ~~to~~ low pressure.

here toward the pulmonary trunk & the atrium ←, To prevent backflow. The papillary muscles also contract, generate tension ~~of~~ on the chordae tendineae. → This prevents the flaps of the valves from being forced into the atria & regurgitation of the blood back into the atria during ventricular contraction.

Trabeculae Carneae → the walls

of the ventricle are lined with trabeculae carneae. These are ridges of cardiac muscle covered by endocardium. In addition to these muscular ridges, a band of cardiac muscle, also covered by endocardium