AP (19.2) (33) 10.16.20
26

Atrioventricular Bundle

arising from the AV → node, the AV or bundle of His → through the interventricular septum before dividing into two ↓ atrioventricular bundle branches → commonly called the left ventricle & the right bundle branch the right ventricle. → Since the ventricle is much larger than the right, the left bundle branches is also {{ why is it larger ?? }) larger than right. Portions of the right bundle branch are found in the moderator bundle & supply the right papillary muscle.

* Because of this connection, each papillary muscle receives the impulse @ approx the same time so they begin to contract stimultaneously just prior to the remainder of the ~~myoc~~ myocardial contractile contraction.

{ This is ~~Bete~~ believed to allow tension to develop on the chordae ~~tin~~ tendineae ~~prot~~ prior to the right ventricular contraction.

→ There is no corresponding moderator band on the left. Both bundle descend & reach the apex of the heart where they connect with the Purkinje fibers. → takes about 25 ms

The Purkinje Fibers

→ are additional myocardial conductive fibers that spread the impulse to the myocardial contractile cells in the ventricles

They extend throughout the myocardium from the Apex of the heart toward the ~~ant~~ atrioventricular septum & the base of the ~~heart~~ heart.

The Purkinje fibers have a fast ~~inherent~~ inherent conduction rate & the electrical impulse reaches all of the ventricular muscle.