


Admissible Continuation

A Mathematical Theory of Persistent Systems


The defining question: How do systems change completely, yet remain identically themselves?
Based on the monograph by Flyxion.



THE UNIVERSAL PROBLEM OF STRUCTURAL PRESERVATION


These systems operate through entirely different physical mechanisms, yet confront an identical mathematical problem: They must execute continuous state transitions without destroying the invariant conditions that make future transitions possible.

BIOLOGY




CONTINUOUS MOLECULAR TURNOVER
WITH INVARIANT CELL WALL INTEGRITY

SOFTWARE





SUCCESSIVE CODE DEPLOYMENTS
WITH PRESERVED API CONTRACTS

INSTITUTIONS




ACTIVE RECONSTRUCTION & AMENDMENT
ON SOLID FOUNDATIONAL STRUCTURE

A fundamental shift in mathematical primitives

	Traditional Dynamics	Admissible Continuation
Fundamental Primitive	States (isolated points in space)	Transitions (relational vectors across time) 
Primary Objective	Optimization/Equilibrium (reaching an ideal end)	Infinite Extendability (preserving future options) 
System Rules	Arbitrary external constraints	Internally derived boundary invariants 
Definition of Failure	Sub-optimal configuration	Irreversible loss of distinguishability 

Persistence begins with mathematical distinguishability




- Evolution is simply the preservation or destruction of a boundary.
- Before mechanics, dynamics, or rules can exist, a system must first be mathematically distinguishable from its environment.
- A transition that destroys this boundary dissolves the system into undifferentiated noise.

Observation partitions the universe into recoverable information

Persistence Ladder

1. DISTINCTION
2. OBSERVATION
3. INVARIANTS
4. ADMISSIBILITY
5. REPAIR
6. RECURSION



1. Distinguishability is always relative to an observer.


2. **Observation maps** partition the configuration space into distinct equivalence classes.

3. **Information** is not an independent primitive—it is the quantitative expression of this distinguishability surviving a transition.

Invariants are the structural boundaries that prevent dissolution

Persistence Ladder

1. DISTINCTION
2. OBSERVATION
3. INVARIANTS
4. ADMISSIBILITY
5. REPAIR
6. RECURSION




- Invariants are not arbitrary rules imposed from the outside.
- They are the structural properties that must survive for the system's defining distinctions to remain recoverable.
- Multiple invariants aggregate to define a **unified persistence threshold**.

Persistence is the existence of infinitely extendable trajectories

Persistence Ladder

1. DISTINCTION
2. OBSERVATION
3. INVARIANTS
4. ADMISSIBILITY
5. REPAIR
6. RECURSION



The primary mathematical object is the **transition**.


A transition is **admissible** only if it preserves the **invariant structures**.

A system **persists** precisely when there exists a **continuation** whose successive transformations preserve the conditions for further continuation.

Robustness requires a rich reachability set

Persistence Ladder

1. DISTINCTION
2. OBSERVATION
3. INVARIANTS
4. ADMISSIBILITY
5. REPAIR
6. RECURSION




Robustness is not resisting physical change or maintaining static equilibrium. It is **Continuation Capacity**: possessing a structurally rich landscape of admissible future options.

Degradation is the loss of recoverable organization

Persistence Ladder

1. DISTINCTION
2. OBSERVATION
3. INVARIANTS
4. ADMISSIBILITY
5. REPAIR
6. RECURSION



- We do not measure how far a system has physically traveled from its origin.
- We measure how much recoverable organization has degraded, pushing the system toward irrecoverable dissolution.

Repair restores future options rather than recreating the past

1. DISTINCTION


2. OBSERVATION

3. INVARIANTS

4. ADMISSIBILITY

5. REPAIR

6. RECURSION




Repair is a geometric projection back onto the admissible manifold. Its goal is to minimize reconstruction defect while maximizing future continuation capacity, completely independent of historical exactness.

A calculus of operators governs the transition lifecycle

$$\Pi = C \circ Q \circ P$$

Persistence Ladder

1. DISTINCTION
2. OBSERVATION
3. INVARIANTS
4. ADMISSIBILITY
5. REPAIR
6. RECURSION



Every complex persistent computation factors into a canonical progression of primitive operators. Unconstrained generation is strictly separated from warranted state commitment.

True persistence is the indefinite continuation of repair

1. DISTINCTION


2. OBSERVATION

3. INVARIANTS

4. ADMISSIBILITY

5. REPAIR


6. RECURSION



- Persistence is not a single successful repair event.
- It emerges when recursive continuation systems continuously self-maintain, ensuring the reconstruction defect remains strictly bounded over time.


Persistent systems survive by continuously evolving their own rules

1. DISTINCTION
2. OBSERVATION
3. INVARIANTS
4. ADMISSIBILITY
5. REPAIR
6. RECURSION




Biological adaptation, institutional reform, and machine learning algorithms all exhibit operator dynamics. True persistence requires adaptation. The repair mechanisms themselves must evolve to maintain admissibility in changing environments.

The Architecture of Persistence



Each mathematical level logically necessitates the preceding structures.
None can be removed without dissolving the system.

Viability through the discipline of preservation



Persistence cannot be understood through reachable states, optimal actions, or generated outputs alone.

Long-term viability emerges not from the unrestricted capacity to generate possibilities, but from the disciplined preservation of the boundaries that permit meaningful continuation across time.