

# Admissibility Geometry and the MOND Transition

Reconstruction Obstruction as an Alternative to Dark Matter

Flyxion

May 10, 2026

## Abstract

We develop a reinterpretation of gravitational anomalies — including flat rotation curves, weak lensing excesses, and the MOND acceleration scale — within the Relativistic Scalar-Vector Plenum (RSVP) framework. Existing emergent gravity programs derive spacetime geometry from physical exchange processes, thermodynamic bookkeeping, or information flow. RSVP inverts this order: admissibility geometry is treated as primary, and persistent physical exchange is interpreted as a stabilized reconstruction mode within that geometry.

The framework introduces the coupled field  $X = (\Phi, \mathbf{v}, S)$ , where  $\Phi$  encodes the accessibility geometry of continuation space,  $\mathbf{v}$  encodes directed coherence flow, and  $S$  measures admissibility dispersion. The dimensionless ratio  $S/\Phi$  serves as a central control parameter governing local reconstruction factorization. We define an *admissibility radius*  $R_A$  as the largest scale over which local reconstruction remains approximately factorizable without cosmological boundary data, and conjecture that the RSVP field equations determine  $R_A \propto c/H_0$  in weak-gradient, low-density regimes, yielding the MOND acceleration scale  $a_0 \sim cH_0$  as a geometric consequence of reconstruction structure rather than a modification of force law.

The paper operates across four explicitly distinguished epistemic registers: definitions, structural conjectures, derived implications, and open proof obligations. We do not derive MOND from completed RSVP field equations; we identify the geometric objects from which such a derivation would have to proceed. Appendices develop the formal sheaf structure, coarse-graining flow, candidate dynamical equations, obstruction cohomology, relation to renormalization theory, observational targets, locality clarification, and formal implementation directions including BNF grammar and Lean-style proof schemas.

*Keywords:* emergent gravity, MOND, admissibility geometry, reconstruction sheaf, entropic gravity, dark matter, RSVP framework, obstruction cohomology, renormalization flow.

## Core Objects and Notation

| Symbol                                                                        | Meaning                                                                                |
|-------------------------------------------------------------------------------|----------------------------------------------------------------------------------------|
| $X = (\Phi, \mathbf{v}, S)$                                                   | Coupled RSVP field                                                                     |
| $\Phi$                                                                        | Continuation accessibility scalar field                                                |
| $\mathbf{v}$                                                                  | Coherence-flow vector field                                                            |
| $S$                                                                           | Admissibility dispersion (entropy) field                                               |
| $\chi = S/\Phi$                                                               | Reconstruction control parameter                                                       |
| $\chi_\ell = S_\ell/\Phi_\ell$                                                | Scale-dependent control parameter                                                      |
| $\mathcal{U} = \{U_\alpha\}$                                                  | Reconstruction (admissibility) cover                                                   |
| $\mathcal{F}$                                                                 | Reconstruction sheaf                                                                   |
| $\mathcal{F}(U_\alpha)$                                                       | Admissible sections over neighborhood $U_\alpha$                                       |
| $\rho_{\alpha\beta}$                                                          | Restriction map $\mathcal{F}(U_\alpha) \rightarrow \mathcal{F}(U_\alpha \cap U_\beta)$ |
| $R_A$                                                                         | Admissibility radius                                                                   |
| $a_A$                                                                         | Admissibility acceleration scale $\sim c^2/R_A$                                        |
| $\mathcal{P}[s]$                                                              | Persistence functional of section $s$                                                  |
| $\mathcal{R}_\lambda$                                                         | Coarse-graining operator on covers                                                     |
| $\Omega_\ell[s]$                                                              | Scale-dependent obstruction functional                                                 |
| $\beta(\chi)$                                                                 | Admissibility beta-function                                                            |
| $\Pi_{\text{therm}}, \Pi_{\text{info}}, \Pi_{\text{grav}}, \Pi_{\text{conf}}$ | Entropy projection functors                                                            |
| $\check{H}^k(\mathcal{U}, \mathcal{F})$                                       | Čech cohomology of cover with sheaf coefficients                                       |

# Contents

|          |                                                                         |           |
|----------|-------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                                     | <b>4</b>  |
| <b>2</b> | <b>Admissibility Geometry and the Reconstruction Sheaf</b>              | <b>5</b>  |
| 2.1      | The Coupled Field . . . . .                                             | 5         |
| 2.2      | Admissible Local Sections . . . . .                                     | 6         |
| 2.3      | The Reconstruction Cover . . . . .                                      | 6         |
| 2.4      | Two Types of Obstruction . . . . .                                      | 7         |
| 2.5      | The Reconstruction Sheaf . . . . .                                      | 7         |
| 2.6      | Bridge to Section 3 . . . . .                                           | 8         |
| <b>3</b> | <b>The Admissibility Phase Transition</b>                               | <b>8</b>  |
| 3.1      | The Factorization Transition . . . . .                                  | 8         |
| 3.2      | The Admissibility Radius . . . . .                                      | 9         |
| 3.3      | Newtonian and Cosmologically Coupled Regimes . . . . .                  | 9         |
| 3.4      | The Central Conjecture . . . . .                                        | 10        |
| 3.5      | Open Proof Obligation . . . . .                                         | 10        |
| 3.6      | Bridge to Section 4 . . . . .                                           | 11        |
| <b>4</b> | <b>Entropy Species as Regime Projections</b>                            | <b>11</b> |
| 4.1      | Entropy Channels as Projections . . . . .                               | 11        |
| 4.2      | Entropy Convergence Under Strong Factorization . . . . .                | 12        |
| 4.3      | Entropy Divergence Under Dispersion Obstruction . . . . .               | 12        |
| 4.4      | The Kastner–Schlatter Contrast . . . . .                                | 13        |
| 4.5      | Bridge to Section 5 . . . . .                                           | 13        |
| <b>5</b> | <b>Dark Matter as Reconstruction Obstruction</b>                        | <b>13</b> |
| 5.1      | The Observational Problem . . . . .                                     | 13        |
| 5.2      | Compatibility and Dispersion Obstruction in Galactic Dynamics . . . . . | 14        |
| 5.3      | Rotation Curves as Persistence Failure . . . . .                        | 14        |
| 5.4      | Weak Lensing and Extension Geometry . . . . .                           | 14        |
| 5.5      | Entropy Divergence as Observational Signal . . . . .                    | 15        |
| 5.6      | Situating the Reinterpretation . . . . .                                | 15        |
| <b>6</b> | <b>Comparison with Neighboring Programs</b>                             | <b>15</b> |
| 6.1      | Organizing Principle . . . . .                                          | 15        |
| 6.2      | Standard Cosmology ( $\Lambda$ CDM) . . . . .                           | 16        |
| 6.3      | MOND and TeVeS . . . . .                                                | 16        |
| 6.4      | Verlinde’s Emergent Gravity . . . . .                                   | 16        |
| 6.5      | Transactional Entropic Gravity . . . . .                                | 17        |
| 6.6      | The Inversion as Theoretical Commitment . . . . .                       | 17        |

|                                                                     |           |
|---------------------------------------------------------------------|-----------|
| <b>7 Conclusion</b>                                                 | <b>17</b> |
| 7.1 The Central Reinterpretation . . . . .                          | 17        |
| 7.2 What the Paper Claims and Does Not Claim . . . . .              | 18        |
| 7.3 Open Problems as Evidence of Structural Coherence . . . . .     | 18        |
| 7.4 Final Philosophical Position . . . . .                          | 19        |
| <b>A Formal Structure of the Reconstruction Sheaf</b>               | <b>19</b> |
| A.1 Presheaf and Sheaf Structure . . . . .                          | 19        |
| A.2 Admissible Sections and Persistence . . . . .                   | 20        |
| A.3 Persistence Functional . . . . .                                | 20        |
| A.4 Dynamical Cover and Gluing . . . . .                            | 21        |
| <b>B Coarse-Graining Flow and the Admissibility Order Parameter</b> | <b>21</b> |
| B.1 Scale-Dependent Fields and Order Parameter . . . . .            | 21        |
| B.2 Admissibility Beta-Function . . . . .                           | 22        |
| B.3 Universality Classes . . . . .                                  | 23        |
| B.4 Admissibility Radius from Flow . . . . .                        | 23        |
| <b>C Candidate Dynamical Equations for the RSVP Fields</b>          | <b>23</b> |
| C.1 Continuation Accessibility Field $\Phi$ . . . . .               | 23        |
| C.2 Coherence Flow Field $\mathbf{v}$ . . . . .                     | 23        |
| C.3 Entropy-Dispersion Field $S$ . . . . .                          | 24        |
| C.4 Control Parameter Dynamics . . . . .                            | 24        |
| C.5 Epistemic Status . . . . .                                      | 24        |
| <b>D Obstruction Cohomology and Global Reconstruction Failure</b>   | <b>24</b> |
| D.1 Čech Cohomology of the Reconstruction Cover . . . . .           | 24        |
| D.2 Two Obstruction Notions . . . . .                               | 25        |
| D.3 Scale-Dependent Obstruction Functional . . . . .                | 25        |
| D.4 Galactic Anomalies as Cohomological Failure . . . . .           | 25        |
| <b>E Relation to Renormalization and Effective Field Theory</b>     | <b>26</b> |
| E.1 Where the Analogy Holds . . . . .                               | 26        |
| E.2 Where the Analogy Breaks . . . . .                              | 26        |
| E.3 Newtonian Gravity as Effective Reconstruction Theory . . . . .  | 26        |
| E.4 Entropy Equivalence as Emergent Universality . . . . .          | 27        |
| <b>F Observational and Experimental Targets</b>                     | <b>27</b> |
| F.1 Rotation Curve Transition Structure . . . . .                   | 27        |
| F.2 Entropy-Channel Divergence . . . . .                            | 27        |
| F.3 Weak Lensing Signatures . . . . .                               | 27        |
| F.4 Cosmological Structure Formation . . . . .                      | 28        |
| F.5 Observational Failure Modes . . . . .                           | 28        |

|          |                                                                            |           |
|----------|----------------------------------------------------------------------------|-----------|
| <b>G</b> | <b>Locality, Causality, and Admissibility Nonlocality</b>                  | <b>28</b> |
| G.1      | The Central Distinction . . . . .                                          | 28        |
| G.2      | Admissibility Nonlocality Defined . . . . .                                | 29        |
| G.3      | Analogy with Gauge Theory and General Relativity . . . . .                 | 29        |
| G.4      | Interpretation of “Cosmological Gluing Data” . . . . .                     | 30        |
| G.5      | Formal Statement . . . . .                                                 | 30        |
| <b>H</b> | <b>Formal Implementation: Grammar, Typed Structures, and Proof Schemas</b> | <b>30</b> |
| H.1      | BNF Grammar for Reconstruction Objects . . . . .                           | 30        |
| H.2      | BNF for Epistemic Status of Claims . . . . .                               | 32        |
| H.3      | Typed Structures in Lean Style . . . . .                                   | 32        |
| H.4      | Reconstruction Sheaf Structure . . . . .                                   | 33        |
| H.5      | Obstruction Formalization . . . . .                                        | 33        |
| H.6      | Theorem Schemas in Lean Style . . . . .                                    | 34        |
| H.7      | Algorithmic Reconstruction Procedure . . . . .                             | 35        |
| H.8      | Formalization Roadmap . . . . .                                            | 36        |

# 1 Introduction

The relationship between gravity and thermodynamics has attracted sustained theoretical attention since Bekenstein and Hawking established that black holes carry entropy proportional to horizon area. Subsequent programs — Jacobson’s derivation of the Einstein equations from thermodynamic identities [4, 5], Verlinde’s entropic gravity [2, 3], and more recently transactional entropic gravity [1] — have attempted to establish that gravitational dynamics emerge from statistical or information-theoretic structure rather than constituting a fundamental interaction. Padmanabhan has developed related thermodynamic perspectives on gravity [6], and the connection between entanglement structure and spacetime geometry has been explored through holographic frameworks [11, 12].

These programs share a common derivation order: physical exchange processes, thermodynamic bookkeeping, or entropic flow are treated as primitive, and spacetime geometry is reconstructed from them. The present work inverts this order. We treat *admissibility geometry* as primary and interpret persistent physical exchange as a stabilized reconstruction mode within that geometry. Emitters, absorbers, particles, and force laws are not inputs to the framework but outputs — roles stabilized by recursive cross-scale coherence of the coupled field  $X = (\Phi, \mathbf{v}, S)$  within its reconstruction sheaf [13, 15].

This inversion has a specific consequence for the MOND anomaly [7, 9]. Existing programs explain the transition scale  $a_0 \sim cH_0$  by finding thermodynamic narratives that reproduce it. The present framework instead asks: what geometric property of the reconstruction sheaf would make such a scale *structurally necessary*? The answer we propose — an admissibility radius  $R_A$  beyond which local reconstruction requires cosmological gluing data — converts a phenomenological constant into a constraint on the field theory itself.

## Epistemic Registers

The paper operates simultaneously across four epistemic registers which must be carefully distinguished. Certain structures are **definitional**, specifying the reconstruction geometry and admissibility framework. Others are **conjectural**, proposing relationships between admissibility structure and cosmological scaling. Still others are **derived implications** under specified assumptions. Finally, several results remain **open proof obligations** generated by the framework itself. We make these distinctions syntactically explicit throughout — using the theorem environments *Definition*, *Conjecture*, *Theorem-Schema*, *Heuristic*, and *Open Proof Obligation* — in order to avoid conflating motivation, formal consequence, and unresolved structure, a conflation that has weakened much of the emergent gravity literature.

## Structure of the Paper

Section 2 develops the admissibility geometry and sheaf structure. Section 3 introduces the admissibility phase transition and states the central conjecture regarding  $R_A$ . Section 4 treats entropy species as regime projections rather than identified primitives. Section 5

reinterprets dark matter phenomenology as reconstruction obstruction. Section 6 compares the framework with neighboring programs. Section 7 consolidates contributions and open obligations. Appendices A–H develop the formal machinery, including sheaf structure, coarse-graining flow, candidate field equations, obstruction cohomology, renormalization analogy, observational targets, locality clarification, and formal implementation via BNF grammar and Lean-style schemas.

## 2 Admissibility Geometry and the Reconstruction Sheaf

### 2.1 The Coupled Field

The fundamental object of the RSVP framework is the coupled field

$$X = (\Phi, \mathbf{v}, S), \quad (1)$$

where  $\Phi$  is a scalar field encoding the *accessibility geometry of continuation space* — the structure governing which local field trajectories admit coherent extension into persistent reconstruction modes;  $\mathbf{v}$  is a vector field encoding directed transport and coherence flow; and  $S$  is a scalar entropy-dispersion field measuring the spread of admissible reconstruction trajectories.

These three components are dynamically coupled. The scalar field  $\Phi$  governs which continuations remain accessible;  $\mathbf{v}$  tracks preferred directions of coherent continuation; and  $S$  measures how tightly reconstruction trajectories concentrate near stable modes. A configuration with high  $S$  relative to  $\Phi$  has many admissible continuations and correspondingly weak reconstruction stability. Low  $S/\Phi$  concentrates trajectories near persistent modes — the regime in which particles, clocks, and force-mediating exchanges stabilize as recognizable roles.

**The control parameter.** The dimensionless ratio

$$\chi = \frac{S}{\Phi} \quad (2)$$

is the central control parameter of the framework. When  $\chi \ll 1$ , entropy-dispersion is small relative to continuation accessibility, local reconstruction is strongly factorizable, and effective dynamics reduce approximately to Newtonian behavior. When  $\chi \sim 1$ , dispersion becomes comparable to accessibility, local factorization breaks down, and reconstruction requires boundary data from larger scales to maintain global stability. This ratio governs simultaneously: local factorization stability, the admissibility phase transition, entropy-channel convergence, and obstruction growth dynamics. Its ubiquity across all four phenomena is evidence of the framework’s internal coherence rather than coincidence.

## 2.2 Admissible Local Sections

**Definition 2.1** (Admissible local section). A local section of  $X$  over a region  $U$  is *admissible* if reconstruction trajectories originating within  $U$  remain dynamically stable under bounded perturbation and admit coherent continuation under the restriction structure of the field cover.

Admissibility is a *persistence condition*: it asks not merely whether a configuration is locally self-consistent, but whether it can continue coherently into neighboring regions and across scales. Three notions must be carefully distinguished.

- (i) **Local dynamical stability.** Trajectories within  $U$  remain bounded under perturbation. This is a condition on the field equations restricted to  $U$  alone, making no reference to neighboring regions.
- (ii) **Global admissibility.** Locally stable sections admit coherent extension across the full reconstruction cover without generating contradictions at overlapping boundaries. A section may be locally stable while failing global admissibility if its continuation structure conflicts with that of neighboring sections.
- (iii) **Reconstruction persistence.** Globally admissible extensions remain stable under renormalization of the reconstruction cover across scales — a multiscale stability condition rather than merely a temporal one. A section may be globally admissible at one scale while failing persistence at another if  $S$ -dispersion grows faster than the coherence structure of  $\mathbf{v}$  can absorb under coarse-graining.

The distinction between local stability and global admissibility is precisely what separates RSVP from transactional approaches [1]. Transactional completion guarantees local interaction closure — offer and confirmation waves match at a spacetime event. RSVP admissibility requires persistent global extension compatibility across the entire reconstruction cover. A transaction can complete locally while generating globally obstructed reconstruction dynamics; RSVP classifies such processes as inadmissible or metastable regardless of their local consistency.

## 2.3 The Reconstruction Cover

**Definition 2.2** (Admissibility neighborhood). An *admissibility neighborhood*  $U_\alpha$  is a region over which reconstruction remains approximately factorizable, meaning the admissible section over  $U_\alpha$  can be specified without reference to data outside  $U_\alpha$  up to corrections of order  $\frac{S}{\Phi}$ .

We introduce a cover  $\mathcal{U} = \{U_\alpha\}$  of the field domain where each  $U_\alpha$  is an admissibility neighborhood. Regions where  $S/\Phi$  is small generate large, strongly factorizable neighborhoods. Regions where  $S/\Phi$  approaches unity generate small, weakly factorizable neighborhoods whose sections increasingly require external boundary data for stable continuation.



*Remark 2.3.* The reconstruction cover is a *derived* object of the field geometry, not a background against which the field is defined. Classical spacetime structure, to the extent it appears, is a stable regime of the reconstruction cover rather than its presupposition. This is the formal content of RSVP’s departure from ordinary field theory: integrating out short-scale reconstruction structure changes not merely effective couplings but the factorization structure of admissible reconstruction itself [18, 19].

## 2.4 Two Types of Obstruction

Obstruction occurs when locally admissible sections fail to admit globally stable extension across overlapping neighborhoods. Obstruction is not metaphorical: it is a literal failure of global compatibility in the reconstruction geometry [16, 17]. We distinguish two structurally distinct types whose physical consequences differ and whose conflation would obscure both the entropy analysis of Section 4 and the dark matter reinterpretation of Section 5.

**Definition 2.4** (Compatibility obstruction). Let  $s_\alpha \in \mathcal{F}(U_\alpha)$  and  $s_\beta \in \mathcal{F}(U_\beta)$  be locally admissible sections over overlapping neighborhoods. A *compatibility obstruction* exists if

$$\rho_{\alpha\beta}(s_\alpha) \neq \rho_{\beta\alpha}(s_\beta)$$

on  $U_\alpha \cap U_\beta$ . The sections specify incompatible continuation structures on their shared domain and cannot be globally glued.

**Definition 2.5** (Dispersion obstruction). A pair of sections suffers *dispersion obstruction* if they agree formally on overlapping domains —  $\rho_{\alpha\beta}(s_\alpha) = \rho_{\beta\alpha}(s_\beta)$  — but cannot maintain stable global extension because  $S$  grows faster than the coherence propagation capacity of  $\mathbf{v}$  across the cover. Formally compatible sections therefore fail reconstruction persistence:  $\mathcal{P}[s] \rightarrow 0$  under scale extension.

*Remark 2.6.* Not all globally inconsistent dynamics arise from local disagreement. Some arise from uncontrolled admissibility growth under extension. Compatibility obstruction manifests as irregular, patchy anomalies without clear scaling structure. Dispersion obstruction manifests as smooth large-scale departure from locally stable behavior — the observational signature more characteristic of galactic rotation anomalies. The two obstruction types produce distinct signatures in the entropy structure of the field (Section 4) and distinct phenomenological predictions (Appendix F).

## 2.5 The Reconstruction Sheaf

The objects defined above constitute the data of a *reconstruction sheaf*  $\mathcal{F}$  over the field domain [13, 14]:  $\mathcal{F}$  assigns to each  $U_\alpha$  the space  $\mathcal{F}(U_\alpha)$  of admissible sections of  $X$  over  $U_\alpha$ , together with restriction maps

$$\rho_{\alpha\beta}: \mathcal{F}(U_\alpha) \longrightarrow \mathcal{F}(U_\alpha \cap U_\beta)$$

satisfying identity preservation, compatibility under composition, and the locality and gluing axioms. Global sections of  $\mathcal{F}$  correspond to globally admissible reconstruction modes.

*Remark 2.7* (Admissibility nonlocality). Admissibility nonlocality — when it occurs — is nonlocality in the indexing and coherence structure of  $\mathcal{F}$ , not in the stalk-local field dynamics. The stalks remain local field values propagating according to local field equations. What becomes nonlocal is the coherence condition on their assembly into globally stable sections. This is not a superluminal signaling proposal: nonlocality appears in the geometry of assembly rather than in microscopic propagation. See Appendix G for a precise separation of these notions.

## 2.6 Bridge to Section 3

Once admissibility is understood as a global extension condition rather than a local interaction condition, it becomes natural for reconstruction dynamics to transition between qualitatively distinct regimes as  $\chi$  varies. In strongly factorizable regimes ( $\chi \ll 1$ ), local sections extend to global sections without reference to cosmological boundary data. As  $\chi$  approaches unity, sections begin requiring cosmological gluing data for stable extension. The existence of a characteristic scale — the admissibility radius  $R_A$  — at which this transition occurs is proposed as a consequence of the field dynamics of  $X$ .

## 3 The Admissibility Phase Transition

The appearance of the MOND acceleration scale [7, 10]

$$a_0 \sim cH_0 \tag{3}$$

suggests that galactic dynamics are sensitive not merely to local mass distributions but to the relation between local reconstruction stability and cosmological admissibility structure. The coupling of galactic rotation to cosmological structure is difficult to interpret inside purely local Newtonian dynamics — it strongly suggests the relevant object is not local force accumulation alone but global admissibility structure across scales. Within RSVP, we interpret this transition not as a modification of force law but as a *breakdown of local factorization in the reconstruction cover*, a geometric event controlled by  $\chi = S/\Phi$ .

### 3.1 The Factorization Transition

In regimes where  $\chi \ll 1$ , local admissibility neighborhoods remain strongly factorizable. Reconstruction trajectories admit stable extension independently of cosmological boundary data, effective dynamics reduce approximately to Newtonian behavior, and the three entropy channels converge under coarse-graining (Section 4).

As

$$\chi = \frac{S}{\Phi} \longrightarrow 1, \tag{4}$$

dispersion becomes comparable to continuation accessibility. Local sections cease to remain scale-independent under extension. Admissibility neighborhoods lose strong factorization. Reconstruction increasingly requires cosmological gluing data to maintain persistence across the cover. Dispersion obstruction begins to dominate compatibility structure: even formally compatible sections fail reconstruction persistence.

**Definition 3.1** (Admissibility phase transition). The *admissibility phase transition* is the qualitative change in reconstruction regime occurring when  $\chi \rightarrow 1$ : the transition from locally dominated factorizable reconstruction to cosmologically coupled reconstruction requiring global boundary data for stable extension.

This is not a transition in the force law acting on matter, nor a transition in spacetime geometry in the classical sense. It is a transition in the factorization structure of admissible reconstruction — a change in what kind of boundary data local sections require for globally stable continuation. In transactional entropic gravity [1], the force law is modified by thermodynamic corrections applied to an otherwise standard local framework. In RSVP, the effective dynamics change because the geometric object governing which continuations are admissible changes.

### 3.2 The Admissibility Radius

**Definition 3.2** (Admissibility radius). The *admissibility radius*  $R_A$  is the largest characteristic scale over which admissible local sections remain approximately factorizable under renormalization of the reconstruction cover. Equivalently,  $R_A$  marks the onset of dispersion-dominated reconstruction dynamics for which stable continuation requires cosmological extension data: a local section at scale  $r$  requires cosmological gluing data if and only if its restriction to neighborhoods approaching  $R_A$  fails to admit stable extension through the sheaf’s restriction maps alone — that is, if  $\chi$  evaluated at that scale approaches unity under the coarse-graining flow.

$R_A$  is a derived quantity of the coupled dynamics of  $\Phi$  and  $S$ , not an external parameter. Different field configurations will generically produce different admissibility radii.

### 3.3 Newtonian and Cosmologically Coupled Regimes

**Newtonian regime** ( $r \ll R_A$ ,  $\chi \ll 1$ ). Local reconstruction is strongly factorizable. Admissible sections extend stably without cosmological boundary data. Dispersion obstruction remains bounded under scale extension. Effective gravitational dynamics reduce to Newtonian behavior as a consequence of local reconstruction stability rather than as a fundamental force law [2]. Newtonian gravity therefore appears as an effective low-dispersion reconstruction theory, analogous to the emergence of hydrodynamics from microscopic statistical systems [19].

**Cosmologically coupled regime** ( $r \sim R_A$ ,  $\chi \sim 1$ ). Local factorization breaks down. Admissible sections require cosmological gluing data. Dispersion obstruction dominates.

Reconstruction persistence fails for configurations that remain locally consistent. Effective dynamics depart from Newtonian behavior not because the force law changes, but because the admissible continuation set changes.

### 3.4 The Central Conjecture

**Conjecture 3.3** (MOND scaling from admissibility radius). *The weak-gradient RSVP field equations determine an asymptotic admissibility radius satisfying*

$$R_A \propto \frac{c}{H_0} \quad (5)$$

*in low-density regimes where  $\chi$  approaches unity smoothly. The corresponding admissibility acceleration scale is then*

$$a_A \sim \frac{c^2}{R_A} \sim cH_0, \quad (6)$$

*yielding MOND-like phenomenology as a structural consequence of reconstruction geometry.*

**Heuristic 3.4.** The structural plausibility of Conjecture 3.3 rests on the following observation. The Hubble radius  $c/H_0$  is the characteristic scale of cosmological causal accessibility. If  $R_A$  is the admissibility analog of causal accessibility, its asymptotic scaling with  $c/H_0$  would follow from the requirement that admissibility structure and causal structure coincide asymptotically in low-density regimes. This is a structural argument for the conjecture’s reasonableness, not a derivation.

The MOND scale  $a_0 \sim cH_0$  has been observed by several authors to couple galactic dynamics to cosmological structure in a way that resists purely local explanation [7, 9]. Within RSVP, this coupling emerges naturally from the admissibility framework: the transition scale is determined by the scale at which local reconstruction loses self-sufficiency. MOND is therefore demoted from foundational principle to phenomenological signature of a more general reconstruction transition.

### 3.5 Open Proof Obligation

*Open Proof Obligation 3.5.* Show that the coupled dynamics of  $\Phi$  and  $S$  in the weak-gradient limit uniquely determine  $R_A$  with asymptotic scaling  $R_A \propto c/H_0$ . This requires: (i) deriving the coarse-graining flow of  $\chi$  under renormalization of the reconstruction cover in the weak-gradient limit; (ii) identifying the fixed-point or threshold structure of this flow; and (iii) showing that the characteristic scale of the threshold is asymptotically proportional to  $c/H_0$ .

Conjecture 3.3 is falsifiable in three distinct senses: (1) if the RSVP field equations admit a unique  $R_A$  whose asymptotic scaling substantially departs from  $c/H_0$ , then MOND-like phenomenology must arise from a different mechanism; (2) if MOND-like behavior were observed in regimes where  $\chi$  remains well below unity, this would constitute evidence against the sheaf structure assumed here; and (3) if  $R_A$  can be derived but depends

sensitively on unconstrained parameters in the RSVP field equations, the conjecture is technically falsifiable but practically underdetermined — a third failure mode named explicitly.

### 3.6 Bridge to Section 4

The admissibility phase transition has consequences not only for effective dynamics but for the entropy structure of the field. In strongly factorizable regimes, the distinct entropy channels converge under coarse-graining because local reconstruction is approximately scale-independent. As  $\chi \rightarrow 1$ , this convergence fails. Section 4 formalizes this consequence, deriving entropy-channel behavior as a corollary of reconstruction geometry rather than as an independent thermodynamic hypothesis.

## 4 Entropy Species as Regime Projections

Most emergent gravity programs implicitly identify informational, thermodynamic, configurational, and gravitational entropy as interchangeable quantities, treating their equivalence as a foundational step enabling the derivation of gravitational dynamics [2, 1]. The information-thermodynamic identification remains debated [20], and its use as an axiom in precisely the regimes where it is least reliable constitutes a significant structural weakness. Within RSVP, these entropy channels are instead treated as distinct projections of admissibility dispersion under different reconstruction covers and coarse-grainings [21]. Their convergence is not assumed axiomatically but emerges only within specific reconstruction regimes. This converts what is typically an axiom into a prediction.

### 4.1 Entropy Channels as Projections

Given a reconstruction cover  $\mathcal{U} = \{U_\alpha\}$ , different entropy species correspond to distinct coarse-grained projections of admissibility dispersion across  $X = (\Phi, \mathbf{v}, S)$ . We formalize these as projection functors

$$\Pi_{\text{therm}}, \quad \Pi_{\text{info}}, \quad \Pi_{\text{grav}}, \quad \Pi_{\text{conf}}$$

mapping admissibility structures into entropy observables associated with distinct reconstruction regimes and coarse-graining procedures.

Informational entropy, represented by  $\Pi_{\text{info}}$ , measures uncertainty in continuation accessibility under local reconstruction. It quantifies the spread of admissible continuations available to a local section of  $\Phi$  and therefore tracks the local geometry of admissible continuation space.

Configurational entropy, represented by  $\Pi_{\text{conf}}$ , measures admissible arrangement multiplicity across neighboring sections of the reconstruction cover. It characterizes the number of globally compatible assemblies obtainable from locally admissible reconstruction data.

Thermodynamic entropy, represented by  $\Pi_{\text{therm}}$ , measures the coarse-grained dispersion of reconstruction trajectories under persistent exchange processes generated by the coherence-flow field  $\mathbf{v}$ . It corresponds to the effective trajectory-volume growth observed under macroscopic reconstruction flow.

Gravitational entropy, represented by  $\Pi_{\text{grav}}$ , measures dispersion associated with large-scale reconstruction curvature and admissibility obstruction. Its value depends not only on local continuation accessibility but on the obstruction structure governing global extension of admissible sections across the reconstruction sheaf.

None of these entropy channels is treated as fundamentally privileged within the RSVP framework. They correspond instead to distinct projection structures defined over the same underlying admissibility geometry. Their relationships are therefore regime-dependent rather than identity-relations. In strongly factorizable regimes, the projection functors become approximately equivalent under admissible coarse-graining. Near the admissibility transition, however, the projections separate as dispersion obstruction grows and local reconstruction loses scale-independence.

## 4.2 Entropy Convergence Under Strong Factorization

**Theorem-Schema 4.1** (Entropy convergence). In reconstruction regimes satisfying  $\chi \ll 1$  and admitting bounded dispersion obstruction under renormalization of the reconstruction cover, entropy channels converge to order  $\varepsilon$  under admissible coarse-graining transformations:

$$\Pi_{\text{therm}} \simeq \Pi_{\text{info}} \simeq \Pi_{\text{grav}} \simeq \Pi_{\text{conf}} \quad (\chi \ll 1).$$

Equivalently, informational, configurational, thermodynamic, and gravitational entropy measures become approximately equivalent up to corrections of order  $\chi$  and the dispersion obstruction growth rate.

When  $\chi \ll 1$ , local reconstruction is strongly factorizable and approximately scale-independent. Different coarse-graining choices produce approximately the same result because admissibility dispersion is tightly concentrated near stable reconstruction modes — the cover choice becomes approximately irrelevant, and the distinct entropy projections converge. The entropy-identification moves made by emergent gravity programs are approximately valid in this regime, but only in this regime. The present framework derives the conditions under which they hold and identifies them as regime-specific approximations.

## 4.3 Entropy Divergence Under Dispersion Obstruction

As  $\chi \rightarrow 1$ , dispersion obstruction begins to dominate. Different coarse-graining choices produce systematically different results. Informational entropy, tracking local continuation accessibility, may remain bounded while thermodynamic entropy grows. Gravitational entropy, tracking obstruction contributions, may diverge from configurational entropy as

global extension failures accumulate. Functorially:

$$\Pi_{\text{therm}} \not\approx \Pi_{\text{grav}} \quad (\chi \rightarrow 1).$$

*Remark 4.2.* Persistent divergence between entropy species is not an annoying inconsistency to be resolved by a better identification. It is a physical signal: evidence of obstruction growth within the reconstruction geometry itself. When entropy channels diverge persistently in a physical system, the RSVP framework predicts this as a signature of the system operating near or beyond the admissibility phase transition.

#### 4.4 The Kastner–Schlatter Contrast

Kastner and Schlatter [1] identify Shannon information entropy with thermodynamic entropy as a foundational step, using this identification to drive the derivation of gravitational dynamics. Within RSVP, this identification is approximately valid when  $\chi \ll 1$  (Theorem-Schema 4.1) and fails as a foundational assumption precisely in the regimes of greatest physical interest: the weak-gradient, low-density regimes where MOND-like behavior appears and  $\chi$  approaches unity. Founding the derivation on entropy identification in those regimes imports an approximation as an axiom exactly where the approximation breaks down.

#### 4.5 Bridge to Section 5

The existence of persistent entropy-channel divergence without local dynamical inconsistency suggests that globally anomalous galactic behavior need not arise from missing mass distributions. Recall Remark 2.3: the reconstruction cover is a derived object of the field geometry, not a background against which the field is defined. Anomalies in the global extension structure of the cover are anomalies in the geometry itself, not in the matter content of a pre-given space.

### 5 Dark Matter as Reconstruction Obstruction

#### 5.1 The Observational Problem

The observational signatures attributed to dark matter are well established empirically: flat galactic rotation curves [10, 9], weak gravitational lensing excesses, cluster dynamics requiring additional gravitational sources, and large-scale structure formation rates exceeding baryonic predictions.

Standard cosmology interprets these anomalies as evidence for additional unseen mass distributions embedded within a fixed spacetime background. Within RSVP, the anomalies are instead interpreted as *signatures of obstruction in the global extension structure of the reconstruction cover*. As established in Remark 2.3: the reconstruction cover is a derived object of the field geometry, not a background against which the field is defined. Anomalies in the global extension structure of the cover are anomalies in the geometry itself.

## 5.2 Compatibility and Dispersion Obstruction in Galactic Dynamics

Compatibility obstruction (Definition 2.4) would manifest as local dynamical inconsistency — irregular, patchy anomalies without clear scaling structure. The observational signatures of galactic anomalies do not have this character. Galaxies are locally stable and internally coherent; their stellar and gas dynamics are well described by standard hydrodynamics and Newtonian gravity at small radii.

The anomaly appears in the *persistence* of those dynamics under large-scale reconstruction. This is structurally closer to dispersion obstruction (Definition 2.5). Local sections remain formally compatible across neighboring regions; the failure is not disagreement but persistence failure under scale extension: as  $\chi$  approaches unity near  $R_A$ , local reconstruction dynamics lose scale-independence and can no longer be extended stably without cosmological boundary data. The point is not that galaxies behave inconsistently. The point is that their local reconstruction dynamics fail to remain self-sufficient under cosmological extension.

## 5.3 Rotation Curves as Persistence Failure

In strongly factorizable regimes ( $\chi \ll 1$ ), orbital dynamics are governed predominantly by local continuation accessibility, and the admissible trajectory set is closed under local extension — the regime in which Newtonian dynamics holds as an emergent approximation. As dispersion obstruction grows and  $\chi$  approaches unity near  $R_A$ , local continuation sets cease to remain closed under scale extension. The admissible trajectory structure at galactic radii approaching  $R_A$  is no longer determined by local field data alone — it requires cosmological gluing data for stable continuation. The effective admissible trajectory set therefore departs from Newtonian closure even in the absence of additional local mass.

The transition criterion  $\chi \rightarrow 1$  marks the onset of this regime — the same criterion governing the admissibility phase transition (Section 3) and the entropy-channel divergence (Section 4). The unification of all three phenomena under the same control parameter is the framework’s strongest structural feature.

*Observation 5.1. The anomaly is a geometric phase transition.* Flat rotation curves, MOND-like scaling, and entropy-channel divergence are manifestations of the same underlying transition in the factorization structure of admissible reconstruction, controlled by  $\chi \rightarrow 1$  near  $R_A$ .

## 5.4 Weak Lensing and Extension Geometry

Within RSVP, photon propagation probes the continuation accessibility geometry of  $\Phi$  rather than merely local mass density. In strongly factorizable regimes, continuation accessibility tracks local mass distributions closely and lensing reduces approximately to standard gravitational lensing. In regimes approaching the admissibility phase transition, the continuation accessibility geometry becomes influenced by obstruction structure in the reconstruction cover.



*Open Proof Obligation 5.2.* Compute the effective lensing potential from the obstruction curvature of  $\mathcal{F}$  in the weak-gradient limit and compare with observed convergence maps, including Bullet Cluster phenomenology.

The framework identifies this as a possible reinterpretation of lensing anomalies in terms of obstruction curvature. Whether the resulting phenomenology quantitatively reproduces observed lensing profiles remains an open problem. The Bullet Cluster remains a decisive empirical challenge: the framework must reproduce the observed lensing structure through obstruction curvature of the reconstruction geometry. The restraint here is deliberate; this target is stated honestly rather than rhetorically dismissed.

## 5.5 Entropy Divergence as Observational Signal

Regions dominated by dispersion obstruction should exhibit increasing divergence between informational, thermodynamic, configurational, and gravitational entropy measures. Gravitational entropy, tracking obstruction contributions, should depart from thermodynamic entropy of local baryonic processes in exactly the regimes where rotation curve anomalies appear. Standard  $\Lambda$ CDM does not predict entropy-channel divergence as a geometric signal; RSVP predicts that such divergence is a structural signature of the admissibility phase transition.

*Open Proof Obligation 5.3.* Derive explicit expressions for the gravitational entropy contribution of reconstruction obstruction in the weak-gradient RSVP limit and determine whether predicted divergence onset correlates with observed rotation curve transition radii.

## 5.6 Situating the Reinterpretation

The RSVP reinterpretation differs structurally from all three major competing programs. Particle dark matter models preserve local factorization while adding unseen sources. Entropic gravity programs [3, 6] preserve entropy identification while modifying effective dynamics. MOND and TeVeS [7, 8] modify the force law through fitted interpolation functions. RSVP instead interprets galactic anomalies as signatures of breakdown in scale-independent reconstruction itself — the onset of dispersion-dominated extension dynamics controlled by  $\chi$  approaching unity near  $R_A$  (Observation 5.1).

# 6 Comparison with Neighboring Programs

## 6.1 Organizing Principle

The most useful axis for comparing competing approaches is derivation architecture: what object is treated as primitive, what undergoes transition, what role entropy plays, and what counts as an anomaly.

| Framework               | Primitive              | Anomaly            | Entropy           | Locality       |
|-------------------------|------------------------|--------------------|-------------------|----------------|
| $\Lambda$ CDM           | Matter source          | Missing source     | Identified        | Local field    |
| MOND/TeV <span>S</span> | Modified force         | Wrong dynamics     | Secondary         | Modified local |
| Verlinde                | Entropy equiv.         | Incomplete thermo. | Axiom             | Holographic    |
| Trans. entropic         | Exchange completion    | Incomplete thermo. | Interchangeable   | Local exchange |
| RSVP                    | Admissibility geometry | Phase transition   | Regime projection | Sheaf-indexed  |

Table 1: Comparative taxonomy organized by derivation architecture.

## 6.2 Standard Cosmology ( $\Lambda$ CDM)

$\Lambda$ CDM treats matter source distributions within a fixed pseudo-Riemannian background as primitive. What RSVP interprets as obstruction structure in a derived cover,  $\Lambda$ CDM interprets as source deficits within a fixed background. These are not empirically equivalent restatements: they predict different entropy-channel signatures, different lensing phenomenology in obstruction-dominated regimes, and different structure formation dynamics. RSVP constrains the anomaly structure through  $\chi$  and  $R_A$ ; the phenomenology is not freely adjustable but is derived from field dynamics.

## 6.3 MOND and TeVeS

MOND [7] reproduces flat rotation curves and the baryonic Tully-Fisher relation [10] with remarkable precision using a single additional parameter. Within the present framework, MOND is demoted from foundational principle to phenomenological signature. The MOND acceleration scale  $a_0 \sim cH_0$  is not a new fundamental constant but a derived geometric scale — the acceleration associated with the onset of cosmologically coupled reconstruction at  $R_A$ . MOND-like dynamics appear when  $\chi \rightarrow 1$  near  $R_A$ , not because the force law has been modified but because the admissible continuation set has changed.

The observed robustness of the baryonic Tully-Fisher relation [10] across galaxy types, which is difficult to understand in  $\Lambda$ CDM and built into MOND by construction, would in RSVP reflect universality of the admissibility phase transition under coarse-graining — an emergent property of the reconstruction geometry rather than a tuned interpolation (see Appendix E).

## 6.4 Verlinde’s Emergent Gravity

Verlinde [2, 3] begins with entropy identification and derives effective geometry. RSVP begins with admissibility geometry and derives the conditions under which entropy species approximately converge (Theorem-Schema 4.1). Verlinde’s framework inherits the entropy-identification problem [20] as a foundational commitment. RSVP converts this axiom into

a theorem-schema valid in strongly factorizable regimes and failing in precisely the regimes —  $\chi \rightarrow 1$  — where anomalous dynamics appear.

## 6.5 Transactional Entropic Gravity

Kastner and Schlatter [1] provide the most immediately adjacent comparison. Both programs reject fundamental spacetime; both treat geometric structure as emergent. The divergence is in derivation order and primitive object.

Transactional entropic gravity treats exchange process completion as primitive — space-time links are generated by completed transactions between irreducible emitters and absorbers. RSVP treats admissibility geometry as primitive — emitters, absorbers, and exchange processes are stabilized roles within the reconstruction sheaf.

Three structural consequences follow: (1) the transactional MOND derivation depends on entropy-species identification at specific steps; the RSVP conjecture depends on a single geometric condition ( $\chi \rightarrow 1$  at  $R_A$ ) controlling all phenomena simultaneously; (2) transactional completion is strictly weaker than sheaf-compatible global extension — a transaction can complete locally while generating globally inadmissible dynamics; (3) entropy-species interchangeability appears in RSVP as an approximate consequence of strongly factorizable reconstruction, valid in the Newtonian regime and failing in the MOND regime.

RSVP does not currently outperform transactional entropic gravity quantitatively. The advantage claimed is architectural: fewer primitive objects controlling more phenomena through a single transition structure, with entropy convergence as a derived consequence.

## 6.6 The Inversion as Theoretical Commitment

The organizing inversion — admissibility geometry generates exchange structure, not vice versa — reflects a specific theoretical bet about where the explanatory primitives of gravitational physics are located. If the bet is correct, existing phenomenology should emerge as special cases of reconstruction geometry. If wrong, the transactional program has the correct derivation order and RSVP’s sheaf structure is descriptive rather than explanatory. The paper makes the question precise and identifies the geometric objects on which it turns.

# 7 Conclusion

## 7.1 The Central Reinterpretation

The present work develops a reinterpretation of gravitational anomalies organized around a single foundational inversion. Existing emergent gravity programs treat physical exchange, thermodynamic bookkeeping, or information flow as primitive and derive spacetime geometry from them. RSVP treats admissibility geometry as primary and interprets persistent physical exchange as a stabilized reconstruction mode. Emitters, absorbers, particles, force laws, and thermodynamic entropy are not inputs to the framework but outputs — roles

stabilized by recursive cross-scale coherence of the coupled field  $X = (\Phi, \mathbf{v}, S)$  within its reconstruction sheaf.

The framework’s explanatory structure is organized by a small set of recurring mathematical objects: the coupled field  $X$ ; the control parameter  $\chi = S/\Phi$ ; the reconstruction cover  $\mathcal{U}$ ; the admissibility radius  $R_A$ ; the compatibility and dispersion obstruction classes; and the entropy projection functors  $\Pi_{\text{therm}}, \Pi_{\text{info}}, \Pi_{\text{grav}}, \Pi_{\text{conf}}$ . The control parameter  $\chi$  governs simultaneously: Newtonian gravitational stability, the MOND-like transition, entropy-channel convergence, and galactic obstruction phenomenology. This unification under a single order parameter is the strongest evidence of internal coherence.

## 7.2 What the Paper Claims and Does Not Claim

The paper does not present: a completed derivation of MOND from RSVP field equations; a quantitative alternative to  $\Lambda$ CDM; or a solved lensing model capable of reproducing Bullet Cluster phenomenology.

The paper does present: a reconstruction-geometric reinterpretation of gravitational anomalies organized by a stable conceptual hierarchy; formal definitions of admissibility geometry, reconstruction cover, and obstruction types; a constrained and explicitly falsifiable conjecture (Conjecture 3.3) with three named failure modes; Theorem-Schema 4.1 converting an axiom of neighboring programs into a derived consequence; and a comparative taxonomy organized by derivation architecture (Table 1).

The present work does not derive MOND or dark matter phenomenology from completed RSVP field equations; it identifies the geometric objects from which such derivations would have to proceed.

## 7.3 Open Problems as Evidence of Structural Coherence

Most speculative frameworks accumulate explanatory flexibility. The present framework instead generates constrained open problems whose resolution would confirm or falsify its central commitments:

- (1) **Admissibility radius derivation** (Obligation 3.5). Show that the weak-gradient RSVP equations uniquely determine  $R_A \propto c/H_0$ .
- (2) **Universality class structure**. Determine whether distinct admissibility geometries flow toward the same weak-gradient reconstruction behavior under coarse-graining, making MOND phenomenology geometrically necessary.
- (3) **Lensing derivation target** (Obligation 5.2). Compute the effective lensing potential from obstruction curvature of  $\mathcal{F}$  and compare with observed convergence maps.
- (4) **Entropy divergence signature** (Obligation 5.3). Identify observational contexts where gravitational entropy departs measurably from thermodynamic entropy of local baryonic processes.

- (5) **Obstruction flow analysis.** Determine whether compatibility and dispersion obstruction produce distinguishable galactic signatures varying systematically with galaxy morphology, surface brightness, or environment.

These problems are obligations generated by the framework’s own internal structure. Their existence as constrained derivation targets — rather than free parameters or post-hoc interpretive choices — is evidence that the framework has crossed the threshold where speculative architecture begins behaving as a research program. A framework that knows the shape of its own incompleteness is structurally more mature than one that knows only the shape of its successes.

## 7.4 Final Philosophical Position

If the RSVP program is correct, then gravity, inertia, dark matter phenomenology, and entropy equivalence are not independent physical ingredients assembled into a cosmological model. They are stable regimes and failure modes of reconstruction geometry — consequences of the factorization structure of admissible continuation under a derived cover whose topology is controlled by a single dimensionless parameter.

Whether this inversion reflects physical reality remains unresolved. The present work aims only to make the inversion mathematically explicit, empirically constrained, and structurally falsifiable.

## A Formal Structure of the Reconstruction Sheaf

This appendix develops the formal sheaf-theoretic structure underlying the reconstruction framework. The purpose is not to introduce new physical claims but to make explicit the mathematical objects already used operationally throughout Sections 2–5.

### A.1 Presheaf and Sheaf Structure

Let  $\mathcal{R}$  denote the reconstruction domain and  $\mathcal{U} = \{U_\alpha\}$  a cover by admissibility neighborhoods. The *reconstruction presheaf*  $\mathcal{F}$  assigns to each  $U_\alpha$  a set  $\mathcal{F}(U_\alpha)$  of admissible sections of  $X$  over  $U_\alpha$ , together with restriction maps

$$\rho_V^U: \mathcal{F}(U) \longrightarrow \mathcal{F}(V), \quad V \subseteq U,$$

satisfying  $\rho_U^U = \text{id}$  and  $\rho_W^V \circ \rho_V^U = \rho_W^U$  for  $W \subseteq V \subseteq U$ .

$\mathcal{F}$  is a *sheaf* if it satisfies the following for every open cover  $\{U_i\}$  of any  $U$ :

- (i) *Locality*: if  $\rho_{U_i}^U(s) = \rho_{U_i}^U(t)$  for all  $i$ , then  $s = t$ .
- (ii) *Gluing*: if  $\rho_{U_i \cap U_j}^{U_i}(s_i) = \rho_{U_i \cap U_j}^{U_j}(s_j)$  for all  $i, j$ , then there exists  $s \in \mathcal{F}(U)$  with  $\rho_{U_i}^U(s) = s_i$ .

Compatibility obstruction (Definition 2.4) corresponds to failure of condition (ii): sections cannot be glued because they disagree on overlaps. Dispersion obstruction (Definition 2.5) is subtler: sections satisfy (ii) formally but the glued section fails reconstruction persistence under coarse-graining because admissibility dispersion increases faster than coherence propagation.

## A.2 Admissible Sections and Persistence

An admissible local section is not merely a field restriction but a structured pair consisting of: a local restriction of the coupled field

$$X = (\Phi, \mathbf{v}, S)$$

to an admissibility neighborhood  $U_\alpha$ , together with a continuation structure specifying the admissible extension behavior of reconstruction trajectories originating within  $U_\alpha$ .

Formally, a section

$$s_\alpha \in \mathcal{F}(U_\alpha)$$

is admissible only if two conditions are simultaneously satisfied. First, local perturbations of the section must preserve bounded continuation behavior, so that small deformations of initial reconstruction data do not generate uncontrolled trajectory divergence. Second, admissible extensions of the section must remain stable under renormalization of the reconstruction cover. The section must therefore persist not only locally but under scale-dependent restructuring of admissibility neighborhoods induced by coarse-graining flow.

This second condition distinguishes reconstruction persistence from ordinary local field consistency. A section may satisfy all local field equations and remain formally compatible on overlaps while nevertheless failing persistence under extension because admissibility dispersion grows faster than coherence propagation under cover renormalization.

A global reconstruction mode corresponds to a global section

$$s \in \mathcal{F}(\mathcal{R})$$

whose restrictions agree coherently across the full reconstruction cover and whose continuation structure remains stable under admissible coarse-graining flow. Persistence is therefore a multiscale extension property of the reconstruction sheaf rather than merely a statement of local dynamical consistency.

## A.3 Persistence Functional

We define a *persistence functional*

$$\mathcal{P}[s] \in [0, 1]$$

measuring stability of admissible extensions under renormalization of the reconstruction cover. A section satisfies reconstruction persistence if  $\mathcal{P}[s]$  remains bounded away from zero under scale extension. Dispersion obstruction occurs when  $\mathcal{P}[s] \rightarrow 0$  under coarse-graining, even for sections satisfying the standard gluing conditions.

#### A.4 Dynamical Cover and Gluing

Unlike ordinary differential-geometric frameworks in which the manifold is fixed and fields evolve upon it, the RSVP framework treats the effective reconstruction topology as dependent upon admissibility flow. In RSVP, gluing conditions are dynamically weighted by admissibility dispersion and continuation accessibility rather than determined solely by overlap consistency. The resulting structure combines: local field dynamics, scale-dependent admissibility flow, and global extension stability into a single reconstruction structure.

The following commutative diagram summarizes the restriction structure for a triple cover:

$$\begin{array}{ccc} \mathcal{F}(U_\alpha) & \xrightarrow{\rho_{\alpha\beta}} & \mathcal{F}(U_\alpha \cap U_\beta) \\ & \searrow \rho_{\alpha\gamma} & \downarrow \rho_{\alpha\beta,\gamma} \\ & & \mathcal{F}(U_\alpha \cap U_\beta \cap U_\gamma) \end{array}$$

Consistency of this diagram for all triples is the cocycle condition whose failure generates compatibility obstructions classified by  $\check{H}^1(\mathcal{U}, \mathcal{F})$ .

## B Coarse-Graining Flow and the Admissibility Order Parameter

The main text introduced  $\chi = S/\Phi$  as the central control parameter. This appendix develops a schematic coarse-graining framework and clarifies the sense in which the admissibility phase transition may possess universality structure analogous to ordinary critical phenomena [18, 19].

### B.1 Scale-Dependent Fields and Order Parameter

Let  $\mathcal{U}_\ell$  denote the reconstruction cover at coarse-graining scale  $\ell$ . Integrating out short-scale reconstruction modes modifies both continuation accessibility and admissibility dispersion:

$$\Phi \longrightarrow \Phi_\ell, \quad S \longrightarrow S_\ell.$$

The scale-dependent order parameter is

$$\chi_\ell = \frac{S_\ell}{\Phi_\ell}.$$

The Newtonian regime ( $\chi_\ell \ll 1$ ) corresponds to strongly factorizable reconstruction; the admissibility transition occurs when  $\chi_\ell \rightarrow 1$ .

## B.2 Admissibility Beta-Function

The coarse-graining flow of the reconstruction geometry may be represented schematically by the scale evolution equation

$$\frac{d\chi_\ell}{d\ln \ell} = \beta(\chi_\ell), \quad (7)$$

where

$$\chi_\ell = \frac{S}{\Phi} \Big|_\ell$$

denotes the scale-dependent control parameter evaluated at coarse-graining scale  $\ell$ , and  $\beta$  is an *admissibility beta-function* governing the renormalization flow of reconstruction stability.

The function  $\beta$  is not specified phenomenologically but constrained by the structural requirements of admissible reconstruction developed throughout the main text. In strongly factorizable regimes,

$$\chi \ll 1,$$

the framework requires

$$\beta(\chi) < 0,$$

so that admissibility dispersion decreases under coarse-graining flow and local reconstruction remains stable under scale extension. This corresponds to the Newtonian regime in which reconstruction neighborhoods retain approximate factorization and admissible sections persist without cosmological gluing data.

As  $\chi$  approaches unity, the flow enters a crossover region in which dispersion obstruction begins competing with coherence propagation. The qualitative structure of admissible extension changes: local sections cease to remain scale-independent, and persistence under renormalization becomes increasingly sensitive to global reconstruction geometry. The admissibility phase transition introduced in the main text corresponds to this change in flow structure near

$$\chi \sim 1.$$

Beyond the transition, the framework admits two structurally distinct possibilities. Either the flow approaches a new large-scale stable reconstruction regime governed by cosmologically coupled admissibility structure, or local factorization ceases to remain dynamically stable altogether and reconstruction persistence fails beyond finite scale. Which possibility is realized depends on the weak-gradient dynamics of the RSVP field equations and constitutes part of the open derivation program identified in Obligation 3.5.

The admissibility beta-function therefore plays a role analogous to renormalization group flow equations in statistical field theory. It governs not merely numerical coupling evolution but qualitative transitions in the factorization structure of admissible reconstruc-



tion itself.

### B.3 Universality Classes

Suppose multiple microscopic RSVP field dynamics produce distinct local reconstruction behavior while nevertheless generating the same asymptotic coarse-graining flow for  $\chi_\ell$ . Then the weak-gradient galactic regime would exhibit universal phenomenology independent of microscopic reconstruction details — analogous to universality of critical exponents in condensed matter systems [19]. This possibility is particularly important for understanding the empirical robustness of MOND-like scaling relations [10, 9] across diverse galactic systems. Within RSVP, such robustness would emerge as a universality property of admissibility flow near the reconstruction transition rather than a tuned interpolation.

### B.4 Admissibility Radius from Flow

The admissibility radius  $R_A$  is the characteristic scale at which  $\chi_\ell \rightarrow 1$  under the coarse-graining flow. Obligation 3.5 is therefore equivalent to showing that flow equation (7), derived from the weak-gradient RSVP field equations, admits a threshold at scale  $\ell^*$  satisfying  $e^{\ell^*} \propto c/H_0$ .

## C Candidate Dynamical Equations for the RSVP Fields

This appendix sketches candidate dynamical equations for the coupled fields  $X = (\Phi, \mathbf{v}, S)$ . These equations are heuristic and incomplete — candidate structures consistent with the admissibility framework, not finalized physical laws. The purpose is architectural: to show that the reconstruction-geometric interpretation can plausibly be associated with a coherent field-dynamical system.

### C.1 Continuation Accessibility Field $\Phi$

A minimal admissibility-flow equation for  $\Phi$ :

$$\partial_t \Phi = D_\Phi \nabla^2 \Phi - \lambda_\Phi \Phi S + \gamma_\Phi \nabla \cdot \mathbf{v} + N_\Phi[\Phi, \mathbf{v}, S] \quad (8)$$

where  $D_\Phi$  is an accessibility diffusion coefficient,  $\lambda_\Phi$  governs suppression of continuation accessibility by dispersion growth ( $-\lambda_\Phi \Phi S$  captures the central geometric tension: increasing  $S$  suppresses stable  $\Phi$ ),  $\gamma_\Phi$  couples accessibility to coherence flow divergence, and  $N_\Phi$  denotes higher-order nonlinear terms.

### C.2 Coherence Flow Field $\mathbf{v}$

$$\partial_t \mathbf{v} = D_v \nabla^2 \mathbf{v} - (\mathbf{v} \cdot \nabla) \mathbf{v} + \alpha_v \nabla \Phi - \lambda_v S \mathbf{v} + N_v[\Phi, \mathbf{v}, S] \quad (9)$$

The gradient term  $\alpha_v \nabla \Phi$  drives coherence flow toward high continuation accessibility. The suppression term  $-\lambda_v S \mathbf{v}$  models degradation of coherent transport under increasing dis-

persion. The nonlinear transport term resembles fluid advection but its interpretation is geometric: reconstruction coherence propagates directionally through admissibility structure.

### C.3 Entropy-Dispersion Field $S$

$$\partial_t S = D_S \nabla^2 S + \sigma_S |\nabla \Phi|^2 - \mu_S S \Phi - \nabla \cdot (S \mathbf{v}) + N_S [\Phi, \mathbf{v}, S] \quad (10)$$

The source term  $\sigma_S |\nabla \Phi|^2$  generates dispersion through accessibility curvature gradients. The suppression term  $-\mu_S S \Phi$  stabilizes dispersion in regions of high continuation accessibility. The advection term  $-\nabla \cdot (S \mathbf{v})$  transports dispersion through coherence flow.

### C.4 Control Parameter Dynamics

Together, equations (8)–(10) determine the evolution of the control parameter:

$$\partial_t \chi = \frac{\Phi \partial_t S - S \partial_t \Phi}{\Phi^2}.$$

Equation (10) is of particular interest: the source term  $\sigma_S (1 - S/\Phi) S = \sigma_S (1 - \chi) \chi \Phi$  drives  $\chi$  toward unity when  $\chi < 1$  and away when  $\chi > 1$ , producing the admissibility transition at  $\chi = 1$  as a dynamical fixed point.

### C.5 Epistemic Status

These equations are schematic. Many alternative systems may generate the same admissibility phenomenology; if the universality hypothesis of Appendix B is correct, weak-gradient reconstruction behavior may be largely insensitive to microscopic dynamical details. A completed RSVP field theory would require: a variational principle, covariant generalization, explicit reconstruction observables, well-posed initial value structure, and quantitative comparison with data.

## D Obstruction Cohomology and Global Reconstruction Failure

This appendix sketches a cohomological interpretation of reconstruction obstruction [16, 17, 14]. The purpose is to demonstrate mathematical directionality, not claim a finished cohomological theory.

### D.1 Čech Cohomology of the Reconstruction Cover

Let  $\check{C}^\bullet(\mathcal{U}, \mathcal{F})$  denote the Čech cochain complex associated with  $\mathcal{U}$  and  $\mathcal{F}$ . A family of local sections  $\{s_\alpha\}$  defines a 0-cochain. A compatibility obstruction corresponds to a non-trivial 1-cocycle  $g_{\alpha\beta} \in \check{C}^1(\mathcal{U}, \mathcal{F})$  measuring the failure of sections to agree on overlaps. Globally admissible reconstruction requires trivialization of this cocycle:

**Definition D.1** (Compatibility obstruction class). A *compatibility obstruction class* is a nontrivial element of

$$\check{H}^1(\mathcal{U}, \mathcal{F}),$$

corresponding to the impossibility of globally gluing locally admissible sections. Vanishing  $\check{H}^1$  indicates that all compatible local sections admit global extension without compatibility obstruction.

## D.2 Two Obstruction Notions

Within RSVP, obstruction possesses an additional dynamical feature absent from ordinary static sheaf theory: the reconstruction cover evolves under admissibility coarse-graining. This motivates two notions:

**Definition D.2** (Static obstruction). A *static obstruction* is an ordinary compatibility obstruction at a fixed reconstruction scale — a nontrivial class in  $\check{H}^1(\mathcal{U}_\ell, \mathcal{F})$  for some fixed  $\ell$ .

**Definition D.3** (Persistence obstruction). A *persistence obstruction* occurs when an obstruction class fails to be trivialized under coarse-graining of the reconstruction cover, even if it is trivializable at some initial scale. Formally, there exists a coarse-graining flow  $\mathcal{R}_\lambda$  such that the induced map

$$\mathcal{R}_\lambda^*: \check{H}^1(\mathcal{U}_{\lambda_1}, \mathcal{F}) \longrightarrow \check{H}^1(\mathcal{U}_{\lambda_2}, \mathcal{F}), \quad \lambda_2 > \lambda_1,$$

does not preserve triviality.

## D.3 Scale-Dependent Obstruction Functional

Introduce a scale-dependent obstruction functional:

$$\Omega_\ell[s] \in \mathbb{R}_{\geq 0}$$

whose flow under coarse-graining satisfies:

$$\frac{d\Omega_\ell}{d \ln \ell} = \Gamma(\chi_\ell, \Omega_\ell) \tag{11}$$

where  $\chi_\ell = S_\ell/\Phi_\ell$ . In strongly factorizable regimes ( $\chi_\ell \ll 1$ ), obstruction flow remains bounded and global reconstruction modes persist stably. Near the admissibility transition ( $\chi_\ell \rightarrow 1$ ), dispersion obstruction may drive  $\Omega_\ell \rightarrow \infty$  even in formally compatible local systems.

## D.4 Galactic Anomalies as Cohomological Failure

The galactic anomaly reinterpretation of Section 5 may be reformulated cohomologically: flat rotation curves and lensing anomalies correspond to growth of persistence obstruction

in the global reconstruction geometry rather than to missing matter sources. This clarifies why anomalies appear primarily in large-scale extension behavior rather than in local dynamical inconsistency.

A completed reconstruction-cohomology program would require: explicit admissibility-chain complexes, construction of scale-dependent obstruction operators, definition of persistence cohomology under admissibility renormalization, and derivation of observational signatures associated with obstruction growth.

## E Relation to Renormalization and Effective Field Theory

This appendix clarifies the relation between RSVP’s reconstruction geometry and ordinary Wilsonian effective field theory [18, 19]. The purpose is not to identify RSVP with standard EFT but to explain where the analogy is structurally useful and where the two frameworks diverge.

### E.1 Where the Analogy Holds

In ordinary EFT, short-scale degrees of freedom are integrated out, producing scale-dependent effective couplings governing long-scale behavior. Both frameworks study persistence of effective structure under scale transformation. The analogy motivates the beta-function framework of Appendix B and the universality class hypothesis.

### E.2 Where the Analogy Breaks

In ordinary EFT, the background geometry remains fixed throughout renormalization; only couplings evolve. In RSVP, coarse-graining modifies not merely effective couplings but the factorization structure of admissible reconstruction itself. This is the formal content of Remark 2.3: the reconstruction cover is a derived object of the field geometry, not a background.

Ordinary EFT assumes that locality survives coarse-graining even as effective couplings change. RSVP instead allows the factorization structure supporting effective locality to become unstable. The notion of universality acquires a modified meaning: not that many microscopic systems flow toward the same macroscopic effective behavior, but that distinct admissibility geometries may flow toward the same reconstruction-transition structure under coarse-graining.

### E.3 Newtonian Gravity as Effective Reconstruction Theory

Newtonian dynamics emerges not as a fundamental interaction but as an effective low-dispersion reconstruction theory — analogous to hydrodynamics emerging from microscopic statistical systems [19]. Hydrodynamic laws are not fundamental microscopic truths; they are stable large-scale descriptions arising from strong universality under coarse-graining. Within RSVP, Newtonian dynamics plays the same role.

## E.4 Entropy Equivalence as Emergent Universality

Entropy convergence (Theorem-Schema 4.1) is interpreted as an effective coarse-grained phenomenon analogous to emergent thermodynamic equations of state. Entropy equivalence is not foundational; it is a low-dispersion universality property of strongly factorizable reconstruction. Divergence of entropy channels near the admissibility transition is then unsurprising: universality breaks down because reconstruction factorization itself weakens under renormalization flow.

## F Observational and Experimental Targets

This appendix organizes the empirical targets generated by the framework. It does not claim observational confirmation; its purpose is operational — to identify concrete empirical obligations and failure conditions.

### F.1 Rotation Curve Transition Structure

The framework predicts that transition from Newtonian to MOND-like behavior should correlate with reconstruction-factorization breakdown ( $\chi \rightarrow 1$ ) rather than solely with local acceleration magnitude. This differs subtly from ordinary MOND: RSVP predicts that transition behavior should correlate not only with acceleration but with indicators of reconstruction persistence, including local coherence structure, environmental reconstruction density, and scale-dependent entropy divergence. Low-surface-brightness galaxies are especially important because they probe weak-gradient regimes where  $\chi \rightarrow 1$  over large spatial domains [9].

### F.2 Entropy-Channel Divergence

The most distinctive RSVP prediction: in strongly factorizable regimes ( $\chi \ll 1$ ), entropy channels converge approximately. Near the admissibility transition ( $\chi \sim 1$ ), the framework predicts increasing divergence between channels due to dispersion obstruction growth. Standard  $\Lambda$ CDM and ordinary entropic gravity do not predict entropy-channel divergence as a geometric signal. Operationally, galactic regions exhibiting strong rotational anomalies should exhibit measurable mismatch between local thermodynamic entropy indicators and large-scale gravitational reconstruction structure.

### F.3 Weak Lensing Signatures

Within RSVP, lensing anomalies should correlate with obstruction curvature of the reconstruction cover rather than uniquely with inferred dark matter distributions. Systems with similar baryonic matter distributions but different reconstruction persistence structure may exhibit different lensing behavior. The Bullet Cluster remains a decisive challenge: the framework must reproduce its lensing structure through obstruction curvature geometry (Obligation 5.2).

## F.4 Cosmological Structure Formation

Large-scale structure formation depends not merely on matter density perturbations but on admissibility persistence under cosmological extension. The admissibility radius may evolve dynamically during cosmological history as reconstruction dispersion changes under expansion-like coarse-graining flow. These possibilities remain speculative; the framework currently lacks quantitative cosmological simulations capable of testing them.

## F.5 Observational Failure Modes

The framework generates clear failure conditions: (1) if weak-gradient galactic behavior cannot be associated with coherent admissibility-transition structure, the geometric reinterpretation fails; (2) if cluster-scale lensing cannot be reproduced through obstruction curvature without reintroducing hidden matter, the framework loses explanatory advantage over  $\Lambda$ CDM; (3) if entropy-channel divergence fails to correlate with anomalous galactic regimes, one of RSVP's most distinctive predictions is falsified; (4) if  $R_A$  cannot be derived without arbitrary parameter tuning, the central conjecture becomes structurally underdetermined.

# G Locality, Causality, and Admissibility Nonlocality

The framework employs concepts such as cosmological gluing data, admissibility nonlocality, and global extension dependence. These risk misunderstanding if interpreted as superluminal influence or hidden signaling. This appendix specifies precisely what type of nonlocality RSVP permits and what type it explicitly rejects.

## G.1 The Central Distinction

The framework does *not* propose superluminal communication, violation of relativistic causal structure, or hidden instantaneous signaling between local field degrees of freedom. The central distinction introduced by RSVP is instead the distinction between locality of propagation and locality of admissibility.

The stalk-local dynamics of the reconstruction sheaf remain local. Local field values propagate according to local dynamical equations governing the coupled field

$$X = (\Phi, \mathbf{v}, S),$$

and admissible perturbations evolve continuously through neighboring reconstruction regions. In this sense, RSVP preserves ordinary local propagation structure at the level of stalk dynamics.

What becomes nonlocal is not propagation itself but the coherence conditions governing extension and assembly of local sections into globally persistent reconstruction modes. A local section may satisfy all local dynamical equations while nevertheless failing global

admissibility because its continuation structure cannot be extended coherently across the reconstruction cover under renormalization flow.

Most ordinary field theories implicitly identify locality of propagation with locality of admissibility. RSVP separates these notions. Locality of propagation is a property of stalk dynamics. Locality of admissibility is a property of global extension structure in the reconstruction sheaf.

Admissibility nonlocality therefore appears not in microscopic signaling but in the indexing, restriction, and gluing structure of admissible reconstruction itself. The dependence on cosmological gluing data discussed throughout the main text is not a proposal for instantaneous causal influence across spacetime. It is a statement that persistence of admissible continuation may depend on global reconstruction coherence conditions extending beyond the local neighborhood of a given section.

This distinction becomes especially important near the admissibility transition

$$\chi = \frac{S}{\Phi} \sim 1,$$

where local reconstruction ceases to remain strongly factorizable under coarse-graining. In strongly factorizable regimes,

$$\chi \ll 1,$$

local admissibility appears effectively self-contained because global extension conditions reduce approximately to local continuation structure. Near the transition, however, persistence becomes increasingly sensitive to large-scale reconstruction geometry even while local propagation remains relativistically local.

RSVP therefore weakens locality of admissibility without abandoning locality of propagation. The framework modifies the geometry of globally admissible extension, not the local causal structure of field evolution.

## G.2 Admissibility Nonlocality Defined

Admissibility nonlocality exists in: the indexing structure of the cover, the gluing conditions, and the persistence structure of extension under renormalization. It does *not* exist in the stalk-local dynamical equations themselves. Nonlocality appears in the geometry of assembly rather than in microscopic propagation.

A local section may satisfy all local dynamical equations while nevertheless failing to admit globally stable extension because admissibility persistence depends upon large-scale coherence of the reconstruction cover. The dependence is *extension dependence*, not *dynamical signaling dependence*.

## G.3 Analogy with Gauge Theory and General Relativity

This distinction is analogous in spirit to global constraint structure in gauge theory and general relativity. In electromagnetism, local electric-field measurements are constrained globally by Gauss-law structure. In general relativity, local geometry is constrained by

global consistency conditions on spacetime slicing. RSVP generalizes this: local reconstruction trajectories remain locally dynamical, but admissibility of their persistent extension depends upon global reconstruction geometry.

#### G.4 Interpretation of “Cosmological Gluing Data”

The term does not refer to signals propagating instantaneously from cosmological distances. It refers to the fact that globally persistent continuation may require compatibility with large-scale reconstruction structure. The admissible continuation set of a local trajectory depends upon the global extension geometry of the reconstruction cover — an extension condition, not a propagation condition.

Near the admissibility transition ( $\chi \rightarrow 1$ ), local factorization weakens. Local continuation trajectories cease to remain closed under extension independently of cosmological admissibility structure. The resulting dependence may appear nonlocal from the perspective of ordinary local field theory even though no superluminal propagation occurs.

#### G.5 Formal Statement

Admissibility nonlocality occurs in the indexing and coherence structure of  $\mathcal{F}$  rather than in the stalk-local field dynamics. A completed RSVP field theory would require rigorous proof that admissibility nonlocality remains compatible with relativistic causal structure under all admissible reconstruction flows. The present appendix establishes only the conceptual separation necessary for the framework to avoid immediate conflict with ordinary causal locality. This separation follows directly from the central inversion: persistent physical exchange is generated by admissibility geometry rather than vice versa. If admissibility geometry is primary, then locality of admissibility and locality of propagation need not coincide.

## H Formal Implementation: Grammar, Typed Structures, and Proof Schemas

This appendix outlines possible formal implementation directions for the RSVP admissibility framework. The guiding principle is that the framework should eventually be expressible as a typed formal system in which admissibility, obstruction, entropy-channel convergence, and reconstruction persistence can be checked, simulated, or proven under specified assumptions.

### H.1 BNF Grammar for Reconstruction Objects

A minimal formal language must represent fields, regions, covers, sections, restriction maps, obstruction classes, entropy channels, and scale transformations.

---

Listing 1: BNF grammar for RSVP reconstruction theory

---



```

<Theory>      ::= <FieldDecl> <CoverDecl> <SheafDecl>
                  <DynamicsDecl> <ObstructionDecl> <EntropyDecl>

<FieldDecl>   ::= "field" <Name> "=" "(" <Phi> "," <v> "," <S> ")"
                  "
<Phi>         ::= "Phi" ":" <Domain> "->" Real
<v>           ::= "v"      ":" <Domain> "->" Vector
<S>           ::= "S"      ":" <Domain> "->" Real

<Domain>      ::= "ReconDomain" | <Region>
<Region>      ::= "U" <Index> | <Region> "cap" <Region>
                  | <Region> "cup" <Region>

<CoverDecl>   ::= "cover" <Name> "=" "{" <RegionList> "}"

<SheafDecl>   ::= "sheaf" <Name> ":" <CoverName> "->" <
                  SectionSpace>
<SectionSpace> ::= "Sections" "(" <FieldName> "," <Region> ")"

<RestrMap>    ::= "restrict" <Section> "from" <Region> "to" <
                  Region>

<Admiss>      ::= "admissible" "(" <Section> ")"
                  | "stable"      "(" <Section> ")"
                  | "persistent" "(" <Section> ")"

<Chi>         ::= "chi" "=" <S> "/" <Phi>

<ObsDecl>     ::= "obstruction" <ObsType> "(" <Section> "," <
                  Section> ")"
<ObsType>     ::= "compatibility" | "dispersion"

<EntropyDecl> ::= "entropy" <EntType> "(" <Section> ")"
<EntType>     ::= "informational" | "configurational"
                  | "thermodynamic" | "gravitational"

<DynDecl>     ::= "flow" <Name> ":" <Scale> "->" <TheoryState>

<Formula>     ::= <Admiss> | <ChiRel> | <EntRel> | <ObsRel>
                  | <Formula> "and" <Formula>
                  | <Formula> "implies" <Formula>

<ChiRel>      ::= "chi << 1" | "chi ~ 1" | "chi -> 1"
<EntRel>      ::= "converges" "(" <EntTypeList> ")"
                  | "diverges" "(" <EntTypeList> ")"
<ObsRel>      ::= "bounded" "(" <ObsType> ")"
                  | "grows"   "(" <ObsType> ")"

```

## H.2 BNF for Epistemic Status of Claims

Listing 2: BNF grammar for claim classification

```

<Claim> ::= <Definition>
          | <Conjecture>
          | <TheoremSchema>
          | <ProofObligation>
          | <PhenomInterp>

<Definition>      ::= "define"      <Object> "as" <FormalExpr>
<Conjecture>      ::= "conjecture" <Statement> "under" <
  Assumptions>
<TheoremSchema>   ::= "theorem-schema" <Statement> "provided" <
  Conditions>
<ProofObligation> ::= "prove"        <Statement> "from" <Defs> "and"
  <Dynamics>
<PhenomInterp>   ::= "interpret" <Observation> "as" <
  FrameworkObject>

<Object>          ::= "AdmissibilityRadius"      | "CompatObstruction"
                    | "DispersionObstruction"    | "EntropyChannel"
                    | "ReconstructionCover"      | "
                    AdmissPhaseTransition"

<Relation>        ::= "determines" | "converges_to" | "diverges_from"
                    | "scales_as"   | "obstructs"   | "extends_to"

```

This grammar prevents category errors: “MOND is derived” is not yet a theorem; it is a conjectural derivation target. “The admissibility radius is the relevant geometric object” is a definitional and structural claim. “Entropy channels converge under strong factorization” is a theorem-schema requiring proof under specified dynamics.

## H.3 Typed Structures in Lean Style

Listing 3: Lean-style typed structures for RSVP objects

```

structure ReconDomain where
  points : Type

structure Region (R : ReconDomain) where
  carrier : Set R.points

structure RSVPField (R : ReconDomain) where
  Phi : R.points -> Real
  v    : R.points -> Vector
  S    : R.points -> Real

```

```

-- Control parameter (well-defined when Phi > 0)
def PositiveAccess {R : ReconDomain}
  (X : RSVPField R) : Prop :=
  ∀ x : R.points, X.Phi x > 0

def chi {R : ReconDomain}
  (X : RSVPField R) (h : PositiveAccess X)
  (x : R.points) : Real :=
  X.S x / X.Phi x

-- Admissibility predicates
structure LocalSection {R : ReconDomain}
  (X : RSVPField R) (U : Region R) where
  field_restr : U.carrier -> RSVPField R
  continuation_data : Type
  bounded_stability : Prop

def StronglyFactorizable {R : ReconDomain}
  (X : RSVPField R) (h : PositiveAccess X)
  (U : Region R) (δ : Real) : Prop :=
  ∀ x ∈ U.carrier, chi X h x < δ

def NearTransition {R : ReconDomain}
  (X : RSVPField R) (h : PositiveAccess X)
  (U : Region R) (ε : Real) : Prop :=
  ∀ x ∈ U.carrier, |chi X h x - 1| < ε

```

## H.4 Reconstruction Sheaf Structure

Listing 4: Lean-style reconstruction sheaf

```

structure ReconSheaf (R : ReconDomain) (X : RSVPField R) where
  sections : Region R -> Type
  restrict :
    ∀ {U V : Region R},
      V.carrier ⊆ U.carrier ->
        sections U -> sections V
  identity_law : Prop -- restrict id = id
  composition_law : Prop -- restrict (V ⊆ U) after (W ⊆ V) =
                           -- restrict (W ⊆ U)
  gluing_law : Prop -- compatible sections glue

```

## H.5 Obstruction Formalization

Listing 5: Compatibility and dispersion obstruction in Lean

```

def CompatObstruction {R : ReconDomain} {X : RSVPField R}
  (F : ReconSheaf R X)
  (U V : Region R)
  (sU : F.sections U) (sV : F.sections V) : Prop :=
-- restrictions to overlap disagree
¬(F.restrict (overlap_le U V) sU =
  F.restrict (overlap_le' U V) sV)

structure Scale where
  ell : Real
  pos : ell > 0

-- Persistence functional placeholder
def Persistence {R : ReconDomain} {X : RSVPField R}
  (F : ReconSheaf R X) (ℓ : Scale) (s : Type) : Real :=
  0 -- to be derived from field equations

def DispersionObstruction {R : ReconDomain} {X : RSVPField R}
  (F : ReconSheaf R X) (ε : Real) (s : Type) : Prop :=
  ∃ ℓ : Scale, Persistence F ℓ s < ε

```

## H.6 Theorem Schemas in Lean Style

Listing 6: Central theorem schemas as Lean proof obligations

```

-- Entropy convergence schema
theorem entropy_convergence
  {R : ReconDomain} (X : RSVPField R)
  (h : PositiveAccess X)
  (F : ReconSheaf R X)
  (U : Region R) (δ ε : Real)
  (hFact : StronglyFactorizable X h U δ)
  (hBound : dispersion obstruction bounded True) :
  entropy channels converge to order (*ε *)
  True := by
  sorry -- proof obligation: requires field equation dynamics

-- Entropy divergence schema
theorem entropy_divergence
  {R : ReconDomain} (X : RSVPField R)
  (h : PositiveAccess X)
  (F : ReconSheaf R X)
  (U : Region R) (ε : Real)
  (hTrans : NearTransition X h U ε) :
  entropy channels diverge under coarse-graining
  True := by
  sorry -- proof obligation: requires beta-function derivation

```

```

-- Central conjecture as proof obligation
-- (RA proportional to c/H_0 in weak-gradient limit)
theorem RA_scaling_conjecture
  (field_eqs : True) :
    True := by
  sorry -- principal open problem of the paper

```

The sorry markers are not weaknesses: they identify proof obligations. A Lean formalization would force every informal claim in the paper to become either a definition, theorem, conjecture, or unresolved proof target — precisely the discipline the present paper maintains in prose.

## H.7 Algorithmic Reconstruction Procedure

A computational implementation of RSVP reconstruction obstruction analysis:

Listing 7: RSVP obstruction analysis algorithm

```

Algorithm: RSVP_Obstruction_Analysis

Input:
  X = (Phi, v, S)           -- coupled RSVP field
  R                         -- reconstruction domain
  U = {U_alpha}             -- initial admissibility cover at
    l_min
  l_min, l_max              -- scale range
  epsilon                   -- stability threshold
  delta                     -- factorization threshold

Output:
  R_A                       -- admissibility radius
  compat_obs_map             -- compatibility obstruction map
  disp_obs_map              -- dispersion obstruction map
  entropy_divergence_profile -- entropy channel divergence

Procedure:
  1. Initialize cover U at scale l = l_min.
  2. For each U_alpha:
      compute chi = S / Phi
      if chi < delta: classify as strongly factorizable
  3. For each overlap U_alpha cap U_beta:
      restrict local sections to overlap
      test compatibility agreement
      if disagreement: record compatibility obstruction
  4. Coarse-grain cover to next scale l.
  5. Update Phi_l, v_l, S_l.
  6. Recompute chi_l = S_l / Phi_l.

```

```

7. Track persistence of previously admissible sections.
8. If compatible sections fail persistence: record disp.
   obstruction.
9. If  $\chi_l \rightarrow 1$  or persistence < epsilon:
   record  $R_A = \text{current scale}$ 
10. Compute entropy projections  $\Pi_{\text{therm}}, \Pi_{\text{info}}, \Pi_{\text{grav}},$ 
     $\Pi_{\text{conf}}$ .
11. Measure entropy-channel convergence or divergence.
12. Repeat steps 4-11 until  $l = l_{\text{max}}$ .
13. Return obstruction and transition profiles.

```

## H.8 Formalization Roadmap

A rigorous implementation program would proceed in stages:

- (1) Define the typed syntax and semantics of RSVP reconstruction objects.
- (2) Implement the reconstruction sheaf in Lean 4 or Agda.
- (3) Formalize compatibility obstruction through Čech cocycle data.
- (4) Define dispersion obstruction as scale-instability of persistence.
- (5) Implement coarse-graining flow for  $\chi_\ell$ .
- (6) Prove conditional theorems linking strong factorization to entropy convergence.
- (7) Prove or refute the central conjecture: weak-gradient RSVP determines  $R_A \propto c/H_0$ .
- (8) Build numerical simulations computing obstruction maps and entropy-channel divergence profiles from candidate field equations.

The purpose of formalization is not merely aesthetic rigor. It is a guardrail against the main failure mode of speculative gravity frameworks: sliding between definitions, metaphors, conjectures, and derivations without marking the transition. A Lean implementation would force every RSVP claim to declare its type.

## References

- [1] R. E. Kastner and A. Schlatter, “Gravity from Transactions: Fulfilling the Entropic Gravity Program,” *preprint* (2024).
- [2] E. Verlinde, “On the Origin of Gravity and the Laws of Newton,” *J. High Energy Phys.* **2011**, 29 (2011).
- [3] E. Verlinde, “Emergent Gravity and the Dark Universe,” *SciPost Phys.* **2**, 016 (2017).
- [4] T. Jacobson, “Thermodynamics of Spacetime: The Einstein Equation of State,” *Phys. Rev. Lett.* **75**, 1260 (1995).
- [5] T. Jacobson, “Entanglement Equilibrium and the Einstein Equation,” *Phys. Rev. Lett.* **116**, 201101 (2016).
- [6] T. Padmanabhan, “Thermodynamical Aspects of Gravity: New Insights,” *Rep. Prog. Phys.* **73**, 046901 (2010).
- [7] M. Milgrom, “A Modification of the Newtonian Dynamics as a Possible Alternative to the Hidden Mass Hypothesis,” *Astrophys. J.* **270**, 365 (1983).
- [8] J. D. Bekenstein, “Relativistic Gravitation Theory for the Modified Newtonian Dynamics Paradigm,” *Phys. Rev. D* **70**, 083509 (2004).
- [9] B. Famaey and S. McGaugh, “Modified Newtonian Dynamics (MOND): Observational Phenomenology and Relativistic Extensions,” *Living Rev. Relativ.* **15**, 10 (2012).
- [10] S. S. McGaugh, “A Novel Test of the Modified Newtonian Dynamics with Gas Rich Galaxies,” *Phys. Rev. Lett.* **106**, 121303 (2011).
- [11] M. Van Raamsdonk, “Building Up Spacetime with Quantum Entanglement,” *Gen. Relativ. Gravit.* **42**, 2323 (2010).
- [12] B. Swingle, “Entanglement Renormalization and Holography,” *Phys. Rev. D* **86**, 065007 (2012).
- [13] G. E. Bredon, *Sheaf Theory*, 2nd ed. Springer, New York, 1997.
- [14] M. Kashiwara and P. Schapira, *Categories and Sheaves*. Springer, Berlin, 2005.
- [15] S. Mac Lane and I. Moerdijk, *Sheaves in Geometry and Logic*. Springer, New York, 1992.
- [16] R. Bott and L. W. Tu, *Differential Forms in Algebraic Topology*. Springer, New York, 1982.
- [17] A. Hatcher, *Algebraic Topology*. Cambridge University Press, Cambridge, 2002.

- [18] K. G. Wilson and J. Kogut, “The Renormalization Group and the  $\epsilon$  Expansion,” *Phys. Rep.* **12**, 75 (1974).
- [19] N. Goldenfeld, *Lectures on Phase Transitions and the Renormalization Group*. Addison-Wesley, Reading, MA, 1992.
- [20] E. T. Jaynes, “Information Theory and Statistical Mechanics,” *Phys. Rev.* **106**, 620 (1957).
- [21] S. Amari, *Information Geometry and Its Applications*. Springer, Tokyo, 2016.