Nothing Shields What Isn't a Field: The Podkletnov Rotating Superconductor Claims

Flyxion
Independent Researcher

Abstract

Eugene Podkletnov's reported observation of a small weight loss above a rapidly rotating superconducting disc has circulated for three decades as evidence that gravitational effects can be locally shielded by a rotating quantum condensate. This paper argues that the claim rests on a category error, treating gravitation as a field with a shieldable flux analogous to electromagnetism, when on an entropic and geometric account gravity is instead the large-scale organization of an ordered configuration with no local flux for any material, superconducting or otherwise, to intercept. The paper first sets out the history of the claim and its theoretical sponsors, then examines the replication record, then separates the mundane artifacts that can mimic a weight anomaly from the structural reason no shielding mechanism is available. It closes by treating the standard defense of the claim, that the effect is real but dependent on an undisclosed recipe, as a pattern that deserves a name of its own. The failure of independent replication is read here not as an experimental shortfall but as the expected outcome of an experiment built on a mechanism that was never available. The paper does not prove that no future observation could show a gravitational coupling to a superconductor; it argues only that the shielding interpretation has no support in the evidence presented for it.

1 Introduction

In 1992, Podkletnov and Nieminen reported in *Physica C* that a small non-magnetic, non-conducting sample suspended above a levitating, rotating disc of $YBa_2Cu_3O_{7-x}$ lost a fraction of its weight, on the order of a few hundredths of a percent in the first report [1]. The authors proposed that a state of energy inside the crystal structure at low temperature altered the usual interaction among electromagnetic, nuclear and gravitational forces, producing what they called a gravitational force shielding. Later accounts raised the figure. A 1997 paper described a composite disc driven by an applied electromagnetic field and reported weight losses reaching roughly two percent, with a cylindrical region of reduced weight said to extend vertically for a considerable distance above the apparatus [2]. A later preprint with Modanese extended the claim to an impulse generator based on a charged superconductor [3].

The claim has never been dismissed by the community that could have dismissed it fastest, because it arrived with a theoretical sponsor. Li and Torr had argued in 1991 that gravitational fields could modify the magnetic attenuation of superconductors, and a rotating lattice of aligned ions was proposed as a means of amplifying a gravitomagnetic response to detectable size [4]. The Podkletnov result looked like the experimental confirmation of a prediction, which is how a single unreplicated measurement acquired a following. Several well-instrumented groups then attempted to reproduce it, and none reported success. The question this paper takes up is not whether those attempts were sufficiently faithful to the original. It is whether the original claim ever had a mechanism that could be faithful or unfaithful to anything.

2 The Record of the Claim

The documentary record of the original experiment is unusually thin for a result of this significance. The 1997 account was published in a journal whose editorial scope did not centrally include gravitation [2]. Reviews of the replication literature note that no photographs or video of the original apparatus operating are known to exist, and that the details of the disc's manufacture were described only in general terms [13]. That absence matters because the effect, if real, was said to depend sensitively on disc structure, rotation, cooling and the applied field. A claim whose enabling conditions are unrecorded cannot be falsified by any particular failure to meet them, and it cannot be confirmed by any particular success either.

The theoretical sponsors did not close this gap. Bull and de Podesta proposed in 1995 an alternative explanation of the screening experiments that did not require any modification of gravity [5]. Unnikrishnan asked the following year whether a superconductor could shield gravity at all, and his analysis was likewise skeptical of the shielding interpretation [7]. Modanese attempted a theoretical analysis of the reported effect and concluded that it has no explanation in standard gravity theories, appealing at most to non-perturbative Euclidean quantum gravity [6]. A sponsor who finds that the standard theory excludes the effect is not supplying a mechanism for it.

3 The Category Error

The shielding analogy borrows its plausibility from electromagnetism, where a superconductor genuinely does expel magnetic flux and can meaningfully be said to shield a region from a magnetic field, because the magnetic field is a local, sourced quantity with field lines that a material can intercept, redirect, or exclude. Gravitation, on the account assumed here, is not this kind of object. It is the geometry induced by the large-scale redistribution of an ordered configuration, tracked through coupled scalar, vector, and entropy fields whose curvature effects emerge at scales where mass-energy content is significant relative to the surrounding structure. A superconducting disc a few centimeters across, however exotic its internal quantum state, does not carry enough mass-energy to source a locally significant curvature in the first place, and more importantly has no flux of the relevant kind to expel, because gravitational curvature is not carried as a local flux through a material the way magnetic field lines are.

Asking a rotating superconductor to shield gravity is, on this account, not asking it to do something difficult; it is asking it to do something that has no available mechanism, in the same way that asking a Faraday cage to block the passage of time is not a difficult engineering problem but a request addressed to the wrong category of thing entirely.

The point can be made without committing to the entropic account. In general relativity, the equivalence of inertial and gravitational mass means that every form of energy couples to the gravitational field in the same proportion. A shield would have to be a material that decouples some body from this field, and the equivalence principle is the statement that no such decoupling exists, since a body that responded differently would reveal itself by falling differently in a uniform field. The interplay is why the history of attempts at gravity shielding, from Majorana's absorption proposals onward, has consistently ended in a requirement that something else in the physics change, such as a violation of the equivalence principle or the existence of a gravitational conductor analogous to an electrical one. No such conductor has been found, and the superconducting case adds nothing to the argument, because the superconducting state modifies how charged carriers move and how magnetic flux is treated, neither of which is the quantity that a weight measurement tracks.

The more modest reading of the Li and Torr program is that rotating or accelerated condensates might generate a gravitomagnetic field through a London-type moment, a prediction that is distinct from shielding [4]. That prediction has been tested directly. Laboratory searches for gravitomagnetic fields near rotating

superconductors, carried out with sensitive accelerometers and gyroscopes, reported at first an anomalous signal whose interpretation depends on instrumental systematics [12]. A weight loss of one or two percent in a sample above a disc would be an effect of a different and far larger kind than the gravitomagnetic signals those instruments searched for, and a null at the instrument level does not by itself rule out a geometry-specific shielding. It does remove the supporting claim that rotating superconductors generally show anomalous gravitational coupling, which leaves the shielding effect resting on its recipe alone.

4 What the Original Measurement Likely Captured

The more mundane candidates for the reported weight anomaly are well documented in the replication literature: air currents induced by the disc's rotation, vibration coupling into the weighing apparatus, and small systematic errors in a measurement setup not originally designed with the rigor a claim of this magnitude would require. A rotating disc cooled with liquid nitrogen sits in a convective environment. Cold gas boiling off the cryogen produces buoyancy and drag forces on a suspended sample, and an electrically driven rotating field produces eddy currents and magnetic forces on any sample with the slightest susceptibility. A balance that reads weight changes of a few hundredths of a percent on a few grams is measuring changes in force of the order of micronewtons, a range in which thermal drift, gas flow and magnetic coupling are all capable of producing signals.

Subsequent attempts using vacuum enclosures, vibration isolation, and higher-precision balances have consistently failed to reproduce an effect above the noise floor. Li and collaborators at the Marshall Space Flight Center, working with type-II YBCO superconductors, found no anomalous gravitational force in a static test [8]. Woods and colleagues, attempting to reproduce the experiment with discs of up to roughly three inches in diameter, reported no effect and argued that several of the conditions required by the original claim had not been met [9]. Hathaway, Cleveland and Bao ran a rotating superconducting disc with radio frequency fields and reported no weight modification down to a level of about one part in 10^5 [10]. Reviews of the field restate this record [13], and a separate set of weight measurements of superconductors passing through their critical temperature, including under extremely-low-frequency radiation, found no anomaly within the accuracy of the equipment [11].

The pattern of these outcomes is informative in itself. A real effect that was small and fragile would be expected to show up occasionally, in a minority of runs, in some laboratories, with magnitudes that scale in a regular way with the parameters varied. What the record shows is an effect that appears in the claim and disappears under controls, which is the pattern expected of an artifact rather than a real, if small, physical effect that merely requires better instruments to see more clearly.

5 Objections

A frequent defense holds that Podkletnov's original apparatus used a specific multi-layer disc geometry that subsequent replication attempts did not faithfully reproduce, and that the effect is real but fragile, dependent on manufacturing details that have never been fully disclosed. This is possible in principle for any exotic material effect, but it is also the standard shape of an unfalsifiable defense, since it places the burden of failure on the replicator's fidelity to an incompletely specified recipe rather than on the underlying claim, and it has not, in three decades, produced a second successful independent construction of the disc from any research group willing to publish the attempt.

A second objection is that institutional replication attempts were underpowered or conducted by groups with a stake in a negative result. This is difficult to maintain given that the principal replicators included people who were sympathetic to the underlying theory and who began with the expectation of confirmation.

Li, Noever and their colleagues were working inside a program that regarded the effect as plausible, and a program that announces a null result against its own expectation is not behaving as a motivated critic [8].

A third objection is that the absence of a conventional mechanism is not evidence of absence, since many discovered effects lacked a theory when first observed. This is correct, and it supports the opposite conclusion to the one intended. Effects that lacked a theory at discovery, superconductivity itself being the standard example, were nevertheless reproduced by independent groups within months because the observation was robust and the recipe was shared. What distinguishes the Podkletnov case is not the lack of theory but the lack of reproduction, and the lack of reproduction is what the shielding hypothesis was meant to survive.

6 What Remains Open

Nothing in this paper establishes that a rotating superconductor has no gravitational coupling of any kind. Small gravitomagnetic couplings to rotating matter are predicted by general relativity, they have been measured in satellite experiments with the appropriate precision, and no argument here excludes that exotic condensates might show deviations at the edge of present sensitivity. What the argument excludes is the particular claim that a benchtop device can reduce the weight of an object by percentage-scale amounts without a corresponding source of the lost momentum and energy. That claim requires a mechanism, and the one on offer does not exist. It also requires that the result survive an apparatus whose construction and operation are recorded in enough detail for another group to build it, and that requirement has been available to satisfy for three decades.

7 Conclusion

The rotating superconductor claim treats gravity as a field with a local flux that a sufficiently exotic material might exclude, an analogy borrowed wholesale from electromagnetism without any independent argument that gravitation actually behaves this way. On an account where gravitational curvature is the geometric bookkeeping of a large-scale ordered configuration rather than a locally fluxed field, there is no shielding mechanism available to a benchtop superconductor regardless of how carefully it is built or how fast it is spun, which is consistent with three decades of failed replication being exactly what one would expect rather than an ongoing engineering shortfall. This paper does not support, and does not claim to prove, any positive hypothesis about how the original observation arose; it states only that the evidence offered for shielding is insufficient, and that the structure of the claim makes it insufficient in a way that further effort in the same direction is unlikely to repair.

References

[1] E. Podkletnov and R. Nieminen, A Possibility of Gravitational Force Shielding by Bulk $YBa_2Cu_3O_{7-x}$ Superconductor, *Physica C* 203 (1992) 441–444.

[2] E. Podkletnov, Weak Gravitation Shielding Properties of Composite Bulk $YBa_2Cu_3O_{7-x}$ Superconductor Below 70 K Under E.M. Field, *Journal of Physics D* 30 (1997) 279, arXiv:cond-mat/9701074.

[3] E. Podkletnov and G. Modanese, Impulse Gravity Generator Based on Charged $YBa_2Cu_3O_{7-y}$ Superconductor with Composite Crystal Structure, arXiv preprint, 2001.

[4] N. Li and D. G. Torr, Effects of a Gravitational Field on the Magnetic Attenuation of Superconductors, *Physical Review D* 43 (1991) 457–459.

[5] M. Bull and M. de Podesta, Alternative Explanation of Gravitational Screening Experiments, *Physica C* 253 (1995) 199–200.

[6] G. Modanese, Theoretical Analysis of a Reported Weak Gravitational Shielding Effect, *Europhysics Letters* 35 (1996) 413, arXiv:hep-th/9505094.

[7] C. S. Unnikrishnan, Does a Superconductor Shield Gravity?, *Physica C* 266 (1996) 133–137.

[8] N. Li, D. Noever, T. Robertson, R. Koczor and W. Brantley, Static Test for a Gravitational Force Coupled to Type II YBCO Superconductors, *Physica C* 281 (1997) 260–267.

[9] R. C. Woods, S. G. Cooke, J. Helme and C. H. Caldwell, Gravity Modification by High-Temperature Superconductors, AIAA Paper 2001-3363, 2001.

[10] G. Hathaway, B. Cleveland and Y. Bao, Gravity Modification Experiment Using a Rotating Superconducting Disk and Radio Frequency Fields, *Physica C* 385 (2003) 488–500.

[11] M. Tajmar, K. Hense, K. Marhold and C. J. de Matos, Weight Measurements of High-Temperature Superconductors during Phase Transition in Stationary, Non-Stationary Condition and under ELF Radiation, arXiv:gr-qc/0404005, 2004.

[12] M. Tajmar, F. Plesescu, B. Seifert and K. Marhold, Measurement of Gravitomagnetic and Acceleration Fields Around Rotating Superconductors, arXiv:gr-qc/0610015, 2006.

[13] A. Gallerati, G. Modanese and G. Ummarino, Interaction Between Macroscopic Quantum Systems and Gravity, arXiv:2206.07574, 2022.