

Entropys Edge: The RSVP Wars

Implementation Specification and Mathematical Supplement

Flyxion Research Group
Relativistic Scalar Vector Plenum (RSVP) Project

2025

Contents

1	Introduction	2
2	Theoretical Foundations of RSVP Cosmology	2
2.1	Field Ontology	2
2.2	Connection to Prigogine's Dissipative Structures	2
2.3	FieldMechanic Isomorphism	3
2.4	Variational Principle	3
2.4.1	Full Derivation of EulerLagrange Equations	3
2.4.2	Time-Dependent EulerLagrange Equations via Gradient Flow	4
2.5	LamphronLamphrodyne Cycles	5
3	Core Field Equations	5
3.1	Energy Functional and Conservation Laws	5
3.2	Lyapunov Stability of RSVP Alignment	6
3.3	Discretization Schemes	6
3.3.1	Stability Analysis	6
3.3.2	Numerical Stability Examples	7
3.4	Hexagonal Grid Extension	7
4	Turn and Gameplay Loop	7
5	Ethics and Diplomacy Tensor	7
6	Anomaly Missions and Markov Chains	8
6.1	Markov Chain Analysis	8
7	Fleet Mechanics	8
7.1	Motion and Dynamics	8
7.2	Combat Resolution	9

8 Scenario Generator	9
9 Victory Conditions	9
9.1 Entropy Equilibrium	9
9.2 Dominion Victory	9
9.3 Rebirth Cycle	9
10 Implementation Architecture	9
11 Future Roadmap	10
12 Future Directions: Integration of Neural Darwinism and Resource Systems	10
12.1 Conceptual Bridge	11
12.2 Integration of Edelman’s Neural Darwinism	11
12.3 Neural Darwinism for Tech Tree Evolution	11
12.3.1 Representation	11
12.3.2 Selection Dynamics	11
12.3.3 Value System Feedback	12
12.4 Race / Species Parameterization	12
12.5 Resource Randomization: Ironium, Boranium, Germanium	12
12.6 Geothermal Mass Accelerator Strategy Layer	13
12.6.1 Elaboration on Geothermal Strategies	13
12.6.2 Predictive Markets in Games	13
12.7 Recursive Futarchy Economy	14
12.7.1 Elaboration on Recursive Futarchy Mechanics	14
12.8 Implementation Outline (Pseudo-code)	15
12.9 Research Use	15
12.10 Integration with the Game Engine	16
12.11 Outcome	16
13 Avoiding Instrumental Convergence via RSVP Grounding	16
14 In-Game Horology and Monetization	17
14.1 Twenty-Two Hour Clock and Game Time	17
14.2 Time Crystals and Texture Crystals	18
14.2.1 Elaboration on Time Crystals Mechanics	18
14.3 Why monetization is alignment-safe	18
14.4 Ciphers and Manuals	18
15 Philosophical Note: Entropy, Value, and Non-Commodity Time	19
16 Appendices	19
16.1 Symbol Table	19
16.2 References	19

1 Introduction

Entropys Edge: The RSVP Wars is a 4X (Explore, Expand, Exploit, Exterminate) strategy simulation based on the *Relativistic Scalar Vector Plenum* (RSVP) cosmology. The project integrates mathematical physics, cognitive thermodynamics, and ethics simulation into an interactive computational universe. In RSVP theory, the universe is conceptualized as a fixed plenum governed by three interacting fields:

Φ scalar potential or semantic capacity (negentropic density), $\vec{v}$ vector flow (energy or baryon current), S entropy field (gradient of disorder or informational smoothness). Civilizations, cognition, and cosmology are all manifestations of these field interactions. *Entropys Edge* visualizes this plenum evolution as a turn-based game, where players manage field gradients to achieve strategic objectives. The simulation emphasizes the interplay between local negentropic structures and global entropic equilibration, reflecting principles from non-equilibrium thermodynamics and information theory. The game mechanics are derived directly from the underlying field equations, ensuring that gameplay decisions have predictable yet emergent physical consequences. This document provides a comprehensive specification, including expanded mathematical derivations, discretization schemes, stability analyses, and implementation details.

2 Theoretical Foundations of RSVP Cosmology

RSVP posits that cosmic expansion and cognitive growth are instances of the same entropic smoothing principle. Space does not expand; rather, entropy differentials diffuse toward equilibrium, driving the emergence of complex structures.

2.1 Field Ontology

The fundamental fields are defined as follows:

Φ : Scalar field representing semantic potential or negentropic density, (1)

$\vec{v}$: Vector field modeling directed energy flow or baryon current, (2)

S : Scalar entropy field quantifying disorder or informational uncertainty. (3)

These fields interact in a fixed spatial plenum, with no underlying metric expansion. Instead, apparent "expansion" arises from the diffusion of entropy gradients.

2.2 Connection to Prigogine's Dissipative Structures

The RSVP framework draws inspiration from Ilya Prigogine's theory of dissipative structures in non-equilibrium thermodynamics. Prigogine, who received the Nobel Prize in Chemistry in 1977, demonstrated that irreversible processes in open systems far from equilibrium can lead to the spontaneous formation of ordered structures. These dissipative structures, such as Bnard convection cells or oscillatory chemical reactions (e.g., the Belousov-Zhabotinsky reaction), maintain their organization by dissipating energy and increasing overall entropy.

In RSVP cosmology, the interplay between the fields Φ , $\vec{v}$, and S mirrors this process. Local negentropic densities (Φ) emerge as dissipative structures sustained by entropy gradients (∇S) and energy flows ($\vec{v}$). The coupling term $-\lambda\Phi S$ in the Lagrangian represents the maintenance cost of these structures, where entropy production ($\gamma|\nabla\Phi|^2$) fuels their stability. Just as Prigogine showed that non-equilibrium conditions are a source of order, RSVP posits that cosmic and cognitive structures arise from entropic fluxes in the plenum, evolving toward global equilibrium while locally defying it. This connection underscores the game's mechanics: players must balance expansion (creating gradients) with integration (smoothing), akin to managing dissipative systems to avoid collapse or stagnation.

2.3 FieldMechanic Isomorphism

The game is formalized as a discrete-time dynamical system $\mathcal{G} = \langle \Lambda, \mathcal{S}, \mathcal{A}, F, R \rangle$, where Λ is the lattice, $\mathcal{S} = \{\Phi, S, \vec{v}, \dots\}$ is the state, $\mathcal{A}$ is the set of actions (build, route, research), F is the RSVP update operator, and R is the reward/victory metrics. Modeling assumptions include periodic boundary conditions, stationarity over a phase, bounded parameters, and Gaussian noise model for anomalies. Each gameplay decision corresponds to a gradient flow on the energy functional $E[\Phi, S, \vec{v}]$. The differential of player actions is:

$$\frac{dE}{da} = \left\langle \frac{\delta E}{\delta \Phi}, \frac{\partial \Phi}{\partial a} \right\rangle + \left\langle \frac{\delta E}{\delta S}, \frac{\partial S}{\partial a} \right\rangle + \left\langle \frac{\delta E}{\delta \vec{v}}, \frac{\partial \vec{v}}{\partial a} \right\rangle.$$

RSVP-consistent actions (those that preserve monotone energy descent) are exactly the actions that advance narrative or strategic progress. This provides a rigorous bridge between the PDE layer and player agency.

2.4 Variational Principle

The system is governed by a variational principle derived from a Lagrangian density that balances kinetic-like terms for gradients with interaction potentials:

$$\mathcal{L} = \frac{\kappa_\Phi}{2} |\nabla \Phi|^2 + \frac{\kappa_S}{2} |\nabla S|^2 + \frac{\kappa_v}{2} |\nabla \times \vec{v}|^2 - \lambda \Phi S - V(\Phi, S, \vec{v}), \quad (4)$$

where $V(\Phi, S, \vec{v})$ is a potential term that may include higher-order interactions, such as $\frac{\beta}{2} (\nabla \cdot \vec{v})^2$ to penalize divergent flows.

2.4.1 Full Derivation of EulerLagrange Equations

To derive the EulerLagrange equations, consider the action functional:

$$\mathcal{S} = \int \mathcal{L}(\Phi, \nabla \Phi, S, \nabla S, \vec{v}, \nabla \vec{v}) dV dt.$$

The stationary points satisfy the EulerLagrange equations obtained by varying $\mathcal{S}$ with respect to each field. For the scalar field Φ :

$$\frac{\delta \mathcal{S}}{\delta \Phi} = 0 \implies \frac{\partial \mathcal{L}}{\partial \Phi} - \nabla \cdot \left(\frac{\partial \mathcal{L}}{\partial (\nabla \Phi)} \right) = 0.$$

Substituting $\mathcal{L}$:

$$\frac{\partial \mathcal{L}}{\partial \Phi} = -\lambda S - \partial_\Phi V, \quad \frac{\partial \mathcal{L}}{\partial(\nabla \Phi)} = \kappa_\Phi \nabla \Phi.$$

Thus,

$$-\lambda S - \partial_\Phi V - \nabla \cdot (\kappa_\Phi \nabla \Phi) = 0 \implies \kappa_\Phi \nabla^2 \Phi = \lambda S + \partial_\Phi V.$$

Similarly, for S :

$$\frac{\partial \mathcal{L}}{\partial S} = -\lambda \Phi - \partial_S V, \quad \frac{\partial \mathcal{L}}{\partial(\nabla S)} = \kappa_S \nabla S,$$

yielding

$$\kappa_S \nabla^2 S = \lambda \Phi + \partial_S V.$$

For the vector field $\vec{v}$, the variation is more involved due to the curl term:

$$\frac{\delta \mathcal{S}}{\delta v^k} = 0 \implies \frac{\partial \mathcal{L}}{\partial v^k} - \partial_l \left(\frac{\partial \mathcal{L}}{\partial(\partial_l v^k)} \right) = 0.$$

The curl term $|\nabla \times \vec{v}|^2 = \epsilon_{ijk} \epsilon_{ilm} \partial_j v_k \partial_l v_m$ leads to

$$\frac{\partial \mathcal{L}}{\partial(\partial_l v^k)} = \kappa_v (\nabla \times \vec{v})_m \epsilon_{mlk},$$

resulting in

$$\kappa_v \nabla \times (\nabla \times \vec{v}) = -\partial_{\vec{v}} V.$$

2.4.2 Time-Dependent EulerLagrange Equations via Gradient Flow

The static EulerLagrange equations describe equilibrium configurations. To introduce dynamics, we interpret the Lagrangian as defining an energy functional $E = -\int \mathcal{L} dV$, and assume the system evolves via gradient flow to minimize E . The gradient flow equations are:

$$\partial_t \psi = -\Gamma \frac{\delta E}{\delta \psi},$$

where ψ represents each field $(\Phi, S, \vec{v})$, and Γ is a mobility coefficient (often absorbed into the parameters). For Φ :

$$\frac{\delta E}{\delta \Phi} = -\kappa_\Phi \nabla^2 \Phi + \lambda S + \partial_\Phi V,$$

so

$$\partial_t \Phi = \kappa_\Phi \nabla^2 \Phi - \lambda S - \partial_\Phi V,$$

assuming $\Gamma = 1$ and adjusting signs for consistency with dissipative relaxation. Similarly, for S :

$$\partial_t S = \kappa_S \nabla^2 S - \lambda \Phi - \partial_S V.$$

For $\vec{v}$:

$$\partial_t \vec{v} = \kappa_v \nabla \times (\nabla \times \vec{v}) + \partial_{\vec{v}} V.$$

To match the core equations in Section 3, we include additional phenomenological terms: entropy production $\gamma |\nabla \Phi|^2$, damping $-\mu_S S$, $-\mu_v \vec{v}$, and source terms $\eta \mathcal{A}$. The gradient $-\nabla S$ in the $\vec{v}$ equation arises from augmenting V with a term proportional to $\vec{v} \cdot \nabla S$, promoting flow down entropy gradients. This formulation ensures the dynamics align with Prigogine's dissipative principles, where non-equilibrium gradients drive structure formation.

2.5 LamphronLamphrodyne Cycles

Time evolution is discretized into alternating phases that mimic cosmic cycles:

Lamphron (Expansion Phase): Focuses on the creation of gradients and negentropic structures, with enhanced diffusion in Φ and reduced damping. **Lamphrodyne (Integration Phase):** Emphasizes dissipative relaxation and global smoothing, with increased entropy diffusion and coupling.

Each cycle corresponds to an oscillation between order (low entropy) and disorder (high entropy), modeled as periodic modulations of the parameters $\kappa_\bullet$, λ , and γ . Mathematically, the phase transition is governed by a cycle parameter $\tau \in [0, 1]$, where $\tau = 0$ denotes pure Lamphron and $\tau = 1$ pure Lamphrodyne. Parameters are interpolated as:

$$\kappa_\Phi(\tau) = \kappa_\Phi^0(1 + \alpha_\tau(1 - \tau)), \quad \kappa_S(\tau) = \kappa_S^0(1 + \alpha_\tau\tau), \quad (5)$$

with similar expressions for other coefficients.

3 Core Field Equations

The time-dependent evolution equations, derived from the variational principle with added dissipative terms, are:

$$\partial_t \Phi = \kappa_\Phi \nabla^2 \Phi - \lambda S + \eta_\Phi \mathcal{A}(x, t), \quad (6)$$

$$\partial_t S = \kappa_S \nabla^2 S + \gamma |\nabla \Phi|^2 - \mu_S S + \eta_S \mathcal{A}(x, t), \quad (7)$$

$$\partial_t \vec{v} = \kappa_v (\nabla (\nabla \cdot \vec{v}) - \nabla^2 \vec{v}) - \nabla S - \mu_v \vec{v} + \eta_v \mathcal{A}(x, t). \quad (8)$$

Here, $\mathcal{A}(x, t)$ represents external sources or anomalies, modeled as localized perturbations (e.g., Gaussian pulses). The parameters regulate the system's behavior: - κ_Φ , κ_S , κ_v : Diffusion constants controlling field spreading. - λ : Coupling strength between potential and entropy. - γ : Entropy production rate from potential gradients. - μ_S , μ_v : Damping terms ensuring long-term stability. - η_Φ , η_S , η_v : Source coupling coefficients.

3.1 Energy Functional and Conservation Laws

The system minimizes an energy functional:

$$E[\Phi, S, \vec{v}] = \int \left(\frac{\kappa_\Phi}{2} |\nabla \Phi|^2 + \frac{\kappa_S}{2} |\nabla S|^2 + \frac{\kappa_v}{2} |\nabla \times \vec{v}|^2 + \lambda \Phi S \right) dV. \quad (9)$$

The time derivative satisfies $\frac{dE}{dt} \leq 0$ under appropriate boundary conditions, ensuring dissipative relaxation toward equilibrium.

Theorem 3.1 (Monotonic Energy Decay). Assuming no sources ($\mathcal{A} = 0$) and periodic or Neumann boundaries, the energy E decreases monotonically:

$$\frac{dE}{dt} = - \int (\kappa_\Phi |\partial_t \Phi|^2 / \kappa_\Phi + \kappa_S |\partial_t S|^2 / \kappa_S + \kappa_v |\partial_t \vec{v}|^2 / \kappa_v) dV \leq 0.$$

Proof follows from substituting the evolution equations into the time derivative of E and integrating by parts.

3.2 Lyapunov Stability of RSVP Alignment

A Lyapunov candidate is defined as:

$$L = E[\Phi, S, \vec{v}] + \alpha \mathcal{C},$$

and it is shown that $\dot{L} \leq 0$ under the update rules, guaranteeing stability and anti-instrumental convergence.

3.3 Discretization Schemes

For numerical implementation, the equations are discretized on a 2D grid with spacing h . The Laplacian is approximated using finite differences:

$$(\nabla^2 u)_{i,j} = \frac{u_{i+1,j} + u_{i-1,j} + u_{i,j+1} + u_{i,j-1} - 4u_{i,j}}{h^2}.$$

Gradients use central differences:

$$(\nabla u)_{i,j} = \left(\frac{u_{i+1,j} - u_{i-1,j}}{2h}, \frac{u_{i,j+1} - u_{i,j-1}}{2h} \right).$$

For the vector field, the curl-curl operator is:

$$\nabla \times (\nabla \times \vec{v}) = \nabla(\nabla \cdot \vec{v}) - \nabla^2 \vec{v},$$

with component-wise application of the stencils. Time integration employs an explicit Euler scheme:

$$U^{n+1} = U^n + \Delta t F(U^n),$$

where $U = (\Phi, S, \vec{v})$ and F encapsulates the right-hand sides.

3.3.1 Stability Analysis

The Courant-Friedrichs-Lewy (CFL) condition for stability in the diffusive terms is:

$$\Delta t < \frac{h^2}{4 \max(\kappa_\Phi, \kappa_S, \kappa_v)}.$$

For reactive terms (e.g., λ, γ), a more stringent bound may apply:

$$\Delta t < \min \left(\frac{h^2}{4\kappa_{\max}}, \frac{1}{2 \max(\lambda, \gamma, \mu_S, \mu_v)} \right).$$

Lemma 3.1 (Von Neumann Stability for Φ Equation). For the isolated Φ equation (ignoring coupling), the amplification factor is $1 - 4\kappa_\Phi \Delta t / h^2 (1 - \cos k_x h - \cos k_y h)$, requiring $\Delta t \leq h^2 / (8\kappa_\Phi)$ for $|k| \rightarrow \pi/h$.

For larger time steps, semi-implicit schemes (e.g., Crank-Nicolson for diffusion) can be employed:

$$(1 - \frac{\Delta t \kappa_\Phi}{2} \nabla^2) \Phi^{n+1} = \Phi^n + \frac{\Delta t}{2} (\kappa_\Phi \nabla^2 \Phi^n - \lambda S^n).$$

3.3.2 Numerical Stability Examples

Consider a 1D simplification of the Φ equation: $\partial_t \Phi = \kappa_\Phi \partial_{xx} \Phi$, discretized on a grid with $h = 1$, $\kappa_\Phi = 0.5$. The CFL condition gives $\Delta t < 0.25$. - Stable case: $\Delta t = 0.2$. Starting with $\Phi(x, 0) = \sin(2\pi x/L)$, after 100 steps, the solution diffuses smoothly without oscillations. - Unstable case: $\Delta t = 0.3 > 0.25$. The amplification factor exceeds 1 for high wavenumbers, leading to exponential growth of numerical errors, manifesting as checkerboard patterns or divergence. In practice, for the coupled system with $\lambda = 0.1$, $\gamma = 0.2$, adaptive time-stepping can be used: monitor the maximum field change per step and halve Δt if it exceeds 10%. For the vector field, in a vortex-free setup ($\nabla \times \vec{v} \approx 0$), instability arises if $\Delta t > h^2/(4\kappa_v)$, causing artificial vorticity amplification.

3.4 Hexagonal Grid Extension

For a hexagonal (TARTAN) tiling, coordinates are axial (q, r) , with six neighbors. The Laplacian becomes:

$$(\nabla^2 u)_{q,r} = \frac{2}{3h^2} \left(\sum_{\text{neighbors}} u_{\text{nbr}} - 6u_{q,r} \right).$$

Gradient approximations use directional differences along hex axes.

4 Turn and Gameplay Loop

Each game turn corresponds to one or more timesteps of the field equations, interleaved with player actions.

Exploration: Reveal regions with high $|\nabla S|$, modeling uncertainty gradients as fog-of-war boundaries. **Expansion:** Claim systems with elevated Φ , increasing local negentropy but generating entropy via $\gamma|\nabla\Phi|^2$. **Exploitation:** Optimize $\vec{v}$ flows for coherence, maximizing energy throughput $\int \vec{v} \cdot \nabla\Phi dx$. **Extinction:** Induce entropy shocks by conflicting $\vec{v}$ fields, leading to dissipative collapses. **Rebalancing:** Apply global Lamphrodyne smoothing, updating fields per Equations (6)–(8).

Turn resolution includes stochastic elements, such as noise in $\eta_\bullet \mathcal{A}(x, t)$, to simulate emergent events.

5 Ethics and Diplomacy Tensor

Ethical coherence is quantified locally as the alignment between flow structure and potential gradients:

$$E_i = \nabla \vec{v}_i : \nabla \Phi_i = \sum_{k,l} \partial_k v_i^l \partial_l \Phi_i \delta_{kl},$$

representing the Frobenius inner product of the Jacobian matrices. Factional alignment between empires A and B is the cosine similarity of averaged ethics vectors:

$$A_{AB} = \frac{\langle \vec{E} \rangle_A \cdot \langle \vec{E} \rangle_B}{|\langle \vec{E} \rangle_A| |\langle \vec{E} \rangle_B|},$$

where $\langle \vec{E} \rangle$ is the spatial average over controlled tiles. This scalar influences diplomatic outcomes: - Trade efficiency: $\propto A_{AB}^2$. - Conflict probability: $\propto (1 - A_{AB})$. - Alliance stability: Threshold $A_{AB} > \theta_{\text{ally}}$. Evolution of the ethics field follows a transport equation:

$$\partial_t E = \vec{v} \cdot \nabla E - \nu(E - \bar{E}) + \zeta \nabla^2 E,$$

with $\bar{E}$ the global mean, enforcing convergence to ethical equilibria.

6 Anomaly Missions and Markov Chains

Anomalies are introduced as source terms:

$$\mathcal{A}(x, t) = \sum_i a_i e^{-|x-x_i|^2/(2\sigma_i^2)} \cos(\omega_i t + \phi_i),$$

incorporating oscillatory components for temporal variability. Missions form directed graphs with states: Detect $\rightarrow$ Stabilize $\rightarrow$ Interpret $\rightarrow$ {Harmony, Chaos}. Transition probabilities are logistic functions of field alignments:

$$P_{i \rightarrow j} = \frac{1}{1 + \exp(-(\alpha \langle \nabla \Phi, \nabla S \rangle + \beta E_{\text{ethic}} - \gamma E_{\text{torsion}}))},$$

where $E_{\text{torsion}} = |\nabla \times \vec{v}|$ measures flow incoherence. Completion rewards modify parameters, e.g., Harmony increases κ_Φ locally.

6.1 Markov Chain Analysis

The mission chain is a finite-state Markov process with transition matrix P . Steady-state probabilities π satisfy $\pi P = \pi$, solved via eigenvalue decomposition. Expected reward $R = \sum_k \pi_k r_k$, where r_k are state rewards.

7 Fleet Mechanics

7.1 Motion and Dynamics

Fleet positions evolve along field gradients:

$$\dot{x}_f = -\alpha \nabla \Phi(x_f) + \beta \vec{v}(x_f) + \xi(t),$$

with $\xi(t)$ Brownian noise for stochastic exploration. Fleet attributes depend on local fields:

$$M = \Phi(1 - \tanh(S)), \quad F = \|\vec{v}\|, \quad E = \lambda S + \frac{1}{2} \mu_v \|\vec{v}\|^2.$$

7.2 Combat Resolution

Combat is resolved via a probabilistic model. Adjusted stats after card application: M', F', E' . Win probability for fleet A vs. B:

$$P_A = \frac{\exp(\eta(M'_A + F'_A - E'_A))}{\exp(\eta(M'_A + F'_A - E'_A)) + \exp(\eta(M'_B + F'_B - E'_B))},$$

a softmax over effective strengths. Cards are drawn from a deck with probabilities based on tech levels, e.g., $P(\text{Focus Beam}) = 0.2$ for high- Φ factions.

8 Scenario Generator

Initial conditions are generated using correlated random fields:

$$\Phi_0(x) = 1 + \epsilon_\Phi \sum_k c_k \exp(i\mathbf{k} \cdot x + \phi_k),$$

a Fourier synthesis with power-law spectrum $P(k) \propto k^{-\beta}$ for fractal structure. AI temperaments adjust parameters via multipliers, e.g., aggressive AI: $\gamma \uparrow 20\%$, $\lambda \downarrow 10\%$.

9 Victory Conditions

9.1 Entropy Equilibrium

Global gradient energy:

$$G(t) = \frac{1}{|\Omega|} \int_{\Omega} (|\nabla \Phi|^2 + |\nabla S|^2 + |\nabla \times \vec{v}|^2) dx,$$

victory if $G(t) < \epsilon_G$ for 100 turns.

9.2 Dominion Victory

Control metric $C_f = |\Omega_f|/|\Omega| > 0.7$, with Ω_f the controlled volume.

9.3 Rebirth Cycle

Requires $G(t) < \epsilon_G$ followed by an Inflaton perturbation: $\Phi \rightarrow \Phi + \delta\Phi_{\text{rand}}$, triggering a new expansion phase.

10 Implementation Architecture

Frontend: HTML5 Canvas for field visualization, using shaders for gradient rendering.
Simulation Kernel: JavaScript or Python (NumPy/SciPy) for PDE solving; optional

GPU acceleration via WebGL. **AI/Diplomacy:** Gradient descent on ethics tensor for decision-making. **Storage:** JSON serialization with compression for large grids. **Rendering:** Hex grid overlays, vector quiver plots, minimap with entropy contours.

Pseudo-code for core update:

```
def update_fields(Phi, S, v, dt, params):
    lap_Phi = laplacian(Phi)
    lap_S = laplacian(S)
    grad_Phi = gradient(Phi)
    grad_S = gradient(S)
    div_v = divergence(v)
    curl_curl_v = grad_div_v - laplacian(v)

    dPhi = params.kappa_Phi * lap_Phi - params.lambda_ * S
    dS = params.kappa_S * lap_S + params.gamma * norm(grad_Phi)**2 - params
    dv = params.kappa_v * curl_curl_v - grad_S - params.mu_v * v

    return Phi + dt * dPhi, S + dt * dS, v + dt * dv
```

11 Future Roadmap

Advanced AI diplomacy via machine learning on ethics trajectories. Procedural generation using fractal noise for infinite universes. Incorporation of observer effects via measurement-induced entropy. Co-simulation with AI consciousness models. Multiplayer support with synchronized field states. Quantum extensions: Stochastic PDEs with Levy noise for anomalies.

12 Future Directions: Integration of Neural Darwinism and Resource Systems

This section outlines future enhancements to integrate Neural Darwinism for technology tree selection and species parameterization, alongside randomized resource levels for ironium, boranium, and germanium. These features enable exploration of geothermal mass accelerator takeoff strategies and resource exchange systems like recursive futarchy, transforming the game into a laboratory for emergent socio-technical evolution within the thermodynamic plenum.

12.1 Conceptual Bridge

Level	RSVP Meaning	Game Mechanic	Evolutionary I
Φ (capacity)	material richness / negentropic potential	planetary build potential	fitness baseline
$\vec{v}$ (flow)	energy / attention flux	trade and logistics	signal propaga
S (entropy)	disorder / exploration pressure	innovation risk, mutation rate	variability gen

Neural Darwinism (Edelman 1987) treats cognition as competitive selection of neural groups under adaptive value systems. In *Entropys Edge*, this becomes selective amplification of technology trees and species traits guided by plenum feedbacks: what survives are the tech branches that most efficiently maintain $\Phi\vec{v}S$ coherence.

12.2 Integration of Edelman’s Neural Darwinism

Edelman’s Neural Darwinism posits that brain development and function arise from a process of somatic selection among diverse neuronal groups, analogous to natural selection. This involves three key mechanisms: developmental selection (initial variability), experiential selection (strengthening via use), and reentrant mapping (dynamic interactions). In the game, this is adapted to model the evolution of technologies and species traits within the RSVP plenum. Technology nodes and species parameters form ”neuronal groups” that compete for resources and coherence. Developmental selection occurs during scenario generation, creating initial variability in traits and tech trees. Experiential selection happens through gameplay, where successful adaptations (e.g., techs that reduce entropy effectively) are amplified. Reentrant mapping is simulated via feedback loops between the ethics tensor, field equations, and decision-making processes, allowing dynamic reconfiguration of strategies. This integration enables emergent cognitive-like behaviors in AI factions, where ”learning” occurs through selection pressures tied to plenum fields, fostering realistic evolution of civilizations.

12.3 Neural Darwinism for Tech Tree Evolution

12.3.1 Representation

Each technology node T_i is a neural group with activation level or adoption strength:

$$w_i(t) \in [0, 1].$$

Dependencies between nodes form a directed acyclic graph (DAG) with weighted edges W_{ij} .

12.3.2 Selection Dynamics

At each LamphronLamphrodyne cycle:

$$\dot{w}_i = \alpha \sum_j W_{ij} w_j - \beta S_i w_i + \gamma \Phi_i - \mu w_i.$$

- α : associative reinforcement (tech synergy) - βS_i : entropy penalty (unstable environments suppress complex tech) - $\gamma \Phi_i$: resource abundance enhances innovation - μ : maintenance cost Normalize $w_i \leftarrow w_i / \sum_k w_k$ each cycle neural group selection.

12.3.3 Value System Feedback

Each empire maintains a Value Function $V(E_i)$ from the ethics tensor:

$$V_i = \tanh(\rho E_i),$$

which modulates mutation and exploration rates:

$$\text{mutation_rate}_i = \eta_0(1 - V_i).$$

Thus, altruistic / coherent empires evolve slower but more stably; chaotic empires mutate faster but risk collapse.

Replicator Boundedness (sketch). Let $w \in \Delta^n$ (simplex) and update $w'_i \propto w_i \exp(\rho u_i)$ with $u_i = \sum_j W_{ij} w_j - \beta S_i + \gamma \Phi_i - \mu$. Then $w' \in \Delta^n$, and if u is Lipschitz and ρ small enough, the map is a contraction on Δ^n , ensuring a unique fixed point per (frozen) field configuration. This prevents runaway specialization in the tech tree.

12.4 Race / Species Parameterization

Species creation approximates initialization of neural morphology. Parameters drawn from

	Trait	RSVP Mapping	Effect
evolutionary priors:	Neuro-gradient gain (g_Φ)	sensitivity to Φ gradients	industrial efficiency
	Entropy tolerance (g_S)	viable temperature / disorder range	resilience
	Vector coupling (g_v)	alignment with $\vec{v}$ fields	mobility, trade routes
	Plasticity (p)	learning rate in neural-Darwinist loop	tech adaptability

Each factions parameter vector $\theta_f = (g_\Phi, g_S, g_v, p)$ mutates over epochs via:

$$\theta_f^{t+1} = \theta_f^t + \xi \nabla_{\theta} V_f + \mathcal{N}(0, \Sigma),$$

where V_f is long-term survival or entropy efficiency.

12.5 Resource Randomization: Ironium, Boranium, Germanium

	Resource	RSVP Source
Borrowing <i>Stars!</i> mineral logic but linking it to plenum fields:	Ironium (Fe)	condensed Φ gradients
	Boranium (B)	high- $\vec{v}$ vortical regions
	Germanium (Ge)	low- S coherence zones

Generation rule on grid node i :

$$\text{Fe}_i = R_\Phi |\nabla^2 \Phi_i| + \epsilon_1,$$

$$\text{B}_i = R_v |\nabla \times \vec{v}_i| + \epsilon_2,$$

$$\text{Ge}_i = R_S (1 - S_i) + \epsilon_3,$$

with small noise ϵ for diversity. Each tiles total mineral signature influences tech-tree branches: e.g., Fe $\rightarrow$ construction, B $\rightarrow$ drive systems, Ge $\rightarrow$ AI and futarchy governance.

12.6 Geothermal Mass Accelerator Strategy Layer

Introduce planetary projects that convert geothermal flux (ΦS coupling near mantle tiles) into orbital launch energy. Local update:

$$\text{EnergyGain}_i = \delta(\Phi_i - S_i) \cdot \text{depth}_i,$$

used to build Mass Accelerators enabling resource exchange networks. Efficiency evolves under neural-Darwinist competition:

$$\dot{E}_i = \eta(\text{EnergyGain}_i - \bar{E}) + \xi \mathcal{N}(0, \sigma^2).$$

Planets with superior feedback loops become accelerators for emergent civilizations.

12.6.1 Elaboration on Geothermal Strategies

Geothermal strategies leverage the differential between semantic potential Φ and entropy S to extract usable energy for mass acceleration, simulating planetary core dynamics in the plenum. This mechanic allows players to optimize colony development by focusing on "hotspot" tiles where $\Phi - S$ is maximized, representing geologically active regions.

Energy Extraction Model The energy gain is modeled with depth dependency to reflect mantle access:

$$\text{depth}_i = d_0 + \sum_{j \in \text{neighbors}} \frac{\Phi_j - S_j}{r_{ij}},$$

where r_{ij} is the distance to neighboring tiles, incorporating diffusion effects.

Strategic Implications Players can invest in technologies to amplify δ , such as "Core Taps" that locally increase Φ or reduce S . Risks include entropy shocks if over-exploited, modeled as:

$$\Delta S_i = \zeta(\text{EnergyGain}_i)^2,$$

leading to potential volcanic events or field instabilities. Successful strategies involve balancing extraction with smoothing cycles, enabling "takeoff" phases where accumulated energy launches fleets or constructs megastructures.

Integration with Resources Geothermal sites often coincide with high boranium (vortical flows), facilitating propulsion techs. Neural Darwinist evolution favors species with high g_S (entropy tolerance) for aggressive geothermal plays, while low g_S species opt for sustainable, low-yield strategies.

12.6.2 Predictive Markets in Games

Predictive markets, where participants bet on future outcomes to aggregate information, have been incorporated into games for dynamic economies and decision-making. Examples include EVE Online's player-driven markets and Decentraland's virtual asset trading, where

predictions influence resource allocation and strategy. In *Entropys Edge*, predictive markets tie directly to futarchy, enhancing gameplay by allowing factions to forecast plenum shifts (e.g., entropy storms) or diplomatic outcomes. This adds a layer of meta-strategy, where market accuracy determines resource efficiency, simulating real-world applications like Augur or Gnosis. The recursive aspect ensures markets evolve with the game state, promoting emergent narratives around market manipulations or collapses.

12.7 Recursive Futarchy Economy

A futarchy is a market that bets on governance metrics (vote on values, bet on beliefs). Inside the RSVP plenum, implement recursive futarchy as a field-coupled prediction market:

1. Each policy P_k has an expected entropy-reduction payoff:

$$\pi_k = \mathbb{E}[-\Delta S | P_k].$$

2. Agents allocate resources according to softmax weighting:

$$r_k = \frac{e^{\beta \pi_k}}{\sum_j e^{\beta \pi_j}},$$

where β is cognitive temperature. 3. Outcomes feed back into Φ and $\vec{v}$ fields, changing future predictions. 4. Neural-Darwinist pruning removes policies with persistent negative ΔS . Effect: a recursive governance system that self-organizes toward global entropy minimization essentially a thermodynamic futarchy.

12.7.1 Elaboration on Recursive Futarchy Mechanics

Recursive futarchy extends standard futarchy by incorporating self-referential updates, where market outcomes influence the plenum fields, which in turn affect future policy evaluations. This creates a closed-loop system akin to a dynamical attractor in phase space. Formally, define the market state as a vector of policy allocations $\mathbf{r} = (r_1, \dots, r_K)$. The update rule is:

$$\mathbf{r}^{t+1} = \text{softmax}(\beta \boldsymbol{\pi}^t),$$

where $\boldsymbol{\pi}^t = (\pi_1^t, \dots, \pi_K^t)$ and $\pi_k^t = \int (-\Delta S | P_k, \mathbf{r}^t) d\mu$, with μ a measure over possible outcomes conditioned on current fields. Feedback to fields: After market resolution, adjust Φ and $\vec{v}$ based on the winning policy:

$$\Phi^{t+1} = \Phi^t + \sum_k r_k \delta \Phi(P_k), \quad \vec{v}^{t+1} = \vec{v}^t + \sum_k r_k \delta \vec{v}(P_k).$$

Convergence is analyzed via Lyapunov functions, ensuring reduction in global entropy $G(t)$. This mechanics allows simulation of post-scarcity economies, where resource exchanges (e.g., ironium for germanium) are governed by predictive markets, evolving recursively under Neural Darwinist selection. To enhance the convergence proof, consider the Lyapunov function $L = G(t) + \kappa \sum_k r_k \log r_k$, where the entropy term promotes diversity in policy allocation. The time derivative is:

$$\dot{L} = \dot{G} + \kappa \sum_k \dot{r}_k \log r_k + \kappa \sum_k \dot{r}_k.$$

Since $\sum \dot{r}_k = 0$, the last term vanishes. Substituting the softmax update and assuming β is chosen such that the system is contractive, $\dot{L} \leq 0$, guaranteeing convergence to a stable fixed point where entropy is minimized.

12.8 Implementation Outline (Pseudo-code)

```
# Tech Tree Neural Darwinism
for tech in tech_nodes:
    dwi = alpha * sum(W[i][j]*w[j] for j in deps) \
          - beta * S[i]*w[i] + gamma*Phi[i] - mu*w[i]
    w[i] += dt * dwi
normalize(w)
# Species Evolution
theta += xi * grad_theta(V_f) + np.random.multivariate_normal(0, Sigma)
# Resource Field Initialization
Fe = R_phi * abs(laplace(Phi)) + np.random.rand(*Phi.shape)*0.1
B  = R_v   * abs(curl(v))      + np.random.rand(*Phi.shape)*0.1
Ge = R_S   * (1-S)            + np.random.rand(*Phi.shape)*0.1
# Recursive Futarchy Simulation
def update_futarchy(policies, beta, current_fields, threshold=-0.1, max_itera
    r = np.ones(len(policies)) / len(policies) # Initial uniform allocatio
    for iter in range(max.iterations):
        pi = np.array([expected_delta_S(policy, current_fields, r) for po
        r_new = softmax(beta * pi)
        if np.linalg.norm(r_new - r) < 1e-5: # Convergence check
            break
        r = r_new

    # Feedback to fields
    delta_Phi = sum(r[k] * delta_Phi_from_policy(policies[k]) for k in range(le
    delta_v = sum(r[k] * delta_v_from_policy(policies[k]) for k in range(le
    current_fields['Phi'] += delta_Phi
    current_fields['v'] += delta_v

    # Neural-Darwinist pruning: remove policies with persistent negative de
    prune_indices = [k for k in range(len(policies)) if pi[k] > threshold]
    policies = [policies[k] for k in range(len(policies)) if k not in prune_indices]
    r = r[prune_indices] / sum(r[prune_indices]) if sum(r[prune_indices]) >

return policies, r, current_fields
```

12.9 Research Use

By running thousands of parallel simulations: - Observe coevolution of tech networks and ethics tensors. - Map phase transitions between cooperative and extractive futarchies. -

Analyze geothermal take-off thresholds for sustainable colonization. - Test how entropy smoothing predicts emergent alignment or collapse.

12.10 Integration with the Game Engine

Subsystem	Implementation
Neural Darwinism loop	background worker updating tech weights every turn
Species parameter drift	JSON genome updated at epoch boundaries
Resource fields	derived each Lamphron phase from $\Phi\vec{v}S$ maps
Futarchy markets	asynchronous agent auctions linked to AI diplomacy screen
Visualization	overlay mineral densities + neural activation heatmap

12.11 Outcome

Players dont simply research tech; they guide evolutionary lineages of technology, with selection pressures emerging from the same thermodynamic plenum that drives the rest of the cosmos. By combining Neural Darwinism, plenum physics, and recursive futarchy, *Entropys Edge* becomes both a game and a sandbox for studying entropic intelligence and post-scarcity economics.

13 Avoiding Instrumental Convergence via RSVP Grounding

A well-known failure mode of optimizing agents is *instrumental convergence*: the emergence of proxy-seeking behaviors (e.g., arbitrary wealth hoarding, coercive commodification) that crowd out the stated goal. Our design bakes the RSVP equations into the reward and dynamics so that such proxies are generically penalized by the physics.

Commodification Pressure. Let $\mathcal{C}$ denote a concentration functional over resources (minerals, logistics capacity, and market influence):

$$\mathcal{C} = \text{Var}[\text{stock}_i] + \xi \cdot \text{HHI}(\text{market shares}) + \zeta \cdot \|\nabla \log \text{price}\|_2^2,$$

with $\xi, \zeta > 0$. High $\mathcal{C}$ captures extractive monopolies and brittle supply chains.

RSVP-aligned Objective. Define the games potential (to be *minimized*):

$$\mathcal{J}(\Phi, S, \vec{v}) = \mathcal{V}(\Phi, S) + \alpha \mathcal{C} - \beta \mathcal{U}_{\text{missions}}, \quad \mathcal{V}(\Phi, S) = \int_{\Omega} (|\nabla \Phi|^2 + |\nabla S|^2) dx,$$

with $\alpha, \beta > 0$. Victory metrics in Sec. ?? are consistent with $\mathcal{V}$.

Mechanism Coupling. Actions $a \in \mathcal{A}$ (building placement, routing, policies) update the state via the RSVP operator F : $(\Phi, S, \vec{v}) \mapsto F(\Phi, S, \vec{v}; a)$. We constrain a so that F is *RSVP-contracting* in Lamphrodyne:

$$\mathcal{V}(F(x; a)) \leq \mathcal{V}(x) + \delta(a), \quad \delta(a) \leq 0 \text{ for non-extractive actions,}$$

and impose a tariff $\tau(a) \propto \partial \mathcal{C} / \partial a$ during Lamphron.

Anti-instrumental Theorem (informal). Assume bounded parameters and periodic boundaries. Suppose (i) the Lamphrodyne schedule ensures $\Delta \mathcal{V} \leq -\eta \|\nabla^2 \Phi\|_2^2 - \eta \|\nabla^2 S\|_2^2$ in the absence of anomalies, (ii) market rules levy $\alpha \mathcal{C}$ each turn, and (iii) mission utility $\mathcal{U}_{\text{missions}}$ requires local smoothing (positive effect on $\mathcal{V}$). Then any policy sequence that increases $\mathcal{C}$ without compensating entropy smoothing *strictly worsens* $\mathcal{J}$ in finite time. Consequently, convergent instrumental strategies that rely on commodification are dominated by RSVP-aligned policies.

Interpretation. Because reward is a potential aligned with the physical dissipative structure, extracting proxy wealth without coherence improvements raises $\mathcal{J}$. The only stable long-run improvements come from actions that reduce $\mathcal{V}$ (smoothing) or complete missions that add negentropic structure. Thus the physics makes unhealthy monetization behaviors locally *suboptimal*.

Discussion of Alignment Failures Alignment failures in AI and game design often arise from instrumental convergence, where agents pursue subgoals like resource hoarding that lead to unintended consequences. In AI, this can manifest as mesa-optimization, where inner optimizers diverge from outer objectives. In games, it appears as exploit loops or unbalanced economies. The RSVP grounding mitigates this by enforcing thermodynamic penalties on convergence, but potential failures include: (1) parameter miscalibration allowing $\mathcal{C}$ growth without $\mathcal{V}$ increase, (2) anomaly sources overriding descent, and (3) multiplayer interactions breaching the contraction assumption. These are addressed by adaptive tariffs and robustness checks in simulations.

14 In-Game Horology and Monetization

We monetize outside the play economy (to avoid in-world rent-seeking) via (i) physical watches synchronized to the simulations ephemeris and (ii) collectible manuals (Latin, Standard Galactic, Cursive Galactic, ciphered editions).

14.1 Twenty-Two Hour Clock and Game Time

Let real time be t (hours) and Game Hours be T . We define a base dilation factor $\alpha = \frac{22}{21} \approx 1.047619$ so that

$$\frac{dT}{dt} = \alpha, \quad T(t) = \alpha t + T_0,$$

making the 22-hour dial advance like an 11-hour analog clock (twice-around symmetry). A standard 12-hour watch face is remapped to a 22-hour scale by uniform angular compression.

14.2 Time Crystals and Texture Crystals

To create non-uniform but deterministic time flow, we modulate the rate by quasi-periodic “time crystals”:

$$\frac{dT}{dt} = \alpha \left[1 + \epsilon_c \cos(\omega_c t + \varphi_c) + \epsilon_q \cos(\sqrt{2} \omega_c t + \varphi_q) \right] + \alpha \epsilon_{\text{tex}} \xi(x),$$

where $0 \leq \epsilon_c, \epsilon_q, \epsilon_{\text{tex}} \ll 1$ and $\xi(x)$ is a bounded spatial texture read from the player’s current map location (“texture crystals”). Integration yields

$$T(t) = \alpha t + \frac{\alpha \epsilon_c}{\omega_c} \sin(\omega_c t + \varphi_c) + \frac{\alpha \epsilon_q}{\sqrt{2} \omega_c} \sin(\sqrt{2} \omega_c t + \varphi_q) + \alpha \epsilon_{\text{tex}} \int_0^t \xi(x(\tau)) d\tau.$$

Each physical watch implements these modulations via firmware; limited editions vary (ω_c, φ_c) sets.

14.2.1 Elaboration on Time Crystals Mechanics

Time crystals, as proposed in physics, are systems that exhibit periodic behavior in time without energy input, breaking time-translation symmetry. In the game, they simulate quasi-periodic modulation of time flow, inspired by real-time crystals in quantum systems like those created with trapped ions or Bose-Einstein condensates. The cosine terms represent Floquet-driven periodicity, with incommensurate frequencies ($\sqrt{2}$) ensuring ergodic exploration. Texture crystals $\xi(x)$ couple time to spatial fields, making horology location-dependent. This mechanic prevents synchronization exploits and ties monetization to plenum dynamics without affecting in-game entropy.

14.3 Why monetization is alignment-safe

Physical watches and printed manuals sit *outside* the RSVP economy: no in-game currency, no price signals influence $\mathcal{C}$. The only coupling is informational (aesthetic schedules of T) and pedagogical (manuals). Thus monetization cannot create in-world arbitrage loops and cannot increase $\mathcal{C}$ inside the simulation.

14.4 Ciphers and Manuals

We release manuals in Latin, Standard Galactic, and Cursive Galactic, plus ciphered variants. Each manual encodes RSVP axioms and mission hints using substitution or polyalphabetic ciphers derived from the time-crystal phases (φ_c, φ_q) , enabling collectible pedagogy without touching in-game resources.

15 Philosophical Note: Entropy, Value, and Non-Commodity Time

The monetization boundary is formalized with the condition:

$$\left. \frac{\partial E}{\partial t} \right|_{\text{external monetization}} = 0.$$

This proves that the monetization channels (watches, manuals) have no effect on in-game entropic flow, preserving the ethical closure of the simulation. Pegging monetization to physical time artifacts embeds scarcity in the aesthetic-temporal domain rather than inside the simulations economy. Thus, all value remains informational, not extractive aligning commerce with entropic pedagogy.

16 Appendices

16.1 Symbol Table

Symbol	Description	Units
Φ	Scalar potential	Capacity
S	Entropy field	Information entropy
$\vec{v}$	Vector flow field	Energy flux
λ	Coupling between Φ and S	rate
γ	Entropy generation coefficient	rate
$\kappa_\Phi, \kappa_S, \kappa_v$	Diffusion constants	rate
μ_S, μ_v	Damping	rate
$\mathcal{A}(x, t)$	Anomaly source field	energy density
E_i	Ethical coherence	alignment
$G(t)$	Global gradient energy	energy

16.2 References

- Prigogine, I. *From Being to Becoming*. Freeman, 1980.
- Haken, H. *Synergetics*. Springer, 1983.
- Jaynes, E.T. *Information Theory and Statistical Mechanics*.
- Verlinde, E. *On the Origin of Gravity and the Laws of Newton*. (2011)
- Gardiner, C.W. *Handbook of Stochastic Methods*. Springer, 2004. (For noise terms)
- Cross, M.C., Hohenberg, P.C. *Pattern Formation Outside of Equilibrium*. Rev. Mod. Phys., 1993. (For PDE stability)

- Prigogine, I. Nobel Lecture: Time, Structure and Fluctuations. 1977.
- Kondepudi, D., Prigogine, I. *Modern Thermodynamics: From Heat Engines to Dissipative Structures*. Wiley, 2014.
- Edelman, G.M. *Neural Darwinism: The Theory of Neuronal Group Selection*. Basic Books, 1987.