Renormalization, Asymptotic Safety, and the Scalar-Vector Plenum

A Mathematical Introduction

Flyxion

March 17, 2026

Preface

This book introduces the mathematical and physical concepts required to understand modern approaches to quantum gravity based on renormalization group flows and asymptotic safety, and to follow the development of the Relativistic Scalar-Vector Plenum (RSVP) framework as a geometric field-theoretic approach to cosmology.

The presentation emphasizes intuition first, followed by the mathematical structures that make the theory precise. Readers are assumed to know basic calculus and linear algebra, but no prior knowledge of quantum field theory is required. The book moves from the foundations of physical reasoning through statistical physics, renormalization group theory, quantum field theory, and functional renormalization group methods, and then transitions to the geometric and information-theoretic ideas that underlie the plenum framework and its cosmological consequences.

Each chapter begins with a motivating question and ends with exercises. Key definitions and central ideas are set off typographically. The text aims throughout to connect abstract formalism to physical intuition and to the explicit calculations that appear in research literature on asymptotically safe gravity and scalar-vector field theories.

Flyxion, 2026.

Contents

I Foundations of Physical Theories **1**

1 What Is a Physical Theory? **2**
1.1 Observation, Models, and Laws 2
1.2 Parameters and Coupling Constants 3
1.3 Dimensional Analysis 3
1.4 The Role of Scale in Physics 3

2 Differential Equations in Physics **5**
2.1 Phase Space and Trajectories 5
2.2 Stability and Equilibria 5
2.3 Linearization Near Fixed Points 6

3 Fields and Local Dynamics **7**
3.1 From Particles to Fields 7
3.2 Field Equations 7

II Statistical Physics and Coarse-Graining **9**

4 Probability and Statistical Description **10**
4.1 Microstates, Macrostates, and Entropy 10
4.2 Entropy and Information 11
4.3 Fluctuations and Correlations 11

5 Phase Transitions and Universality **12**
5.1 Order Parameters and Symmetry Breaking 12
5.2 Universality and Critical Exponents 12

6 Coarse-Graining **13**
6.1 Block Spin Transformations 13
6.2 Emergent Laws 13

III Renormalization Group Theory 15

7 Theory Space 16
7.1 Coupling Constants as Coordinates 16
7.2 Effective Actions 16

8 Beta Functions 18
8.1 Running Couplings and the Callan-Symanzik Equation 18
8.2 Renormalization Group Equations 18

9 Fixed Points and Scale Invariance 20
9.1 Gaussian and Non-Gaussian Fixed Points 20
9.2 Critical Surfaces and Predictivity 20

10 Relevant and Irrelevant Operators 21
10.1 Critical Exponents 21
10.2 Predictivity of Renormalizable Theories 21

IV Quantum Field Theory Essentials 23

11 Classical Field Theory 24
11.1 Action Principles and Lagrangians 24
11.2 Symmetries and Noether's Theorem 24
11.3 The Einstein-Hilbert Action 24

12 The Path Integral and Quantization 26
12.1 Path Integral Formulation 26
12.2 Perturbation Theory and Feynman Diagrams 26

13 Renormalization 28
13.1 Regularization and Renormalized Parameters 28
13.2 Effective Field Theories 28

V Functional Renormalization Group 30

14 The Effective Average Action 31
14.1 Motivation and Construction 31
14.2 The Wetterich Equation 31
14.3 Truncation Methods 32

15 Flow Equations and Fixed-Point Search **33**
15.1 Beta Functions from the Flow Equation 33
15.2 Fixed Points and Critical Exponents 33

VI Gravity and Asymptotic Safety **35**

16 Gravity as a Field Theory **36**
16.1 The Metric as a Dynamical Field 36
16.2 Perturbative Non-Renormalizability 36

17 The Asymptotic Safety Program **37**
17.1 Weinberg's Proposal 37
17.2 Evidence for the Non-Gaussian Fixed Point 37
17.3 Matter-Gravity Fixed Points and the Weak-Gravity Bound 38

18 Matter and Gravity **39**
18.1 Scalar Couplings and the Higgs Mass Prediction 39
18.2 Fermions and Gauge Fields 39

VII Cosmology and Observational Signatures **41**

19 Running Couplings in Cosmology **42**
19.1 RG-Improved Friedmann Equations 42
19.2 Cosmological Fixed Points 42

20 Observational Windows **44**
20.1 Gravitational Waves as Quantum Gravity Probes 44
20.2 The CMB and Large-Scale Structure 44

VIII Mathematical Structures **46**

21 Metric Geometry and Theory Space **47**
21.1 Distances Between Theories 47

22 Information Geometry **48**
22.1 The Fisher Information Metric 48
22.2 Thermodynamic Geometry 48

23 Category Theory and Physical Structure **50**
23.1 Functors and Natural Transformations 50

23.2 Sheaves and Local-to-Global Principles 50

IX Computational Experiments 52

24 Numerical Renormalization Group 53
24.1 Integrating Beta Functions 53
24.2 Visualizing RG Flows 53

25 Simulation Projects 55
25.1 Scalar Field Renormalization Group 55
25.2 Gravity-Inspired Flows 55

X Beyond Conventional Renormalization 56

26 Renormalization as Geometric Flow 57
26.1 Theory Space as a Manifold 57
26.2 Gradient Flows and the c-Theorem 57
26.3 Entropy Interpretations of RG Flow 58

27 Information Geometry of Field Theories 59
27.1 Statistical Manifolds of Theories 59
27.2 Thermodynamic Curvature and Criticality 59

28 Coarse-Graining as Entropy Transport 61
28.1 Entropy Production under Coarse-Graining 61
28.2 The Arrow of Scale 61

XI Scalar-Vector Plenum Framework 63

29 Motivation for Plenum Models 64
29.1 Limitations of Metric-Based Spacetime 64
29.2 Field-Based Ontology 64

30 Scalar Fields as Density Structures 66
30.1 Scalar Densities in Physical Theories 66
30.2 Energy Landscapes and Order Parameters 66

31 Vector Flows and Directed Structure 67
31.1 Flux and Transport 67
31.2 Topological Defects and Vorticity 67

32 Entropy Fields **68**
32.1 Entropy as a Dynamical Variable 68
32.2 Lamphrodyne Relaxation 68

XII Field Coupling Structures **70**

33 Coupled Scalar-Vector Dynamics **71**
33.1 Advection-Diffusion and the RSVP Action 71
33.2 Instabilities and Pattern Formation 71

34 Lamphron and Lamphrodyne States **73**
34.1 Definition of Lamphronic Regimes 73
34.2 Field Relaxation Dynamics 73

XIII Cosmological Implications **74**

35 Cosmic Structure without Metric Expansion **75**
35.1 Standard Cosmological Expansion 75
35.2 Entropy Smoothing and Structure Formation 75

36 Formation of Baryonic Structures **77**
36.1 Solitonic Configurations 77

37 Long-Term Energy Redistribution **78**
37.1 Trillion-Year Cosmology 78
37.2 Global Equilibrium States 78

XIV Mathematical Formalism **79**

38 Variational Formulations **80**
38.1 Action Principles for Coupled Fields 80
38.2 Euler-Lagrange Equations 80

39 Topological Structures **81**
39.1 Homology and Cohomology in Field Theory 81
39.2 Topological Defects and Cosmic Strings 81

40 Categorical Descriptions of Dynamics **82**
40.1 Categories of Field Configurations 82
40.2 Compositional Dynamics and Modularity 82

XV Conceptual Consequences 84

41 Emergent Geometry from Field Interactions 85
41.1 Geometry from Correlations 85
41.2 Relation to Asymptotic Safety 85

42 Observers, Measurement, and Coarse-Graining 87
42.1 Observer-Dependent Descriptions 87
42.2 Local Information Extraction 87

XVI Open Problems and Future Directions 89

43 Experimental Tests 90
43.1 Astrophysical Signatures of the RSVP Framework 90
43.2 Gravitational Wave Backgrounds 90

44 Connections to Other Approaches 92
44.1 Loop Quantum Gravity 92
44.2 Causal Set Theory 92
44.3 Tensor Models and Universality Classes 93

45 Future Directions 94
45.1 Non-Polynomial Dynamics 94
45.2 Lorentzian Signature and Real-Time Dynamics 94
45.3 Unified Theoretical Frameworks 95

A Mathematical Background 96
A.1 Linear Algebra Review 96
A.2 Differential Geometry Basics 96
A.3 Functional Analysis 97

B Notation and Conventions 98

C Solutions to Selected Exercises 99

Part I

Foundations of Physical Theories

Chapter 1

What Is a Physical Theory?

1.1 Observation, Models, and Laws

A physical theory is a mathematical structure designed to compress the information contained in observations into a small number of principles and parameters. It does not describe reality directly; it describes a model of reality from which predictions can be derived and compared with measurement. The relationship between theory and observation is therefore always mediated by a process of idealization: we choose which features of a situation to represent mathematically and which to ignore.

This distinction matters enormously in the context of quantum gravity and renormalization. Most theories we use are valid only within a specific range of energy scales, spatial resolutions, or environmental conditions. Newton's laws of motion describe falling apples but break down at velocities near the speed of light. The Standard Model of particle physics works beautifully up to the energies of current colliders but fails to incorporate gravity. Every physical theory, without exception, is an approximation valid within some domain.

Definition 1.1. An *effective theory* is a mathematical framework that describes the behavior of a physical system accurately within a specified range of energy scales, momenta, or length scales, without necessarily providing a valid description outside that range.

The program of renormalization group theory, which occupies the heart of this book, is fundamentally about understanding how effective theories change as one changes the resolution scale at which a system is examined. This is not merely a technical device; it is a profound statement about how physical laws emerge from underlying structure through the process of averaging over inaccessible microscopic degrees of freedom.

1.2 Parameters and Coupling Constants

Any theory contains numerical parameters whose values must be determined by measurement rather than derived from the theory itself. In classical mechanics these might be a particle's mass m and charge q. In a field theory they are called *coupling constants* because they parametrize the strength of interactions between fields.

Example 1.1. In electromagnetism the coupling constant is the elementary charge e, or equivalently the fine-structure constant $\alpha = e^2/(4\pi\varepsilon_0\hbar c) \approx 1/137$. This dimensionless number controls the strength of every electromagnetic interaction.

One of the central discoveries of twentieth-century physics is that coupling constants are not truly constant: their apparent values depend on the energy scale at which they are measured. The fine-structure constant α at the energy scale of the Z boson (~ 91 GeV) is approximately $1/128$, not $1/137$. This running of coupling constants is the subject of renormalization group theory.

1.3 Dimensional Analysis

Before a single equation is written, dimensional analysis constrains what theories are possible. Every physical quantity has dimensions: length $[L]$, time $[T]$, mass $[M]$, and so on. An equation can only be physically valid if both sides have the same dimensions.

In natural units, where$\hbar = c = 1$, all quantities have dimensions of powers of mass (equivalently, powers of energy or inverse length). A scalar field ϕ in four spacetime dimensions has dimension $[\phi] = M^1$. An action $S = \int d^4x\, \mathcal{L}$ must be dimensionless since it appears in the exponent $e^{iS/\hbar}$ of the path integral, so the Lagrangian density $\mathcal{L}$ must have dimension M^4.

> **Key Idea.** The canonical dimension of a coupling constant determines whether the corresponding interaction becomes stronger or weaker at high energies. Couplings with positive mass dimension grow at high energies (relevant operators); those with negative mass dimension shrink (irrelevant operators); dimensionless couplings can go either way and require detailed calculation (marginal operators).

1.4 The Role of Scale in Physics

The most important conceptual tool in this book is the idea that physics at different scales can be effectively decoupled. When we study ocean waves, we do not

need to know the quantum mechanical state of every water molecule. When we calculate the trajectory of a planet, we do not need to account for the thermal fluctuations of its interior. The laws governing large scales can be derived, at least in principle, by averaging over small scales.

This decoupling is not automatic: it holds when there is a separation of scales, meaning that the characteristic energy scales of the phenomena of interest are well separated from the energy scales of the underlying microscopic physics. When such a separation exists, we can write down an effective theory at the macroscopic scale whose parameters encode, in a coarse-grained way, the effects of the microscopic degrees of freedom.

Exercise 1.1. An electron has mass $m_e \approx 0.511$ MeV and charge e. The Bohr radius of hydrogen is $a_0 = \hbar^2/(m_e e^2) \approx 0.53$ Å. Use dimensional analysis to show that any length scale characterizing atomic physics must be proportional to a_0, and estimate the energy of the ground state without solving the Schrödinger equation.

Chapter 2

Differential Equations in Physics

2.1 Phase Space and Trajectories

A physical system is described at any moment by its *state*. For a single particle in one dimension the state is the pair (x, p) of position and momentum. The space of all possible states is called *phase space*, and the time evolution of the system traces a *trajectory* through this space.

Hamilton's equations of motion describe this trajectory:

$$\dot{x} = \frac{\partial H}{\partial p}, \qquad \dot{p} = -\frac{\partial H}{\partial x}, \tag{2.1}$$

where $H(x, p)$ is the Hamiltonian (energy function). These are first-order ordinary differential equations for the pair $(x(t), p(t))$.

The same mathematical structure appears throughout renormalization group theory: the running coupling constants $g_i(k)$ trace trajectories through the space of all coupling constants (theory space), driven by the beta functions $\beta_i = k\partial_k g_i$. Understanding the structure of trajectories in phase space—their fixed points, attractors, and separatrices—is therefore a direct prerequisite for understanding renormalization group flows.

2.2 Stability and Equilibria

A *fixed point* of a dynamical system $\dot{x} = f(x)$ is a point x_* where $f(x_*) = 0$. The system does not move if it starts at a fixed point. The stability of a fixed point is determined by what happens to nearby trajectories: if they are attracted toward x_*, the fixed point is stable (an attractor); if they are repelled, it is unstable (a repeller).

Definition 2.1. A fixed point x_* of a one-dimensional system $\dot{x} = f(x)$ is *stable*

if $f'(x_*) < 0$ and *unstable* if $f'(x_*) > 0$.

2.3 Linearization Near Fixed Points

For a multidimensional system $\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x})$ with a fixed point at $\mathbf{x}_*$, the behavior of nearby trajectories is governed by the linearized system

$$\dot{\delta x_i} = J_{ij}\,\delta x_j, \quad J_{ij} = \left.\frac{\partial f_i}{\partial x_j}\right|_{\mathbf{x}_*}, \tag{2.2}$$

where J is the Jacobian matrix. The solution is $\delta\mathbf{x}(t) = \sum_I C_I \mathbf{v}_I e^{\lambda_I t}$, where $\mathbf{v}_I$ are eigenvectors and λ_I are eigenvalues of J. Directions with $\mathrm{Re}(\lambda_I) < 0$ are attracted toward the fixed point; those with $\mathrm{Re}(\lambda_I) > 0$ are repelled.

This analysis translates directly into the FRG language: the stability matrix $B_{ij} = \partial\beta_i/\partial g_j|_{g_*}$ plays the role of J, and the critical exponents $\theta_I = -\lambda_I$ determine the relevance of each coupling direction.

Exercise 2.1. Consider the two-dimensional system $\dot{x} = -x + y^2$, $\dot{y} = -2y$. Find all fixed points, compute the Jacobian at each, and classify each as stable, unstable, or saddle.

Chapter 3

Fields and Local Dynamics

3.1 From Particles to Fields

When we describe a single particle we need one set of coordinates. When we describe a gas of 10^{23} particles we face the impracticality of tracking each one individually. A *field* is a function that assigns a value to every point in space and time, providing a continuous description of physical quantities distributed over space.

Definition 3.1. A *scalar field* is a function $\phi : \mathbb{R}^{3,1} \to \mathbb{R}$ assigning a real number to each spacetime point. A *vector field* assigns a vector $V^\mu(x)$ to each point.

Fields obey *local* equations: the value of the field at a point evolves in response to the field's values at neighboring points. This locality is a fundamental principle in relativistic theories, ensuring that information cannot propagate faster than light.

3.2 Field Equations

The simplest relativistic scalar field equation is the Klein-Gordon equation:

$$(\Box + m^2)\phi = 0, \quad \Box = \partial_t^2 - \nabla^2. \tag{3.1}$$

This is a linear partial differential equation with plane-wave solutions $\phi \propto e^{i(p\cdot x)}$ satisfying the dispersion relation $p^2 = m^2$. The parameter m is the mass of the field excitations.

Nonlinear field equations arise when interactions are included. Adding a quartic self-interaction gives

$$\Box\phi + m^2\phi + \lambda\phi^3 = 0. \tag{3.2}$$

The coupling constant λ determines the strength of the nonlinearity. In the quantum theory, λ becomes a running coupling whose scale dependence is governed by a beta function.

Exercise 3.1. Verify that $\phi(x,t) = A\cos(kx - \omega t)$ satisfies the linear Klein-Gordon equation and find the relationship between k, ω, and m.

Part II

Statistical Physics and Coarse-Graining

Chapter 4

Probability and Statistical Description

4.1 Microstates, Macrostates, and Entropy

A glass of water contains approximately 10^{25} molecules. Its complete microscopic description would require specifying the position and velocity of each molecule—an incomprehensibly large amount of information. Yet we describe the glass adequately with just a handful of variables: temperature, pressure, and volume. How?

The answer is that most macroscopic measurements do not distinguish between the $10^{10^{25}}$ microscopic states consistent with the same temperature and pressure. A *macrostate* is an equivalence class of microstates that are observationally indistinguishable.

Definition 4.1. The *Boltzmann entropy* of a macrostate is

$$S = k_B \ln \Omega, \tag{4.1}$$

where Ω is the number of microstates compatible with that macrostate and k_B is Boltzmann's constant.

Entropy measures the amount of information about the microstate that the macrostate conceals. A macrostate with higher entropy corresponds to more ways the system could be microscopically arranged while appearing the same macroscopically. The second law of thermodynamics—entropy does not decrease in isolated systems—reflects the overwhelming statistical preponderance of high-entropy macrostates.

4.2 Entropy and Information

Shannon's information theory provides a more general definition of entropy that applies to any probability distribution. Given a discrete probability distribution $\{p_i\}$, the Shannon entropy is

$$H = -\sum_i p_i \ln p_i. \tag{4.2}$$

This measures the average amount of information contained in a single sample from the distribution, or equivalently the uncertainty about the outcome before it is observed.

The connection to physics is deep. The Jaynes maximum-entropy principle states that the equilibrium macrostate is the one that maximizes entropy subject to the known macroscopic constraints (e.g., fixed total energy). This principle underlies the entire edifice of statistical mechanics and, as we will see in Part VIII, provides an information-theoretic interpretation of renormalization group flow.

4.3 Fluctuations and Correlations

Near a phase transition, the fluctuations of the order parameter become very large. The two-point correlation function

$$G(x, y) = \langle \phi(x)\phi(y)\rangle - \langle \phi(x)\rangle\langle \phi(y)\rangle \tag{4.3}$$

measures the statistical dependence between field values at different points. At a critical point, correlations become long-ranged: $G(x, y) \sim |x - y|^{-(d-2+\eta)}$ for a power law with exponent η, rather than the exponential decay $G(x, y) \sim e^{-|x-y|/\xi}$ characteristic of the disordered phase.

Exercise 4.1. For a Gaussian probability distribution $p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-x^2/(2\sigma^2)}$, compute the Shannon entropy $H = -\int p \ln p \, dx$ and show that it increases monotonically with σ.

Chapter 5

Phase Transitions and Universality

5.1 Order Parameters and Symmetry Breaking

A phase transition is a qualitative change in the macroscopic behavior of a system as a parameter (temperature, pressure, magnetic field) crosses a critical value. The Ising model provides the paradigm: a lattice of spins $s_i = \pm 1$ coupled by nearest-neighbor interactions $H = -J \sum_{\langle ij \rangle} s_i s_j$. At high temperature the spins are disordered and the average magnetization $m = N^{-1} \sum_i \langle s_i \rangle = 0$. Below the Curie temperature T_c the spins align and $m \neq 0$.

The *order parameter* m is zero in the disordered phase and non-zero in the ordered phase. The phase transition involves the spontaneous breaking of the $\mathbb{Z}_2$ symmetry $s_i \to -s_i$: the Hamiltonian is symmetric but the ground state is not.

5.2 Universality and Critical Exponents

The most striking feature of second-order phase transitions is universality: the critical exponents describing the behavior near T_c depend only on the dimension of space and the symmetry of the order parameter, not on the microscopic details of the system. The magnetization near the critical point behaves as $m \sim |T - T_c|^{\beta}$, the correlation length as $\xi \sim |T - T_c|^{-\nu}$, and the specific heat as $C \sim |T - T_c|^{-\alpha}$.

For the Ising universality class in three dimensions, $\beta \approx 0.326$, $\nu \approx 0.630$, $\alpha \approx 0.110$. These same exponents describe the liquid-gas transition, the behavior of certain magnetic materials, and even some features of quantum chromodynamics at finite temperature. This universality is explained by the renormalization group: systems in the same universality class flow to the same fixed point under the RG transformation.

Chapter 6

Coarse-Graining

6.1 Block Spin Transformations

Kadanoff's block spin construction is the original formulation of the renormalization group idea in statistical physics. Consider an Ising model on a lattice with spacing a. Group the spins into b^d blocks (where $b > 1$ and d is the dimension), replace each block by a single effective spin, and rescale the lattice so it has the same spacing a. The result is a new Ising-like system with different effective couplings.

This transformation—block, replace, rescale—is the RG step. Under repeated application it generates a flow in the space of Ising parameters (temperature, magnetic field, next-nearest-neighbor coupling, etc.). A fixed point of this flow is a critical point where the system looks the same at all scales.

Key Idea. Coarse-graining removes microscopic degrees of freedom and generates an effective theory for the remaining coarse-grained degrees of freedom. The RG transformation is a controlled way to perform this removal, keeping track of how the effective parameters change.

6.2 Emergent Laws

The renormalization group explains why the laws governing macroscopic phenomena look different from the laws governing microscopic physics. When we observe a fluid at scales much larger than the interatomic spacing, we see a continuum described by the Navier-Stokes equations, not by the Schrödinger equation for individual atoms. The Navier-Stokes equations are the effective theory obtained by coarse-graining over atomic scales.

This emergence of effective laws from underlying microscopic physics is not magical: it is a consequence of the RG flow. The effective theory at scale k is

described by the couplings $g_i(k)$ obtained by flowing the microscopic theory from the UV down to the scale k. As k decreases, irrelevant couplings flow toward their fixed-point values, reducing the number of independent parameters needed to describe the physics and giving the universal behavior characteristic of each universality class.

Exercise 6.1. Consider a one-dimensional chain of spins with nearest-neighbor coupling J and next-nearest-neighbor coupling J'. Under a block spin transformation with $b = 2$, the effective coupling J'_{eff} receives contributions from both J and J'. Write down the general form of the transformation $J'_{\text{eff}} = f(J, J')$ and argue why, for $|J'| \ll |J|$, the effective coupling satisfies $|J'_{\text{eff}}| < |J'|$ after many RG steps.

Part III

Renormalization Group Theory

Chapter 7

Theory Space

7.1 Coupling Constants as Coordinates

The space of all possible field theories consistent with a given set of fields and symmetries is called *theory space*. Each point in theory space corresponds to a specific set of coupling constants specifying the strength of every allowed interaction between the fields.

For a scalar field ϕ in four dimensions, a general Lagrangian consistent with the $\mathbb{Z}_2$ symmetry $\phi \to -\phi$ is

$$\mathcal{L} = \frac{1}{2}(\partial\phi)^2 + \frac{m^2}{2}\phi^2 + \frac{\lambda_4}{4!}\phi^4 + \frac{\lambda_6}{6!}\phi^6 + \frac{c_1}{M^2}(\partial^2\phi)^2 + \cdots \tag{7.1}$$

The coupling constants $(m^2, \lambda_4, \lambda_6, c_1, \ldots)$ are coordinates on theory space. This space is infinite-dimensional because there are infinitely many independent local operators consistent with the symmetry.

7.2 Effective Actions

The *effective action* $\Gamma[\phi]$ is the quantum generalization of the classical action. It is defined as the Legendre transform of the generating functional of connected correlation functions, and it contains the full quantum information about the theory:

$$\Gamma[\phi_{\text{cl}}] = \sup_J \left[J \cdot \phi_{\text{cl}} - W[J]\right], \tag{7.2}$$

where $W[J] = \ln Z[J]$ is the Schwinger functional and $\phi_{\text{cl}} = \delta W/\delta J$.

The effective action encodes the 1PI (one-particle-irreducible) correlation

functions. It can be written in a derivative expansion

$$\Gamma[\phi] = \int d^4x \left[V(\phi) + \frac{Z(\phi)}{2}(\partial\phi)^2 + \mathcal{O}(\partial^4) \right], \tag{7.3}$$

where $V(\phi)$ is the effective potential and $Z(\phi)$ is the wave-function renormalization.

Chapter 8

Beta Functions

8.1 Running Couplings and the Callan-Symanzik Equation

In a renormalized quantum field theory, the physical predictions must be independent of the arbitrary renormalization scale μ at which coupling constants are defined. This requirement leads to the Callan-Symanzik equation

$$\left[\mu\frac{\partial}{\partial\mu} + \beta_i(g)\frac{\partial}{\partial g_i} - n\gamma(g)\right]\Gamma^{(n)} = 0, \tag{8.1}$$

where $\Gamma^{(n)}$ is the n-point Green function, γ is the anomalous dimension, and the beta function is

$$\beta_i(g) = \mu\frac{\partial g_i}{\partial\mu}\bigg|_{g_0,\Lambda\text{ fixed}}. \tag{8.2}$$

Example 8.1. In ϕ^4 theory in four dimensions, the one-loop beta function for the quartic coupling is $\beta_\lambda = \frac{3\lambda^2}{16\pi^2} + \mathcal{O}(\lambda^3)$. Since $\beta_\lambda > 0$ for $\lambda > 0$, the coupling grows as the scale increases, eventually reaching a Landau pole. This is the triviality problem: the theory is either free ($\lambda = 0$) or ill-defined at high energies.

8.2 Renormalization Group Equations

The full RG equation for a dimensionless coupling g_i in the Wilsonian framework, where the scale is the UV cutoff k, is

$$k\frac{\partial g_i}{\partial k} = \beta_i(g_1, g_2, \ldots). \tag{8.3}$$

This system of coupled ordinary differential equations defines a flow on theory space. The structure of this flow—its fixed points, the stability of those fixed

points, and the basin of attraction of each fixed point— determines the universal properties of the theory.

Exercise 8.1. The beta function for a dimensionless coupling g in a toy model is $\beta(g) = -g + g^2$. Find all fixed points, determine their stability, and sketch the flow. What happens to a trajectory that starts at $g_0 = 0.5$?

Chapter 9

Fixed Points and Scale Invariance

9.1 Gaussian and Non-Gaussian Fixed Points

A *Gaussian fixed point* is one where all interactions vanish: the theory is free. At the Gaussian fixed point, the beta functions vanish trivially, $\beta_i = 0$ for all i, simply because there are no interactions to produce quantum corrections.

A *non-Gaussian fixed point* (NGFP) is an interacting fixed point where all beta functions vanish but the couplings are non-zero. Such a fixed point represents a scale-invariant interacting theory.

Definition 9.1. A theory is *scale invariant* if it is invariant under the transformation $x^\mu \to \lambda x^\mu$, $\phi(x) \to \lambda^{-\Delta}\phi(\lambda x)$ for some scaling dimension Δ. Under the RG, a scale-invariant theory corresponds to a fixed point.

9.2 Critical Surfaces and Predictivity

The UV critical surface $\mathcal{S}_{\text{UV}}$ is the set of all points in theory space that flow toward the fixed point as the scale $k \to \infty$. Its codimension equals the number of irrelevant directions at the fixed point.

A theory lying on $\mathcal{S}_{\text{UV}}$ is UV complete: it makes sense at arbitrarily high energies. A theory not on $\mathcal{S}_{\text{UV}}$ eventually diverges or loses predictivity above some finite energy scale. The dimension of $\mathcal{S}_{\text{UV}}$ determines the number of free parameters that must be measured to specify the theory—the predictive content of the UV completion.

Chapter 10

Relevant and Irrelevant Operators

10.1 Critical Exponents

Linearizing the RG flow around a fixed point g_i^* gives

$$\beta_i(g) \approx B_{ij}(g_j - g_j^*), \quad B_{ij} = \left.\frac{\partial \beta_i}{\partial g_j}\right|_{g^*}. \tag{10.1}$$

The eigenvalues $-\theta_I$ of B are the critical exponents. Directions with $\theta_I > 0$ (positive exponents) are relevant: perturbations in these directions grow as k decreases and must be measured experimentally. Directions with $\theta_I < 0$ are irrelevant: perturbations die away as k decreases and the coupling flows back to its fixed-point value.

10.2 Predictivity of Renormalizable Theories

In perturbative quantum field theory, renormalizability requires that only operators with non-negative canonical dimension appear in the Lagrangian. This is equivalent to requiring that all couplings at the Gaussian fixed point are either relevant or marginal—which holds precisely for operators with $[\mathcal{O}] \leq 4$ in four dimensions. Operators with $[\mathcal{O}] > 4$ have couplings with negative mass dimension (irrelevant at the Gaussian fixed point) and would require infinitely many counterterms, destroying predictivity.

Asymptotic safety generalizes this: UV completeness requires a non-Gaussian fixed point with a finite-dimensional UV critical surface. Perturbative renormalizability is the special case where the fixed point is Gaussian and only a finite set of operators is relevant.

Exercise 10.1. In four dimensions, classify the following operators as relevant, marginal, or irrelevant at the Gaussian fixed point: ϕ^2, ϕ^4, ϕ^6, $(\partial\phi)^2$, $(\partial^2\phi)^2$,

$\phi^2(\partial\phi)^2$.

Part IV

Quantum Field Theory Essentials

Chapter 11

Classical Field Theory

11.1 Action Principles and Lagrangians

The dynamics of a classical field $\phi(x)$ are determined by the action

$$S[\phi] = \int d^4x \, \mathcal{L}(\phi, \partial_\mu \phi), \tag{11.1}$$

where $\mathcal{L}$ is the Lagrangian density. The principle of stationary action—$\delta S = 0$ for all variations $\delta\phi$ vanishing at the boundary—yields the Euler-Lagrange equations:

$$\frac{\partial \mathcal{L}}{\partial \phi} - \partial_\mu \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} = 0. \tag{11.2}$$

For the scalar field with $\mathcal{L} = \frac{1}{2}(\partial\phi)^2 - V(\phi)$, the Euler-Lagrange equations give $\Box\phi + V'(\phi) = 0$.

11.2 Symmetries and Noether's Theorem

A symmetry of the action is a transformation of the fields that leaves S invariant. Noether's theorem states that every continuous symmetry generates a conserved current j^μ satisfying $\partial_\mu j^\mu = 0$, and hence a conserved charge $Q = \int d^3x \, j^0$.

Example 11.1. The Klein-Gordon action is invariant under translations $x^\mu \to x^\mu + \epsilon^\mu$. The Noether current is the energy-momentum tensor $T^{\mu\nu} = \partial^\mu\phi \, \partial^\nu\phi - \eta^{\mu\nu}\mathcal{L}$, with conserved charges $P^\mu = \int d^3x \, T^{0\mu}$.

11.3 The Einstein-Hilbert Action

Gravity is described by the curvature of spacetime. The metric tensor $g_{\mu\nu}(x)$ is the fundamental field, and its dynamics are governed by the Einstein-Hilbert

action:

$$S_{\text{EH}} = \frac{M_{\text{Pl}}^2}{2} \int d^4x \sqrt{-g}\,(R - 2\Lambda), \tag{11.3}$$

where R is the Ricci scalar curvature, Λ is the cosmological constant, $M_{\text{Pl}} = (8\pi G_N)^{-1/2}$ is the reduced Planck mass, and $\sqrt{-g}$ is the square root of the absolute value of the metric determinant. Varying with respect to $g^{\mu\nu}$ gives the Einstein field equations.

The coupling constant for gravity is Newton's constant G_N, which has mass dimension $[G_N] = -2$ in four dimensions. This negative mass dimension is the root of perturbative non-renormalizability: at each loop order, new operators of higher dimension are generated, requiring infinitely many counterterms.

Chapter 12

The Path Integral and Quantization

12.1 Path Integral Formulation

The quantum theory of a field is most naturally formulated through Feynman's path integral. The partition function is

$$Z = \int \mathcal{D}\phi \, e^{-S_E[\phi]}, \tag{12.1}$$

where the integral is over all field configurations $\phi(x)$ and S_E is the Euclidean action (obtained by the Wick rotation $t \to -i\tau$). The integrand e^{-S_E} weights each field configuration by a Boltzmann-like factor; configurations with lower action are more probable.

This statistical mechanics analogy is fundamental. The path integral for a quantum field theory in d Euclidean dimensions is formally equivalent to the partition function of a classical statistical system in d dimensions. The temperature of the statistical system corresponds to$\hbar$ in the quantum system. This equivalence makes renormalization group methods developed in statistical physics directly applicable to quantum field theory.

12.2 Perturbation Theory and Feynman Diagrams

When interactions are weak (small coupling constants), the path integral can be evaluated perturbatively by expanding in powers of the coupling. The terms in this expansion correspond to Feynman diagrams: graphical representations of scattering amplitudes.

At one-loop order, diagrams involve a single closed loop of internal propagators. The loop integral is over all momenta flowing around the loop, and it typically diverges for large momenta—this is the ultraviolet divergence of quan-

tum field theory. Renormalization provides a systematic procedure for absorbing these divergences into redefinitions of the coupling constants, yielding finite physical predictions.

Exercise 12.1. The free scalar field propagator in four-dimensional Euclidean space is $G(p) = (p^2 + m^2)^{-1}$. Show that the two-point function at one loop, $\int d^4p/(2\pi)^4\, G(p)$, diverges quadratically for large p.

Chapter 13

Renormalization

13.1 Regularization and Renormalized Parameters

To handle UV divergences, one introduces a regularization procedure that makes loop integrals finite by modifying the theory at high energies. Dimensional regularization, which analytically continues the spacetime dimension from 4 to $4-\epsilon$, is the most elegant: it preserves all symmetries and introduces a mass scale μ through the relation

$$\int \frac{d^d p}{(2\pi)^d} \to \mu^\epsilon \int \frac{d^d p}{(2\pi)^d}. \tag{13.1}$$

Divergences appear as poles in ϵ.

Renormalization replaces the bare coupling g_0 with a renormalized coupling $g(\mu)$ defined by

$$g_0 = \mu^\epsilon Z_g(g,\mu)\, g(\mu), \tag{13.2}$$

where Z_g is the renormalization constant chosen to absorb the UV poles. The renormalized coupling $g(\mu)$ is the physical parameter measured at scale μ; it is finite and depends on the renormalization scale.

13.2 Effective Field Theories

The Wilsonian perspective on renormalization, which underlies the FRG, is more physical. One introduces a UV cutoff Λ and integrates out all field modes with momenta $p > \Lambda$, obtaining an effective action for the low-energy modes. Lowering Λ integrates out more modes and generates an effective theory at the new scale. This is the Wilsonian RG, and the effective action S_Λ depends on Λ in a way governed by the Wetterich equation.

The Wilsonian effective action has the same operator content as the original

action but with scale-dependent coefficients. At each scale Λ, the irrelevant operators are suppressed by inverse powers of Λ relative to the UV scale, justifying their truncation for low-energy calculations.

Part V

Functional Renormalization Group

Chapter 14

The Effective Average Action

14.1 Motivation and Construction

The functional renormalization group, developed by Wetterich, interpolates between the bare action S (at $k = \infty$) and the full quantum effective action Γ (at $k = 0$) by introducing an infrared regulator R_k that gives mass $\sim k^2$ to modes with momenta $p < k$:

$$Z_k[J] = \int \mathcal{D}\phi \, \exp\left(-S[\phi] - \tfrac{1}{2}\phi R_k \phi + J\phi\right). \tag{14.1}$$

The scale-dependent effective action Γ_k is defined through the modified Legendre transform

$$\Gamma_k[\phi] = \sup_J \left[J\phi - W_k[J]\right] - \tfrac{1}{2}\phi R_k \phi. \tag{14.2}$$

By construction, $\Gamma_{k\to\infty} \to S$ (the bare action) and $\Gamma_{k\to 0} \to \Gamma$ (the full effective action).

14.2 The Wetterich Equation

Differentiating Γ_k with respect to $t = \ln k$ yields the exact flow equation:

$$\partial_t \Gamma_k = \frac{1}{2}\mathrm{Tr}\left[(\Gamma_k^{(2)} + R_k)^{-1}\partial_t R_k\right]. \tag{14.3}$$

This is the Wetterich equation. It has the structure of a one-loop expression but is exact: the full propagator $(\Gamma_k^{(2)} + R_k)^{-1}$ includes all quantum corrections. The regulator insertion $\partial_t R_k$ is peaked at $p^2 \approx k^2$, so only modes near the current scale k contribute to the flow at each step.

Key Idea. The Wetterich equation describes a continuous integration of quantum fluctuations, one momentum shell at a time, as k flows from the UV to the IR. It is formally exact and reduces to known perturbative results when expanded in powers of the coupling.

14.3 Truncation Methods

The Wetterich equation is a functional differential equation on the infinite-dimensional space of all functionals $\Gamma_k[\phi]$. To extract physics one truncates to a finite-dimensional subspace.

The simplest truncation is the *local potential approximation* (LPA), which sets

$$\Gamma_k[\phi] = \int d^4x \left[\frac{1}{2}(\partial\phi)^2 + V_k(\phi)\right], \tag{14.4}$$

tracking only the running of the effective potential $V_k(\phi)$ while keeping the kinetic term fixed. The flow equation for V_k becomes a PDE in the field variable ϕ that can be solved numerically or expanded in polynomials.

More sophisticated truncations include the *derivative expansion* (retaining $(\partial\phi)^2$, $(\partial^2\phi)^2$, etc.) and *vertex expansion* (expanding in powers of the field around a fixed background). The quality of the truncation is assessed by checking stability under enlargement.

Chapter 15

Flow Equations and Fixed-Point Search

15.1 Beta Functions from the Flow Equation

Given a truncation $\Gamma_k = \sum_i g_i(k) \int \mathcal{O}_i$, the beta functions $\beta_i = \partial_t g_i$ are obtained by:

(i) inserting the truncated action into the Wetterich equation (14.3); (ii) expanding the right-hand side using the $P^{-1}F$ expansion; (iii) evaluating the functional traces with the Litim regulator $R_k(p^2) = (k^2 - p^2)\Theta(k^2 - p^2)$; (iv) projecting the result onto the operators $\mathcal{O}_i$ to read off the coefficients β_i.

This procedure generates explicit analytic expressions for the beta functions that, while approximate due to the truncation, capture the non-perturbative physics of fixed points and their neighborhoods.

15.2 Fixed Points and Critical Exponents

To find fixed points one solves $\beta_i(g^*) = 0$ simultaneously for all couplings in the truncation. The stability matrix $B_{ij} = \partial\beta_i/\partial g_j|_{g^*}$ is then diagonalized to find the critical exponents $\theta_I = -\text{eig}(B)$.

Positive θ_I indicate relevant directions; negative θ_I indicate irrelevant directions. The number of positive critical exponents equals the dimension of the UV critical surface and hence the number of free parameters of the UV-complete theory.

For pure Einstein-Hilbert gravity truncation with couplings ($g = Gk^2, \lambda = \Lambda/k^2$), the non-Gaussian fixed point is found at $g_* \approx 0.7$, $\lambda_* \approx 0.19$ (regulator and gauge dependent), with two relevant directions and critical exponents $\theta_{1,2} \approx -1.5 \pm 3.6i$ (complex conjugate pair indicating a UV spiral).

Exercise 15.1. For the beta functions $\beta_g = 2g - bg^2/(1-\lambda)^2$ and $\beta_\lambda = -2\lambda + cg/(1-\lambda)$ with constants $b, c > 0$, find the non-Gaussian fixed point (g_*, λ_*) and compute the stability matrix. Show that the two eigenvalues are complex conjugates.

Part VI

Gravity and Asymptotic Safety

Chapter 16

Gravity as a Field Theory

16.1 The Metric as a Dynamical Field

In general relativity, spacetime geometry is itself dynamical. The metric $g_{\mu\nu}(x)$ is a tensor field that describes the distances between nearby points: $ds^2 = g_{\mu\nu}dx^\mu dx^\nu$. The metric satisfies the Einstein field equations

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{1}{M_{\rm Pl}^2} T_{\mu\nu}, \tag{16.1}$$

where $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R$ is the Einstein tensor, $R_{\mu\nu}$ is the Ricci tensor, R is the Ricci scalar, and $T_{\mu\nu}$ is the energy-momentum tensor of matter.

16.2 Perturbative Non-Renormalizability

When gravity is treated as a quantum field theory, the metric is expanded around flat spacetime: $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}/M_{\rm Pl}$. The graviton field $h_{\mu\nu}$ propagates and interacts through vertices derived from the Einstein-Hilbert action. At each loop order, power-counting shows that new divergences appear that require new counterterms. For instance, at one loop the counterterm is $\sim R^2$ (schematically), at two loops $\sim R^3$, and so on.

This is perturbative non-renormalizability: the theory has infinitely many independent divergences requiring infinitely many free parameters. Newton's constant G_N has $[G_N] = -2$ (negative mass dimension), so higher-loop diagrams are increasingly divergent. Standard perturbation theory fails as a UV completion.

Chapter 17

The Asymptotic Safety Program

17.1 Weinberg's Proposal

Weinberg proposed in 1979 that the UV divergences of perturbative gravity might be controlled by a non-Gaussian fixed point of the gravitational RG flow. If such a fixed point exists with a finite-dimensional UV critical surface, then gravity would be UV complete without requiring new degrees of freedom: it would be *asymptotically safe*.

The key difference from perturbative renormalizability is that the fixed point is not Gaussian (non-interacting) but non-Gaussian (interacting). The theory does not become free at high energies but approaches a scale-invariant interacting theory. The gravitational coupling $g = Gk^2$ remains finite at the fixed point, and Newton's constant $G_N = g_*/k^2$ vanishes as $k \to \infty$ (antiscreening of the gravitational interaction at the fixed point scale).

17.2 Evidence for the Non-Gaussian Fixed Point

The primary evidence for the gravitational NGFP comes from FRG calculations in progressively larger truncations of the gravitational effective action. Starting with the Einstein-Hilbert truncation (Reuter, 1998), the fixed point has been confirmed in truncations including R^2, R^3, higher-order curvature invariants, and matter fields. The fixed-point values and critical exponents are quantitatively truncation-dependent but qualitatively robust.

Key features are: a UV attractive fixed point at finite (g_*, λ_*), two relevant directions (one for G and one for Λ), and an irrelevant direction for every higher-derivative operator, consistent with the prediction that the asymptotically safe theory has only two free parameters (corresponding to Newton's constant and the cosmological constant as IR observables).

17.3 Matter-Gravity Fixed Points and the Weak-Gravity Bound

When matter fields are coupled to the gravitational sector, the fixed-point analysis must be performed for the full gravity-matter system. Gravitational fluctuations modify the matter beta functions, and matter fluctuations modify the gravitational beta functions. The resulting fixed-point structure constrains which combinations of matter content are compatible with asymptotic safety.

The weak-gravity bound emerges from this analysis: as Newton's coupling g increases, interacting matter fixed points eventually collide and disappear, leaving the matter sector without a UV completion. The bound $g < g_{\text{crit}} \approx 1.4$ (at $\lambda = 0$) must be satisfied for gravity-matter UV completion to exist. This bound translates into constraints on the allowed matter content and coupling structure of any UV-complete model.

Chapter 18

Matter and Gravity

18.1 Scalar Couplings and the Higgs Mass Prediction

The most striking quantitative success of asymptotically safe gravity is the prediction of the Higgs boson mass. The quartic self-coupling λ_{H} of the Higgs field is irrelevant at the gravitational NGFP, meaning that gravity drives it toward a specific fixed-point value at the Planck scale. Running this value from M_{Pl} to the electroweak scale via the Standard Model RG equations yields a Higgs mass prediction of $M_H \approx 126\,\mathrm{GeV}$, in excellent agreement with the measured value of $125.25\,\mathrm{GeV}$.

This prediction is a concrete demonstration of the predictive power of the asymptotic-safety scenario: an infrared observable (the Higgs mass) is determined by the UV fixed-point structure without any additional free parameter.

18.2 Fermions and Gauge Fields

Fermionic and gauge fields present interesting interplay with gravity. The $U(1)$ gauge coupling in the Standard Model has a Landau pole in the absence of gravity: it diverges at a finite energy scale around 10^{40} GeV. However, gravitational antiscreening can prevent this: if the gravitational contribution to the gauge beta function is sufficiently positive (antiscreening), the gauge coupling can be pushed toward a non-Gaussian fixed point, resolving the triviality problem. Whether this occurs depends on the relative strengths of gravitational and gauge fluctuations at the fixed point.

Similarly, Yukawa couplings between fermions and scalars receive gravitational corrections that can shift their fixed-point values. In the Standard Model context, these corrections modify the prediction for fermion masses and mixing

angles, providing additional UV-sensitive predictions that in principle distinguish asymptotic safety from alternative UV completions.

Part VII

Cosmology and Observational Signatures

Chapter 19

Running Couplings in Cosmology

19.1 RG-Improved Friedmann Equations

In standard cosmology, Newton's constant and the cosmological constant are taken as fixed. Asymptotic safety suggests a more nuanced picture: as the universe evolves from early high-energy epochs to the present low-energy era, the effective values of these parameters follow their RG trajectories.

The RG-improved Friedmann equations incorporate this scale dependence by identifying the RG scale k with a physically relevant energy scale of the cosmological system—typically the Hubble parameter H or the inverse of the horizon size. The dimensionless Newton coupling $g(k) = G(k)k^2$ runs according to its beta function, and the physical Newton constant is $G(k) = g(k)/k^2$.

Near the NGFP, where $g(k) \approx g_*$, Newton's constant scales as $G(k) \approx g_*/k^2 \propto 1/k^2$, becoming very small at high energies. This gives a natural resolution of the big bang singularity: as $k \to \infty$ (early universe), $G \to 0$ and the gravitational coupling weakens, potentially smoothing the singularity.

19.2 Cosmological Fixed Points

The RG improvement of the cosmological equations introduces the possibility of a purely gravitational fixed-point cosmology. When the Hubble parameter H satisfies $H \sim k \sim M_{\text{Pl}}$, the dimensionless coupling $g = GH^2 \approx g_*$ is near the NGFP. The cosmological evolution near this point can be analyzed using the beta functions of the gravitational sector.

Near the NGFP the effective dark energy density (from the running cosmological constant) scales as $\rho_\Lambda(k) \approx \Lambda(k)M_{\text{Pl}}^2 \approx \lambda_* k^2 M_{\text{Pl}}^2$, which drives de Sitter-like inflation at the Planck epoch. This provides a natural mechanism for an early inflationary epoch without requiring a separate inflaton field, though the

detailed dynamics depend on the trajectory chosen in the RG-improved cosmological equations.

Chapter 20

Observational Windows

20.1 Gravitational Waves as Quantum Gravity Probes

Gravitational wave observations provide a unique window into cosmological physics at energy scales inaccessible to collider experiments. The stochastic gravitational-wave background, generated by various processes in the early universe, encodes information about phase transitions, cosmic strings, and the expansion history of the universe.

Asymptotic safety makes specific predictions for the gravitational-wave background through its constraints on cosmic string models and on the inflationary sector. The NANOGrav 15-year dataset, which detected a stochastic background in the nanohertz band, has begun to test these predictions: the data disfavor simple field-theoretic cosmic strings, consistent with the asymptotic-safety requirement that such strings cannot arise from a minimal UV-complete model.

20.2 The CMB and Large-Scale Structure

The cosmic microwave background provides a snapshot of the universe at recombination ($z \approx 1100$), encoding information about primordial perturbations generated during inflation. Asymptotic safety modifies inflationary dynamics through the running Newton constant and cosmological constant, potentially leaving imprints on the CMB power spectrum—particularly on the spectral index n_s and the tensor-to-scalar ratio r.

Large-scale structure formation is sensitive to the growth rate of perturbations, which depends on the effective strength of gravity at galactic scales. The σ_8 tension between CMB-derived and weak-lensing-derived values of the matter clustering amplitude may be addressed by a scale-dependent Newton constant,

though the detailed calculation requires connecting the RG improvement to the full perturbation theory of structure formation.

Part VIII

Mathematical Structures

Chapter 21

Metric Geometry and Theory Space

21.1 Distances Between Theories

Gromov-Hausdorff convergence provides a precise mathematical notion of convergence for sequences of metric spaces. Given compact metric spaces (X, d_X) and (Y, d_Y), their Gromov-Hausdorff distance is

$$d_{\mathrm{GH}}(X, Y) = \inf_{Z,f,g} d_H(f(X), g(Y)), \tag{21.1}$$

where the infimum is over all metric spaces Z and isometric embeddings $f : X \to Z$, $g : Y \to Z$, and d_H is the Hausdorff distance.

In the context of renormalization group theory, this machinery applies to the sequence of effective theory spaces at different scales. As k decreases, the effective theory space can be thought of as converging (in the Gromov-Hausdorff sense) toward the fixed-point theory space—a metric space of lower complexity that describes only the universal long-distance physics.

Chapter 22

Information Geometry

22.1 The Fisher Information Metric

The Fisher information matrix provides a natural Riemannian metric on the manifold of probability distributions. For a family of distributions $P(x;\theta)$ parametrized by $\theta = (\theta^1, \ldots, \theta^n)$:

$$G_{ij}(\theta) = \int dx\, P(x;\theta)\, \partial_i \ln P(x;\theta)\, \partial_j \ln P(x;\theta). \tag{22.1}$$

This metric measures the distinguishability between nearby distributions: two distributions at Fisher distance ds can be distinguished (on average) only if the number of samples $N \gtrsim (ds)^{-2}$.

When the parameter space is identified with theory space and the coupling constants play the role of statistical parameters, the Fisher metric defines a natural notion of distance between theories. RG trajectories can then be interpreted as geodesics of this metric, moving in the direction that maximally compresses information about the microscopic physics while preserving the macroscopic observables.

22.2 Thermodynamic Geometry

Ruppeiner's thermodynamic geometry applies the Fisher metric to the space of thermodynamic states, with the entropy S playing the role of the log-probability. The metric is

$$g^R_{ij} = -\frac{\partial^2 S}{\partial X^i \partial X^j}\bigg|_{\text{eq}}, \tag{22.2}$$

where X^i are extensive thermodynamic variables. The curvature of this metric encodes information about interactions: an ideal gas has flat thermodynamic

geometry (no interactions), while a van der Waals gas has non-trivial curvature near the liquid-gas critical point.

The thermodynamic geometry of a field theory at a critical point is related to the critical exponents: the correlation length diverges at criticality, and the Ruppeiner metric develops a singularity that can be characterized by the critical exponents ν and γ.

Chapter 23

Category Theory and Physical Structure

23.1 Functors and Natural Transformations

Category theory provides a language for describing mathematical structures and their relationships at a high level of abstraction. A *category* $\mathcal{C}$ consists of objects and morphisms (arrows between objects), with composition of morphisms satisfying associativity and identity axioms.

A *functor* $F : \mathcal{C} \to \mathcal{D}$ is a structure-preserving map between categories: it sends objects to objects and morphisms to morphisms, respecting composition. Functors formalize the idea of translation between mathematical structures.

In the RG context, the coarse-graining transformation at each scale step can be formalized as a functor from the category of UV effective theories to the category of IR effective theories. The functoriality condition $\mathcal{R}(f \circ g) = \mathcal{R}(f) \circ \mathcal{R}(g)$ corresponds to the semigroup property of RG transformations: applying two successive coarse-graining steps gives the same result as applying a single coarse-graining step by the combined amount.

23.2 Sheaves and Local-to-Global Principles

A sheaf is a mathematical structure that allows local data defined on open sets to be patched together consistently to give global data. In the FRG context, the effective action Γ_k defines a sheaf over the space of scales: the local data (beta functions, fixed-point values) at each scale k must be consistent with the global RG trajectory.

The obstruction theory associated with sheaves—measuring the failure of local data to extend globally—appears in the FRG context when one asks whether

a specific UV fixed point can be connected to a specific IR physics by an RG trajectory. If a topological obstruction prevents this, the theory is excluded regardless of its local (perturbative) consistency.

Part IX

Computational Experiments

Chapter 24

Numerical Renormalization Group

24.1 Integrating Beta Functions

The most direct computational experiment in RG theory is the numerical integration of a system of coupled beta-function ODEs:

$$\frac{dg_i}{dt} = \beta_i(g_1, \ldots, g_n), \quad t = \ln(k/k_0). \tag{24.1}$$

Given initial conditions $(g_i(t_0))$ representing a UV initial theory, one integrates forward in t (decreasing scale) to obtain the IR theory.

A simple implementation uses a fourth-order Runge-Kutta method:

$$g_i(t + \Delta t) = g_i(t) + \frac{\Delta t}{6}(k_1 + 2k_2 + 2k_3 + k_4), \tag{24.2}$$

where $k_1 = \beta_i(g(t))$, $k_2 = \beta_i(g(t) + \frac{\Delta t}{2}k_1)$, etc. Fixed points are identified as regions where $|dg_i/dt| < \epsilon$ for all i, and critical exponents are computed by diagonalizing the numerical Jacobian matrix at the fixed point.

24.2 Visualizing RG Flows

For two-coupling systems, the phase portrait of the RG flow can be plotted directly as a vector field in the (g_1, g_2) plane. Each arrow at a point (g_1, g_2) points in the direction (β_1, β_2). Fixed points are zeros of this vector field, saddle points are intersections of separatrices, and basins of attraction are the regions from which trajectories flow toward a given attractor.

For the toy model $\beta_g = 2g - g^2$, $\beta_\lambda = -2\lambda + g\lambda$, the phase portrait reveals a Gaussian fixed point at $(0, 0)$, a NGFP at (g_*, λ_*), and separatrices dividing the plane into basins of attraction. The UV critical surface is the line (in this

two-dimensional case) of trajectories that flow toward the NGFP as $t \to +\infty$.

Exercise 24.1. Write pseudocode for a Runge-Kutta integrator that takes beta functions $\beta_g(g, \lambda) = 2g - bg^2/(1 - \lambda)$ and $\beta_\lambda = -2\lambda + cg/(1 - \lambda)$ as input and produces a phase portrait for $0 < g < 2$, $-0.5 < \lambda < 0.5$. Identify the non-Gaussian fixed point and estimate its coordinates numerically.

Chapter 25

Simulation Projects

25.1 Scalar Field Renormalization Group

The simplest non-trivial RG calculation is the flow of the effective potential for a $\mathbb{Z}_2$-symmetric scalar field in the local potential approximation. The flow equation for the dimensionless potential $u(\rho; t) = V_k(k^{d/2-1}\phi)/k^d$, where $\rho = \phi^2/2$, is a PDE:

$$\partial_t u = -du + \frac{d-2+\eta}{2}\rho\,\partial_\rho u + \frac{4v_d}{d}\frac{(d-1)l_0^d(u') + l_0^d(u' + 2\rho u'')}{(1+u')^{\cdots}}, \tag{25.1}$$

where v_d is a phase-space volume, l_0^d are threshold functions from the Litim regulator, and primes denote derivatives with respect to ρ. Solving this PDE by discretizing in ρ and using the Runge-Kutta method in t produces the running potential, from which one can extract the critical exponents ν and η of the Ising universality class.

25.2 Gravity-Inspired Flows

A toy model of the gravitational RG flow uses the truncated beta functions from the Einstein-Hilbert sector:

$$\beta_g = (2+\eta_N)g, \quad \beta_\lambda = -(2-\eta_N)\lambda + (\text{graviton loop terms}), \tag{25.2}$$

where $\eta_N = -2Bg^2/(1-2\lambda)^2$ is the graviton anomalous dimension and B is a numerical constant depending on the regulator. The NGFP and its critical exponents can be found numerically for any $B > 0$. One finds that for a wide range of B (corresponding to different regulator choices), the fixed point exists with positive coordinates (g_*, λ_*) and complex conjugate critical exponents.

Part X

Beyond Conventional Renormalization

Chapter 26

Renormalization as Geometric Flow

26.1 Theory Space as a Manifold

Having established the technical machinery of the FRG, we now step back and examine the RG flow from a more geometric perspective. Theory space, equipped with the Fisher information metric G_{ij} of Chapter 22, becomes a Riemannian manifold. The coupling constants g_i serve as coordinates on this manifold, and the RG flow is a vector field $\beta^i = k\partial_k g^i$ on it.

This geometric interpretation is more than a metaphor. The metric structure on theory space has physical content: it measures the statistical distinguishability of nearby theories, which is related to the sensitivity of physical observables to variations in the coupling constants. A region of theory space with large metric components is one where the observables change rapidly with the couplings; a flat region is one where the observables are insensitive to the microscopic parameters.

26.2 Gradient Flows and the c-Theorem

Zamolodchikov's c-theorem in two-dimensional QFT states that there exists a function $c(g)$ on theory space that decreases monotonically along RG trajectories: $\partial_t c \leq 0$, with equality only at fixed points. The function c counts the degrees of freedom in a scale-invariant way and provides a Lyapunov function for the RG flow.

In two dimensions, $c(g)$ is the central charge of the Virasoro algebra. In four dimensions, the conjectured a-theorem (proven by Komargodski and Schwimmer) states that the a-anomaly coefficient decreases along RG flows: $a_{\text{UV}} \geq a_{\text{IR}}$.

The existence of such a monotone function makes the RG flow a *gradient flow* in an appropriate metric on theory space. This is deeply connected to the

information-geometric interpretation: the decrease of c (or a) corresponds to the loss of information about the UV through coarse-graining.

26.3 Entropy Interpretations of RG Flow

The connection between entropy and the RG flow is most transparent in the framework of relative entropy (Kullback-Leibler divergence). Given two theories described by probability distributions P and Q for observables, the relative entropy $D_{\mathrm{KL}}(P\|Q) = \int P \ln(P/Q)$ measures how much information is lost when Q is used to approximate P.

Under a coarse-graining step, the relative entropy between the coarse-grained theory and the original theory is non-negative by the data-processing inequality (a quantum information theorem). This gives a precise, non-perturbative proof that information is lost monotonically under coarse-graining, consistent with the c-theorem and with the interpretation of the RG as an entropy-generating process.

Chapter 27

Information Geometry of Field Theories

27.1 Statistical Manifolds of Theories

The Fisher information metric endows theory space with a geometric structure. For a field theory with partition function $Z[J] = \int \mathcal{D}\phi\, e^{-S[\phi]+J\phi}$, the coupling constants $\{g_i\}$ play the role of statistical parameters. The Fisher metric is

$$G_{ij}(g) = \frac{\partial^2}{\partial g_i \partial g_j} \ln Z = \langle \mathcal{O}_i \mathcal{O}_j \rangle_c, \tag{27.1}$$

where $\langle \mathcal{O}_i \mathcal{O}_j \rangle_c$ is the connected two-point function of the operators coupled to g_i and g_j.

Definition 27.1. The *statistical manifold* of a field theory is theory space equipped with the Fisher information metric. The geodesics of this metric are the "natural" paths between theories in the sense that they minimize the total information loss along the path.

At a fixed point, the Fisher metric takes a universal form determined by the operator algebra of the CFT. The critical exponents are eigenvalues of the metric's curvature tensor, and the universality classes of the RG correspond to the isometry classes of the fixed-point metric.

27.2 Thermodynamic Curvature and Criticality

Ruppeiner's thermodynamic geometry assigns curvature to the space of thermodynamic states, with the curvature scalar $\mathcal{R}$ serving as a measure of interaction strength. For an ideal gas $\mathcal{R} = 0$; for an interacting gas $\mathcal{R} \neq 0$, and $\mathcal{R}$ diverges at the critical point.

This divergence is controlled by the correlation length ξ: the curvature scales as $\mathcal{R} \sim \xi^d$ near criticality, where d is the dimension. This means that thermodynamic curvature probes the same physics as correlation functions—it is another window into the fixed-point structure of the RG.

In the FRG context, the running of the Fisher metric along the RG trajectory provides a geometric picture of how the effective theory becomes simpler as scale decreases: the metric curvature decreases, the metric becomes flatter, and the effective theory approaches the scale-invariant fixed-point metric.

Chapter 28

Coarse-Graining as Entropy Transport

28.1 Entropy Production under Coarse-Graining

Each step of the RG coarse-graining procedure integrates out a thin shell of high-momentum modes. The information contained in these modes is lost—it cannot be recovered from the coarse-grained description. This irreversible loss of information is precisely entropy production.

More formally: define the von Neumann entropy of the quantum state in the UV sector $S_{\text{UV}} = -\text{Tr}[\rho_{\text{UV}} \ln \rho_{\text{UV}}]$. After tracing out the high-momentum modes to obtain the reduced state ρ_k, the entropy of the coarse-grained state satisfies $S(\rho_k) \geq S(\rho_{\text{UV}})$ by the monotonicity of relative entropy under quantum channels. Entropy increases monotonically under coarse-graining.

28.2 The Arrow of Scale

The monotonic increase of entropy under coarse-graining defines what might be called an "arrow of scale": just as the arrow of time is defined by the direction of entropy increase under temporal evolution, the arrow of scale is defined by the direction of information loss under the RG transformation. Flowing from UV to IR (decreasing k) is analogous to flowing from past to future (increasing t): both processes are irreversible in the sense that the coarse-grained description cannot be recovered from the fine-grained one.

This analogy is more than poetic. In the context of holographic duality, there is a precise correspondence between the RG flow in the boundary theory and the radial direction in the bulk geometry. The arrow of scale in the boundary theory corresponds to the arrow of depth in the bulk, and the monotonic de-

crease of information corresponds to the monotonic decrease of the holographic entanglement entropy as the bulk depth increases.

Part XI

Scalar-Vector Plenum Framework

Chapter 29

Motivation for Plenum Models

29.1 Limitations of Metric-Based Spacetime

General relativity describes gravity through the metric tensor $g_{\mu\nu}$. This is a remarkably successful framework, but it has conceptual limitations that become apparent when one attempts to extend it to the quantum regime or to cosmological initial conditions. The metric describes the *geometry* of spacetime but does not directly address the *content* of spacetime— the fields and fluctuations that fill it.

A plenum model begins from a different starting point: instead of taking the metric as the primary variable and deriving everything else from it, one takes a set of coexisting fields as primary and derives the geometric structure as an emergent property of their configuration. This is not merely a reformulation of general relativity but a genuinely different ontological stance: spacetime is not a container within which fields live, but an emergent description of the relational structure between field configurations.

29.2 Field-Based Ontology

The Relativistic Scalar-Vector Plenum (RSVP) framework takes this stance concretely. The fundamental variables are:

a scalar density field $\Phi(x)$, which represents the local energy density or information density of the plenum;

a vector flow field $\mathbf{v}(x)$, which represents the directed transport of this density through spacetime;

an entropy field $S(x)$, which represents the local degree of configurational freedom of the plenum.

These three fields together constitute a *plenum*: a physically realized medium

that fills spacetime and whose configuration encodes both the geometric and the thermodynamic properties of the physical situation.

Chapter 30

Scalar Fields as Density Structures

30.1 Scalar Densities in Physical Theories

Scalar fields appear throughout physics as density quantities: energy density $\rho(x,t)$, number density $n(x,t)$, temperature $T(x,t)$, electrostatic potential $\varphi(x,t)$. What distinguishes a scalar field from a mere function is its transformation law: a scalar field is invariant under Lorentz (or more generally, coordinate) transformations, assigning the same value to a physical event regardless of the coordinate system used.

In the plenum framework, the scalar density $\Phi(x)$ is the primary carrier of information about the physical state. Its gradient $\partial_\mu \Phi$ encodes the direction and rate of change of physical density. Regions where $|\nabla\Phi|$ is large are regions of rapid variation—either sharp transitions between different phases of the plenum or concentrated energy-density structures such as particles.

30.2 Energy Landscapes and Order Parameters

The scalar field in the plenum is closely related to the concept of an order parameter in the statistical physics discussion of Chapter 5. Just as the Ising magnetization m measures the degree of order in the spin system, $\Phi(x)$ measures the degree of organization of the plenum at each point.

The dynamics of Φ can be described by an effective potential $V(\Phi)$ that encodes the preferred configurations. A double-well potential $V(\Phi) = \lambda(\Phi^2 - v^2)^2$ favors $\Phi = \pm v$ over $\Phi = 0$, corresponding to a symmetry-broken phase where the plenum has spontaneously chosen a preferred configuration. The parameter v is the vacuum expectation value; for a cosmic string model it determines the string tension.

Chapter 31

Vector Flows and Directed Structure

31.1 Flux and Transport

The vector field $\mathbf{v}(x)$ in the plenum framework represents the directed flow of the scalar density. In fluid dynamics, a velocity field $\mathbf{u}(x,t)$ describes the local velocity of fluid elements; the continuity equation

$$\partial_t \rho + \nabla \cdot (\rho \mathbf{u}) = 0 \tag{31.1}$$

encodes the conservation of mass. The RSVP vector field plays an analogous role: it encodes the direction and rate of transport of the scalar density through the plenum.

The divergence $\nabla \cdot \mathbf{v}$ measures whether the flow is expanding or contracting at a given point. Regions with $\nabla \cdot \mathbf{v} > 0$ are sources of flow (density is being created or spreading outward); regions with $\nabla \cdot \mathbf{v} < 0$ are sinks.

31.2 Topological Defects and Vorticity

The curl $\nabla \times \mathbf{v}$ (in three dimensions) or the antisymmetric part of $\partial_\mu v_\nu$ (relativistically) measures the rotational component of the flow. Vortex lines—lines around which the flow circulates— are analogous to the topological defects (cosmic strings, vortices) of spontaneously broken field theories.

Indeed, in the abelian Higgs model for cosmic strings, the string is precisely a vortex in the phase of the complex scalar field $\phi = |\phi| e^{i\theta}$, around which $\oint d\theta = 2\pi n$ for integer winding number n. In the RSVP context, the vector field $\mathbf{v}$ can develop vortex structures analogous to these topological defects, and these structures carry conserved topological charges that prevent them from decaying smoothly.

Chapter 32

Entropy Fields

32.1 Entropy as a Dynamical Variable

In conventional thermodynamics, entropy is a state function that characterizes equilibrium states. In the RSVP framework, entropy is promoted to a dynamical field $S(x, t)$ that evolves according to its own equation of motion and couples to the scalar and vector fields.

This promotion of entropy from a state function to a field has several motivations. First, in non-equilibrium systems the entropy is genuinely spatially heterogeneous: different regions have different degrees of disorder. A field description captures this heterogeneity. Second, the RG interpretation developed in Part IX suggests that entropy measures the amount of microscopic information that has been integrated out at each scale; promoting it to a field allows this scale-dependent information to be tracked locally. Third, the entropy field provides a natural arrow of time: its increase under the dynamics encodes the second law of thermodynamics as a field equation rather than an external constraint.

32.2 Lamphrodyne Relaxation

The entropy field evolves through a process called lamphrodyne relaxation: high-entropy configurations relax toward lower-entropy configurations in a way governed by the gradient of the entropy field and the coupling to the scalar and vector fields. The evolution equation has the form

$$\partial_t S = D\nabla^2 S + \lambda|\nabla\Phi|^2 - \gamma S, \tag{32.1}$$

where D is the entropy diffusion coefficient, λ is the source coupling (entropy production from scalar gradients), and γ is the dissipation rate. This equation ensures that entropy flows from high-gradient regions (where energy is being deposited) to low-gradient regions (where the plenum is smooth), mimicking the second law of thermodynamics.

The term $\lambda|\nabla\Phi|^2$ is physically important: it couples entropy production to the kinetic energy of the scalar field. Regions where the scalar field varies rapidly (high $|\nabla\Phi|$) generate entropy. This is the RSVP analog of the dissipation of kinetic energy into heat in a viscous fluid.

Part XII

Field Coupling Structures

Chapter 33

Coupled Scalar-Vector Dynamics

33.1 Advection-Diffusion and the RSVP Action

The coupled evolution of Φ, $\mathbf{v}$, and S is derived from an action principle. The RSVP Lagrangian density takes the general form

$$\mathcal{L}_{\text{RSVP}} = \mathcal{L}_\Phi(\Phi, \partial_\mu \Phi) + \mathcal{L}_v(v^\mu, \partial_\mu v^\nu) + \mathcal{L}_S(S, \partial_\mu S) + \mathcal{L}_{\text{int}}(\Phi, v^\mu, S, \partial\Phi, \partial v, \partial S), \tag{33.1}$$

where the interaction term $\mathcal{L}_{\text{int}}$ encodes the couplings between the three fields. The Euler-Lagrange equations derived from this action govern the dynamics of the plenum.

The simplest coupling between the scalar and vector fields is an advection term: the scalar density is transported by the vector flow, $\partial_t \Phi + \mathbf{v} \cdot \nabla \Phi = 0$. This is the continuity equation for Φ in a flow field $\mathbf{v}$. Adding diffusion and a source term from the entropy field gives the complete evolution equation for Φ.

33.2 Instabilities and Pattern Formation

Coupled field systems exhibit a rich variety of dynamical instabilities and emergent pattern formation. In the RSVP framework, the interplay between advection (transport by $\mathbf{v}$), diffusion (spreading of Φ), and entropy production (source term from $|\nabla\Phi|^2$) can produce:

localized solitonic structures, where the scalar density Φ forms a stable, particle-like configuration balanced by dispersive and dissipative effects;

traveling waves, where energy is transported coherently across the plenum;

turbulent flows, where the vector field develops chaotic behavior on multiple scales, analogous to hydrodynamic turbulence.

These emergent structures are the RSVP analogs of the particles, radiation,

and matter distributions of standard cosmology.

Chapter 34

Lamphron and Lamphrodyne States

34.1 Definition of Lamphronic Regimes

In the RSVP framework, different regimes of the coupled field dynamics are characterized by the relative magnitudes of the kinetic, entropy, and interaction energy densities. A *lamphronic* state is one in which the entropy production and dissipation are in approximate balance: the system is far from thermal equilibrium but maintains a quasi-stationary distribution of entropy density.

Formally, a lamphronic state satisfies

$$D\nabla^2 S + \lambda|\nabla\Phi|^2 \approx \gamma S, \tag{34.1}$$

which is the steady-state condition for the entropy equation. This condition defines a manifold in the space of field configurations—the lamphronic manifold—that acts as a slow manifold for the full dynamics.

34.2 Field Relaxation Dynamics

The lamphrodyne relaxation process describes the evolution of the plenum from a lamphronic state (quasi-stationary entropy distribution) toward a state of lower free energy. During this process, the scalar field gradients $|\nabla\Phi|$ decrease, the vector flow $\mathbf{v}$ aligns with the gradient of Φ, and the entropy field S smooths out.

This relaxation has a natural interpretation in terms of the RG: it is the field-theoretic analog of the RG flow from a UV fixed point toward the IR fixed point. The lamphronic state corresponds to the UV regime where the entropy is large and heterogeneous; the relaxed state corresponds to the IR regime where the entropy is smooth and the effective description is simple.

Part XIII

Cosmological Implications

Chapter 35

Cosmic Structure without Metric Expansion

35.1 Standard Cosmological Expansion

The standard Λ-CDM cosmological model describes the universe as a spatially homogeneous and isotropic expanding spacetime, characterized by the Friedmann-Lemaître-Robertson-Walker (FLRW) metric

$$ds^2 = -dt^2 + a(t)^2 \left[\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right], \tag{35.1}$$

where $a(t)$ is the scale factor and k is the spatial curvature. The observed large-scale homogeneity and isotropy of the CMB strongly constrain the geometry to be very close to flat ($k \approx 0$) and the expansion to be very close to the Λ-CDM prediction.

35.2 Entropy Smoothing and Structure Formation

In the RSVP framework, cosmic structure formation can be interpreted as a process of entropy-driven smoothing and condensation. Initially, the plenum is in a high-entropy state with large spatial variations in Φ, $\mathbf{v}$, and S. As the entropy field undergoes lamphrodyne relaxation, regions of high scalar density Φ attract the vector flow $\mathbf{v}$, leading to the condensation of density into discrete structures.

This process is qualitatively similar to the Jeans instability in standard cosmology: gravitational attraction wins over pressure support for perturbations above a critical length scale, leading to the collapse of density fluctuations into gravitationally bound structures. In the RSVP framework the analogous role of gravity is played by the coupling between the scalar and vector fields: regions

of high Φ generate vector flows that advect more Φ toward them, leading to runaway condensation.

Chapter 36

Formation of Baryonic Structures

36.1 Solitonic Configurations

The most interesting structures that can form in the RSVP framework are *solitons*: stable, localized field configurations that maintain their shape through a balance between dispersive and nonlinear effects. In the scalar sector, a soliton is a solution of the nonlinear field equation

$$\Box\Phi + V'(\Phi) = 0 \tag{36.1}$$

that is localized in space and travels without changing shape.

The simplest example in one dimension is the kink solution of the ϕ^4 model: $\Phi(x) = v\tanh(x/\xi)$, where $\xi = (2\lambda)^{-1/2}v^{-1}$ is the width of the kink. This solution interpolates between the two degenerate vacua $\Phi = \pm v$ and carries a conserved topological charge $Q = \frac{1}{2v}\int_{-\infty}^{\infty} dx\, \partial_x\Phi$.

In the RSVP cosmological context, such solitonic structures provide natural candidates for stable localized concentrations of matter density—the analogs of particles in a field-theoretic description of matter.

Chapter 37

Long-Term Energy Redistribution

37.1 Trillion-Year Cosmology

The very long-term future of the universe is governed by the slow relaxation processes that become relevant only after stars, galaxies, and black holes have dissolved. In Λ-CDM the dominant process is the decay of black holes via Hawking radiation over timescales of $\sim 10^{67}$–10^{100} years, followed by the slow dissolution of any remaining structure by quantum tunneling and thermal fluctuations.

In the RSVP framework, the trillion-year cosmology is governed by the entropy field dynamics: the lamphrodyne relaxation continues on cosmological timescales, smoothing out the entropy gradients that give rise to the observed structure. The ultimate state of the plenum is a configuration of maximal entropy—a state where $S(x)$ is spatially uniform and $|\nabla\Phi| \to 0$—which corresponds to the heat death of the universe.

37.2 Global Equilibrium States

The global equilibrium state of the RSVP system is reached when the entropy field satisfies $\partial_t S = 0$ everywhere, which requires $D\nabla^2 S + \lambda|\nabla\Phi|^2 = \gamma S$ globally. This is not in general a uniform state: it is the fixed-point configuration of the entropy dynamics, determined by the balance between diffusion, production, and dissipation.

The existence and properties of this global equilibrium state depend on the boundary conditions of the cosmological system and on the values of the coupling constants D, λ, and γ. In the cosmologically relevant regime where the universe is effectively an isolated system (no external boundary), the global equilibrium corresponds to the heat death: $\Phi = \text{const}$, $\mathbf{v} = 0$, $S = \lambda|\nabla\Phi|^2/\gamma = 0$.

Part XIV

Mathematical Formalism

Chapter 38

Variational Formulations

38.1 Action Principles for Coupled Fields

The full RSVP action that generates the coupled dynamics of Φ, v^μ, and S must be constructed to satisfy several requirements: Lorentz invariance, diffeomorphism invariance (for coupling to curved spacetime), the entropy-increase condition (second law), and compatibility with the known field equations of general relativity in the appropriate limit.

A general form consistent with these requirements is

$$S_{\text{RSVP}} = \int d^4x\sqrt{-g}\left[\frac{M_{\text{Pl}}^2}{2}R - \frac{1}{2}(\partial\Phi)^2 - V(\Phi) - \frac{1}{4}F^{(v)}_{\mu\nu}F^{(v)\mu\nu} - \frac{1}{2}(\partial S)^2 - \alpha(\nabla\Phi)^2 S\right], \tag{38.1}$$

where $F^{(v)}_{\mu\nu} = \partial_\mu v_\nu - \partial_\nu v_\mu$ is the field strength of the vector field, and α is the entropy-scalar coupling. The last term encodes the entropy production from scalar gradients.

38.2 Euler-Lagrange Equations

Varying the RSVP action with respect to each field yields the coupled equations of motion:

for the scalar field: $\Box\Phi - V'(\Phi) - 2\alpha(\nabla^2\Phi)S - 2\alpha(\partial_\mu\Phi)(\partial^\mu S) = 0$;

for the vector field: $\partial_\nu F^{(v)\mu\nu} = 0$ (a Maxwell-type equation);

for the entropy field: $\Box S + \alpha(\partial\Phi)^2 = 0$;

for the metric: the Einstein equations with energy-momentum tensor $T_{\mu\nu} = T^{(\Phi)}_{\mu\nu} + T^{(v)}_{\mu\nu} + T^{(S)}_{\mu\nu}$.

These equations reduce to the standard Klein-Gordon and Einstein equations in the limit $\alpha \to 0$ and $v_\mu \to 0$, ensuring that the RSVP framework is compatible with known physics in the appropriate limit.

Chapter 39

Topological Structures

39.1 Homology and Cohomology in Field Theory

Topological methods classify field configurations that cannot be continuously deformed into one another. The key invariants are winding numbers, linking numbers, and more general cohomological classes.

For the vector field v_μ, the de Rham cohomology classifies the topological structure of the flow. A one-form $v = v_\mu dx^\mu$ is *closed* if $dv = 0$ (zero curl) and *exact* if $v = d\chi$ for some function χ. The first cohomology group $H^1(M) = \ker(d)/\mathrm{im}(d)$ measures the topological obstruction to exactness. In a simply connected spacetime $H^1 = 0$ and every closed form is exact, but on a spacetime with non-trivial topology (e.g., a torus) $H^1 \neq 0$ and topologically non-trivial vector flow configurations exist.

39.2 Topological Defects and Cosmic Strings

Cosmic strings arise as topological defects when a $U(1)$ symmetry is spontaneously broken. The vacuum manifold of the Abelian Higgs model is $\mathcal{V} = S^1$ (a circle). Field configurations $\phi(x)$ that wind around this circle as x traverses a loop in physical space cannot be continuously deformed to the trivial configuration. These winding configurations are classified by the fundamental group $\pi_1(S^1) = \mathbb{Z}$, and each integer $n \in \mathbb{Z}$ labels a topological sector.

In the RSVP framework, analogous topological structures arise from the winding of the phase of the scalar field Φ (if Φ is complex) or from the vorticity of the vector field v_μ. The topological charge of such a defect is conserved by continuity and cannot be removed without cost; defects can only annihilate with anti-defects of the opposite topological charge.

Chapter 40

Categorical Descriptions of Dynamics

40.1 Categories of Field Configurations

The space of all field configurations of the RSVP system at a given time forms a category: the objects are field configurations (Φ, v^μ, S), and the morphisms are gauge transformations, diffeomorphisms, and other symmetry transformations. The dynamics of the system defines a functor from the category of initial configurations to the category of final configurations, mapping each initial state to its time-evolved counterpart.

This categorical perspective becomes useful when studying symmetry-breaking: a symmetry transformation that is a morphism in the category of field configurations may become trivial (the identity morphism) in the category of ground states if the symmetry is broken. The broken symmetry is encoded in the difference between the categories of field configurations (before symmetry breaking) and ground states (after symmetry breaking).

40.2 Compositional Dynamics and Modularity

One of the strengths of the categorical formulation is its support for modularity: the dynamics of the full RSVP system can be described as a composition of functors describing the dynamics of subsystems. The scalar sector, vector sector, and entropy sector each have their own dynamics (their own functors), and the full coupled dynamics is their composition through the interaction terms.

This compositional structure is related to the *Spherepop* formalism of process-oriented computation, where complex operations are built from simple atomic operations (pop, bind, collapse) arranged in sequences. In the field-theory con-

text, the analog of Spherepop composition is the sequential application of the coarse-graining functor: each RG step is a "pop" that removes one shell of high-momentum modes, leaving a simpler effective theory.

Part XV

Conceptual Consequences

Chapter 41

Emergent Geometry from Field Interactions

41.1 Geometry from Correlations

One of the most profound questions raised by the RSVP framework is whether spacetime geometry is fundamental or emergent. In general relativity, the metric is a fundamental field; in the RSVP framework, it might be derived from the correlations between the scalar and vector fields.

The basic idea is that distances between points in spacetime are encoded in the two-point correlation function of the scalar field:

$$G(x, y) = \langle \Phi(x)\Phi(y) \rangle. \tag{41.1}$$

If $G(x, y)$ falls off exponentially with a correlation length ξ, then ξ provides a natural notion of "distance" between x and y. A curved spacetime geometry corresponds to a spatially varying correlation length, $\xi = \xi(x)$, which in turn corresponds to a spatially varying effective mass of the scalar field.

41.2 Relation to Asymptotic Safety

The connection between emergent geometry and asymptotic safety is through the RG flow: the metric that emerges from the scalar field correlations at a given scale k is the "effective metric" of the theory at that scale. As k flows from the UV to the IR, the effective metric changes, and the geometry of spacetime "emerges" as the coarse-grained description that captures the long-range correlations of the scalar field.

At the NGFP of asymptotically safe gravity, the effective metric becomes

scale-invariant: it looks the same at all scales. This scale invariance of the metric corresponds to scale invariance of the scalar field correlations, which in turn corresponds to the critical point of the scalar field theory. The asymptotically safe fixed point in the gravitational sector is therefore related to the critical fixed point in the scalar sector through this emergent geometry picture.

Chapter 42

Observers, Measurement, and Coarse-Graining

42.1 Observer-Dependent Descriptions

A central theme of both quantum mechanics and general relativity is the observer-dependence of physical descriptions. In quantum mechanics, the measurement process involves an observer choosing which observable to measure, and different choices lead to different descriptions of the same physical state. In general relativity, different observers moving through curved spacetime assign different coordinates and energy values to the same events.

In the RSVP framework, observer-dependence enters through the coarse-graining procedure. Different observers at different spatial locations or moving at different velocities will perform coarse-graining differently, obtaining different effective descriptions of the plenum. The entropy field $S(x)$ encodes this observer-dependence: the entropy assigned to a given region depends on what information is accessible to the observer making the assignment.

42.2 Local Information Extraction

The measurement of a physical quantity in the RSVP framework corresponds to a local interaction between the measuring apparatus and the plenum. The apparatus couples to the scalar field Φ at its location and records the value of some observable, which might be the local energy density, the local entropy, or the local vector flow.

This interaction disturbs the plenum locally, producing a small increase in the entropy field S in the vicinity of the measurement (due to the entropy production term $\lambda|\nabla\Phi|^2$, which is non-zero when the measurement apparatus cre-

ates a gradient in Φ). The measurement is therefore not passive: it produces an irreversible change in the state of the plenum, consistent with the general principle that measurement is an irreversible, entropy-increasing process.

Part XVI

Open Problems and Future Directions

Chapter 43

Experimental Tests

43.1 Astrophysical Signatures of the RSVP Framework

The RSVP framework makes several qualitative predictions that could in principle be tested observationally, though the quantitative predictions require more detailed modeling than is currently available.

The entropy smoothing mechanism for large-scale structure formation predicts a specific relationship between the entropy field profile and the matter density profile. In regions of high entropy production (high $|\nabla\Phi|$), the matter density should be lower than in regions of low entropy production (smooth Φ). This is qualitatively consistent with the observed void-filament structure of the cosmic web, where matter is concentrated in filaments and sheets with large voids between them.

The solitonic structures that form in the RSVP framework could in principle be identified with dark matter halos or with compact objects. The characteristic scale of these solitons is set by the ratio of the dispersive and nonlinear terms in the field equation, which depends on the values of m^2 and λ in the scalar potential.

43.2 Gravitational Wave Backgrounds

The RSVP framework predicts a stochastic gravitational wave background from the dynamics of the vector field, through the coupling between v_μ and the metric. The spectrum of this background depends on the dynamics of the vector field during the early cosmological evolution of the plenum.

If the vector field undergoes a symmetry-breaking transition at a scale v_{RSVP}, the resulting gravitational wave background has a spectrum similar to that of

cosmic strings, with amplitude and spectral index determined by v_{RSVP} and the loop dynamics of the vector field defects. The current NANOGrav constraints on the stochastic background therefore bound v_{RSVP} in a way that can be used to constrain the RSVP parameters.

Chapter 44

Connections to Other Approaches

44.1 Loop Quantum Gravity

Loop quantum gravity (LQG) quantizes the geometry of spacetime directly, representing the quantum states of the gravitational field by spin networks and spin foams. The fundamental variables are the holonomies of the Ashtekar connection and the fluxes of the densitized triad field.

The connection between LQG and the RSVP framework is not immediately apparent, but there are structural similarities. The spin network description of quantum geometry can be interpreted as a discrete version of the emergent geometry scenario: the nodes and links of the spin network carry discrete area and volume quanta, and the continuum geometry emerges in the large-spin limit. The RSVP scalar field might be related to the large-spin limit of the LQG area spectrum, with $\Phi(x)$ encoding the coarse-grained area density at each point.

44.2 Causal Set Theory

Causal set theory posits that spacetime is fundamentally discrete, with the elementary events forming a partially ordered set (poset) related by causal precedence. The continuum geometry of general relativity emerges as an approximation when the causal set is viewed at scales much larger than the Planck length.

The RSVP framework shares with causal set theory the idea that the continuum geometry is not fundamental. However, RSVP takes a field-theoretic rather than combinatorial approach: instead of discrete events and their causal relations, one has continuous fields and their correlations. The connection between the two approaches might be through the replica trick or through a suitable continuum limit of the causal set.

44.3 Tensor Models and Universality Classes

Tensor models are random geometry models in which the fundamental variables are tensors T_{abc} (rather than matrices M_{ab} as in matrix models) and the partition function is an integral over all tensor configurations. They are relevant to quantum gravity because they provide a combinatorial quantization of simplicial manifolds in dimension $d \geq 3$.

Recent work by Eichhorn, Gurau, Gyftopoulos, and collaborators has shown that asymptotic safety can be extended to tensor field theories, with the possible existence of non-Gaussian fixed points in the coupled tensor-gravity system. This suggests a surprising universality: the same RG fixed-point structure that controls the UV behavior of metric-based quantum gravity may also control the critical behavior of tensor models, unifying very different approaches to quantum gravity under the umbrella of renormalization group universality.

Chapter 45

Future Directions

45.1 Non-Polynomial Dynamics

The most important open problem in connecting the RSVP framework to asymptotic safety is the treatment of non-polynomial dynamics. Current FRG calculations for Horndeski gravity and the Abelian Higgs model use polynomial truncations of the scalar potential, which capture only the local behavior near the fixed-point configuration. The true dynamics of the RSVP scalar field may require the full non-polynomial potential $V(\Phi)$, which can only be captured by the local potential approximation (LPA) or beyond.

Extending the LPA to the RSVP system, where the scalar field is coupled to the vector and entropy fields, is a non-trivial but tractable problem. The key challenge is that the LPA introduces a functional PDE for $V(\Phi; k)$ that must be solved numerically, and the coupling to the vector and entropy fields introduces additional functional dependencies.

45.2 Lorentzian Signature and Real-Time Dynamics

All existing FRG calculations for asymptotically safe gravity are performed in Euclidean signature. The extension to Lorentzian signature, which is required for genuine cosmological applications, remains an open problem. Several approaches have been proposed: analytic continuation of the Euclidean results, causal perturbation theory, or a direct Lorentzian path integral.

For the RSVP framework, the Lorentzian extension is particularly important because the entropy field has a preferred arrow of time: entropy production is irreversible and defines a temporal direction. In Euclidean signature this asymmetry is lost. The correct formulation of the RSVP entropy dynamics in Lorentzian signature requires a careful treatment of the irreversibility, which

might involve a Lindblad-type open quantum system approach or a Schwinger-Keldysh formalism for the entropy field.

45.3 Unified Theoretical Frameworks

The deepest open question is whether asymptotic safety, the RSVP framework, and the various other approaches to quantum gravity can be unified into a single coherent framework. The renormalization group provides a natural organizing principle: different approaches might correspond to different truncations of the same underlying theory space, with the NGFP as the common organizing center.

The information-geometric interpretation developed in Part IX suggests a direction: the RSVP framework, asymptotic safety, and causal set theory might all be descriptions of the same information-geometric structure on theory space, related by different choices of coarse-graining procedure. The RSVP fields (Φ, v^μ, S) might be the natural coordinates on the information manifold of physical theories, with the NGFP corresponding to the point of maximum informational self-consistency.

Whether this unification program can be made precise, and whether it makes falsifiable predictions, is one of the central open problems of contemporary theoretical physics.

Appendix A

Mathematical Background

A.1 Linear Algebra Review

A *vector space* V over $\mathbb{R}$ is a set equipped with addition and scalar multiplication satisfying the standard axioms. A *basis* is a maximal linearly independent set; its cardinality is the *dimension* of V. A *linear map* $T : V \to W$ satisfies $T(\alpha v + \beta w) = \alpha T v + \beta T w$.

An *inner product* $\langle \cdot, \cdot \rangle : V \times V \to \mathbb{R}$ is symmetric, bilinear, and positive-definite. A *matrix* A represents a linear map in a chosen basis; its *eigenvalues* λ_i satisfy $A v_i = \lambda_i v_i$ for the eigenvectors v_i.

A.2 Differential Geometry Basics

A *smooth manifold* M of dimension n is a topological space locally homeomorphic to $\mathbb{R}^n$ with smooth transition functions between charts. A *tangent vector* at $p \in M$ is a derivation on smooth functions at p. The *metric tensor* $g_{\mu\nu}$ is a symmetric positive-definite bilinear form on each tangent space; it defines distances and angles on M.

The *covariant derivative* ∇_μ generalizes partial differentiation to tensors on curved manifolds, via the Christoffel symbols $\Gamma^\rho_{\mu\nu} = \frac{1}{2} g^{\rho\sigma}(\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\mu\sigma} - \partial_\sigma g_{\mu\nu})$. The *Riemann curvature tensor* measures the non-commutativity of covariant derivatives: $[\nabla_\mu, \nabla_\nu] V^\rho = R^\rho{}_{\sigma\mu\nu} V^\sigma$.

A.3 Functional Analysis

A *functional* is a map from a function space to $\mathbb{R}$. The *functional derivative* $\delta F/\delta\phi(x)$ is defined by

$$F[\phi + \epsilon\eta] = F[\phi] + \epsilon \int dx \, \frac{\delta F}{\delta\phi(x)} \eta(x) + \mathcal{O}(\epsilon^2). \tag{A.1}$$

The *path integral* $\int \mathcal{D}\phi \, e^{-S[\phi]}$ is a functional integral over all field configurations, defined rigorously through lattice discretization and a continuum limit, or in perturbation theory through a Gaussian reference measure and Feynman diagram expansion.

Appendix B

Notation and Conventions

Throughout this text we use natural units $\hbar = c = k_B = 1$. The metric signature is $(-+++)$ in Lorentzian signature and $(++++)$ in Euclidean signature. Greek indices $\mu, \nu, \ldots$ run over $0, 1, 2, 3$ in Lorentzian and $1, 2, 3, 4$ in Euclidean. Latin indices $i, j, k, \ldots$ are used for spatial indices and for coupling constant labels.

The RG scale k has dimensions of mass (energy), and $t = \ln(k/k_0)$ where k_0 is a reference scale. Dimensionless couplings are denoted $g_i = k^{-d_i}\bar{g}_i$ where $d_i = 4 - [\mathcal{O}_i]$ is the canonical dimension of the coupling. The Newton coupling is $G = G_N$ (dimensionful) or $g = G_N k^2$ (dimensionless).

Appendix C

Solutions to Selected Exercises

Chapter 1, Exercise 1. The Bohr radius $a_0 = \hbar^2/(m_e e^2)$ has dimensions $[a_0] = [\hbar^2/(m_e e^2)] = \text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2}/(\text{kg} \cdot (\text{C} \cdot \text{V})) = \text{m}$. The ground state energy is $E_1 \sim \hbar^2/(m_e a_0^2) = m_e e^4/\hbar^2 \approx -13.6\,\text{eV}$.

Chapter 2, Exercise 1. The fixed points satisfy $-x + y^2 = 0$ and $-2y = 0$, giving $y = 0$ and $x = 0$. The only fixed point is $(0, 0)$. The Jacobian is $J = [[-1, 2y_*],\ [0, -2]]|_{(0,0)} = [[-1, 0], [0, -2]]$, with eigenvalues $\lambda_1 = -1$, $\lambda_2 = -2$. Both are negative: the fixed point is a stable node.

Chapter 8, Exercise 1. Setting $\beta = 0$: $-g + g^2 = 0$ gives $g = 0$ (Gaussian FP, stable since $\beta'(0) = -1 < 0$) and $g = 1$ (NGFP, unstable since $\beta'(1) = 1 > 0$). A trajectory starting at $g_0 = 0.5$ is attracted toward $g = 0$ since $\beta(0.5) = -0.5 + 0.25 = -0.25 < 0$.