

Point Estimation of the Central Orientation of Random Rotations

Bryan Stanfill*

Ulrike Genschel

and

Heike Hofmann

Department of Statistics, Iowa State University

October 12, 2012

Abstract

Data in the form of random three-dimensional rotations have application in computer science, kinematics and materials sciences, among other areas. Estimating the central orientation from a sample of such data is an important problem, which is complicated by the fact that several different approaches exist for this, motivated by various geometrical and decision-theoretic considerations. However, little is known about how such estimators compare, especially on common distributions for location models with random rotations. We examine four location estimators, three of which are commonly found in different literatures and the fourth estimator (a projected median) is newly introduced. Our study unifies existing literature and provides a detailed numerical investigation of location estimators for three commonly used rotation distributions in statistics and materials science. While the data-generating model influences the best choice of an estimator, the proposed projected median emerges as an overall good performer, which can be suggested without particular distributional assumptions. We illustrate the estimators and our findings with data from a materials science study by approximating the central orientation of cubic crystals on the micro-surface of a metal.

Keywords: Cayley distribution, Electronic Backscatter Diffraction, geodesic distance, Matrix Fisher distribution, Projected median, Rotation Group

*The authors gratefully acknowledge the Technometrics reviewers and associate editor for providing us with valuable comments to strengthen this manuscript.

1 Introduction

Data in the form of 3×3 rotation matrices find application in several scientific areas, such as biomedical engineering, computer visioning, and geological and materials sciences, where such data represent the positions of objects within some three-dimensional reference frame. For example, ? examine rotation matrix data in studying body positions whilst operating machinery. ? consider this type of orientation data in magnetic resonance imaging and in shape analysis; similar examples can be found in ?, ?, ?, or ?. The data in our illustrative example to follow arise from a study in materials science, where 3×3 rotations represent the orientations of cubic crystals on the micro-surface of a metal specimen as measured through electron backscatter diffraction (EBSD) and “grains” within metals are composed of crystals which roughly share a common orientation; see ? for details on EBSD data.

From a sample of orientations, an important interest is often the estimation of a main or central orientation \mathbf{S} . That is, letting the rotation group $SO(3)$ denote the collection of all 3×3 rotation matrices, observations $\mathbf{R}_1, \dots, \mathbf{R}_n \in SO(3)$ can be conceptualized as a random sample from a *location model*

$$\mathbf{R}_i = \mathbf{S}\mathbf{E}_i, \quad i = 1, \dots, n, \quad (1)$$

where $\mathbf{S} \in SO(3)$ is the *fixed* parameter of interest indicating an orientation of central tendency, and $\mathbf{E}_1, \dots, \mathbf{E}_n \in SO(3)$ denote i.i.d. *random* rotations which symmetrically perturb \mathbf{S} . The data-generating model in (??) is a rotation-matrix analog of a location model for scalar data $Y_i = \mu + e_i$, where $\mu \in \mathbb{R}$ denotes a mean and $e_i \in \mathbb{R}$ denotes an additive error symmetrically distributed around zero. This representation (??) for orientations is quite common and, in fact, a variety of parametric models exist for describing symmetrically distributed rotations \mathbf{E}_i , such as the symmetric matrix Fisher distribution (?), the symmetric Cayley distribution (?) and the circular-von Mises-based rotation distribution (?) in the statistics literature, as well as the Bunge distribution (?), the isotropic Gaussian distribution (??) and the de la Vallée Poussin distribution (?) in the materials science literature. Our goal in this paper is to summarize and compare the most frequently proposed approaches for point estimation of \mathbf{S} based on a sample of orientation data generated by

(??). Depending on the scientific literature, the approaches can be quite different.

The topic of location estimation has received considerable attention for directional data on circles or spheres, (see ?????????), but less is known about estimator properties with rotation data. As a compounding factor, several current approaches to estimating \mathbf{S} have arisen out of literatures having differing statistical and geometrical emphases. In the applied sciences literature, estimators of \mathbf{S} are typically based on distance measures involving *non-Euclidean* (i.e., Riemannian) geometry, such as the *geometric mean* (????) or, more recently, the *geometric median* (??). Preferences may depend on outliers in the data, but such suggestions for estimating \mathbf{S} often do *not* consider the potential impact of the underlying data-generating mechanism. On the other hand, approaches in the statistics literature tend to motivate an estimator for \mathbf{S} through likelihood or moment-estimation principles applied to a specifically assumed distributional model (e.g., matrix Fisher or Cayley distribution) for the symmetric rotation errors \mathbf{E}_i (????). Almost always, this estimator turns out to be a *projected arithmetic mean* based on *Euclidean* geometry. Hence, in addition to potential distributional assumptions, more fundamental divisions in estimation approaches may be attributable to different geometrical perspectives with rotation data.

Considering the potential effects of an underlying data generation model as well as the choice of geometry (i.e., Euclidean vs. Riemannian), the above discussion indicates a need to investigate and identify good point estimators for rotation data. In particular, because estimators in the applied sciences literatures are often selected without decision-theoretical considerations based on underlying distributions, it is of interest to understand how different location estimators behave across common distributions for rotations. In this paper, we evaluate four common estimators for \mathbf{S} in the context of the location model (??). These are either mean- or median-based, and either Euclidean- or Riemannian-based in geometry; the Euclidean-based median estimator is new here but its inclusion is natural and its performance can be generally quite good and (as will be demonstrated) broadly recommendable. Through simulation, we compare how these estimators perform with respect to three common probability models for symmetric rotation errors as defined in (??), namely the circular-von Mises-based distribution, the symmetric matrix Fisher distribution and the symmetric Cayley distribution. The matrix Fisher is arguably the

most common distribution in the statistics literature (see ?). While not noted previously, the symmetric Cayley and the de la Vallée Poussin distribution are in fact the same; the de la Vallée Poussin distribution has been advocated in the material science literature (?). The circular-von Mises-based distribution is included because the distribution is non-regular and has been applied to EBSD data (?). We describe how error distribution assumptions for rotation data, in particular their variability and tail behavior, translate into performance differences among point estimators based on varying geometric considerations.

The remainder of the manuscript is organized as follows. Section ?? provides a brief background on the geometry of rotations and different distance metrics. Section ?? then describes the location estimators for rotation data and compares their geometric underpinnings, which serves to unify some of the existing estimation literature. Section 4 explains the design of the simulation study followed by a summary of our main findings in Section 5. Section 6 provides an illustration of the estimation methods for EBSD data in a materials science application. We provide concluding remarks and future research possibilities in Section 7.

2 Background

2.1 Geometry of Three-dimensional Orientations

Three-dimensional orientation data consist of observations belonging to the group $SO(3)$ of all 3×3 rotation matrices, where an element \mathbf{R} in $SO(3)$ is an orthogonal 3×3 matrix (i.e., $\mathbf{R}^\top \mathbf{R} = \mathbf{I}_{3 \times 3}$) with determinant one. As $SO(3)$ is a Lie group, its elements live on a differentiable manifold. This aspect is helpful in understanding the two different geometric approaches for estimating the central location $\mathbf{S} \in SO(3)$ from a sample of orientation data, referred to here as the *intrinsic* and the *embedding* estimation approaches (see also ? and ? for analogs with directional data).

The rotation group $SO(3)$ is not closed under routine addition or scalar multiplication (i.e., operations natural to statisticians). Hence, statistical estimation approaches often *embed* the rotation group into the higher-dimensional linear space consisting of all 3×3 real matrices, denoted as $\mathcal{M}(3)$. Doing so enables the use of the familiar Euclidean

geometry (and “averaging” notions) to define standard distance measures and loss criteria for obtaining location estimators (see Section ?? and the estimators given in Sections ?? and ??). This embedding technique has been largely applied by statisticians, typically resulting in the projected arithmetic mean of Section ?. See, for example, ?? and ??; the Bayesian estimator used in ? is also a concrete example of this approach.

Alternatively, *intrinsic* estimation approaches use Riemannian geometry to define distances that account for the innate topology or curvature of the space $SO(3)$. In the intrinsic approach, each rotation from $SO(3)$ is associated with a skew-symmetric matrix $\Phi(\mathbf{W})$, defined as

$$\Phi(\mathbf{W}) = \begin{bmatrix} 0 & -w_3 & -w_2 \\ w_3 & 0 & -w_1 \\ w_2 & w_1 & 0 \end{bmatrix}$$

for $\mathbf{W} = (w_1, w_2, w_3)^\top \in \mathbb{R}^3$. That is, through a so-called exponential operator, we map $\Phi(\mathbf{W})$ to a rotation matrix as

$$\exp[\Phi(\mathbf{W})] = \mathbf{I}_{3 \times 3} \cos(r) + \sin(r)\Phi(\mathbf{U}) + (1 - \cos r)\mathbf{U}\mathbf{U}^\top$$

where $r = \|\mathbf{W}\|$ and $\mathbf{U} = \mathbf{W}/\|\mathbf{W}\|$. The space $\mathfrak{so}(3)$ of all skew-symmetric matrices forms the tangent space (Lie-algebra) of $SO(3)$, which is closed under familiar summation and scalar multiplication operations in the usual (i.e., element-wise) manner. The fact that $SO(3)$ is a differentiable manifold allows a distance measure (i.e., the geodesic distance in Section ??) to be defined between points in $SO(3)$ according to Riemannian geometry. The resulting geodesic distance underlies the “geometric” location estimator for \mathbf{S} commonly found in computer science (???) and engineering applications (?); see Sections ?? and ??.

Before leaving this section, it is helpful to note that each rotation matrix \mathbf{R} can be uniquely associated with a pair (r, \mathbf{U}) , where $r \in (-\pi, \pi]$ and $\mathbf{U} \in \mathbb{R}^3$, $\|\mathbf{U}\| = 1$, through

$$\mathbf{R} = \mathbf{R}(r, \mathbf{U}) = \exp[\Phi(\mathbf{U}r)] \in SO(3). \quad (2)$$

This is the so-called axis-angle representation of \mathbf{R} , where \mathbf{R} is represented by rotating the coordinate axis $\mathbf{I}_{3 \times 3}$ about the axis $\mathbf{U} \in \mathbb{R}^3$ by the angle r . In the materials science

literature, \mathbf{U} and r are commonly referred to as the misorientation axis and misorientation angle of \mathbf{R} with respect to $\mathbf{I}_{3 \times 3}$; see ?.

3 Location Estimators

This section begins with a description of the distance metrics the result as a consequence of which geometry is chosen. It then describes four estimators for the location parameter $\mathbf{S} \in SO(3)$ from orientation data generated by (??) based on the two geometric choices (i.e., embedding (??) in Sections ?? and ?? or intrinsic (??) in Sections ?? and ??) and two decision-theoretic loss functions (i.e., squared distances in Sections ?? and ?? or pure distances in Sections ?? and ??). The extent to which the choice of geometry or loss function matters in the estimation of \mathbf{S} will be an important aspect explored in Section ??. A summary of the four estimators and their properties is given in Table ??.

3.1 Choice of Distance Metrics

The choice of geometry, i.e. Riemannian (non-Euclidean) or Euclidean, results in two different metrics to measure the distance between two rotation matrices \mathbf{R}_1 and $\mathbf{R}_2 \in SO(3)$. Under the embedding approach, the natural distance metric between two random matrices is the Euclidean distance, d_E , which is induced by the Frobenius norm

$$d_E(\mathbf{R}_1, \mathbf{R}_2) = \|\mathbf{R}_1 - \mathbf{R}_2\|_F, \quad (3)$$

where $\|\mathbf{A}\|_F = \sqrt{\text{tr}(\mathbf{A}^\top \mathbf{A})}$ denotes the Frobenius norm of a matrix \mathbf{A} and $\text{tr}(\cdot)$ denotes the trace of a matrix. The Euclidean distance between two rotation matrices corresponds to the shortest cord in $\mathcal{M}(3)$ that connects them. If $r \in [0, \pi]$ denotes the misorientation angle in the angle-axis representation (??) of $\mathbf{R}_1^\top \mathbf{R}_2 \equiv \mathbf{R}_1^\top \mathbf{R}_2(r, \mathbf{U})$ (so that $\text{tr}(\mathbf{R}_1^\top \mathbf{R}_2) = 1 + 2 \cos r$), then $d_E(\mathbf{R}_1, \mathbf{R}_2) = 2\sqrt{(1 - \cos r)}$ holds.

By staying in the Riemannian space $SO(3)$ under the intrinsic approach, the natural distance metric becomes the Riemannian (or geodesic) distance, d_R , by which the distance

between two rotations $\mathbf{R}_1, \mathbf{R}_2 \in SO(3)$ is defined as

$$d_R(\mathbf{R}_1, \mathbf{R}_2) = \frac{1}{\sqrt{2}} \|\text{Log}(\mathbf{R}_1^\top \mathbf{R}_2)\|_F = r, \quad (4)$$

where $\text{Log}(\mathbf{R})$ denotes the principle logarithm of \mathbf{R} (i.e., $\text{Log}(\mathbf{R}) = \text{Log}(\mathbf{R}(\mathbf{U}, r)) = \Phi(r\mathbf{U})$ in (??)) and $r \in [0, \pi]$ is the misorientation angle of $\mathbf{R}_1^\top \mathbf{R}_2$. The Riemannian distance corresponds to the length of the shortest path that connects \mathbf{R}_1 and \mathbf{R}_2 *within* the space $SO(3)$. For this reason, the Riemannian distance is often considered the more natural metric on $SO(3)$; see ? for this discussion along with more details on exponential/logarithmic operators related to $SO(3)$.

To make the difference between the Euclidean and Riemannian metrics more concrete, consider the lower dimension example given in Figure ??. Here we plot two points, $\mathbf{R}_1, \mathbf{R}_2$, to visualize the endpoints of two rotations on $SO(2)$. The Riemannian distance between them, $d_R(\mathbf{R}_1, \mathbf{R}_2)$, is indicated by the thick black curved line. As $\mathbf{R}_1 \mathbf{v}$ rotates to $\mathbf{R}_2 \mathbf{v}$ it does so along this curve therefore this distance is considered to stay in the space. The Euclidean distance, $d_E(\mathbf{R}_1, \mathbf{R}_2)$, is illustrated by the straight gray line. It is impossible for $\mathbf{R}_1 \mathbf{v}$ to move along this line, meaning we must leave $SO(2)$ in order to compute this distance.




Figure 1: An illustration of the difference between the Euclidean and Riemannian distance metrics on $SO(2)$.

3.2 The projected arithmetic mean

We begin with the definition of the arithmetic mean for orientation data, as its analog is most frequently encountered in the statistical literature for directional data (e.g. see ?). For a sample of n random rotations $\mathbf{R}_i \in SO(3)$, $i = 1, 2, \dots, n$, this mean-type estimator is defined as

$$\hat{\mathbf{S}}_E = \arg \min_{\mathbf{S} \in SO(3)} \sum_{i=1}^n d_E^2(\mathbf{R}_i, \mathbf{S}) = \arg \max_{\mathbf{S} \in SO(3)} \text{tr}(\mathbf{S}^\top \bar{\mathbf{R}}) \quad (5)$$

where $\bar{\mathbf{R}} = \frac{1}{n} \sum_{i=1}^n \mathbf{R}_i$. The estimator is obtained by minimizing the sum of the squared distances in the Euclidean sense in the ambient space $\mathcal{M}(3)$, which then is projected back into $SO(3)$. ?, who studied the mathematical characteristics of this estimator in detail, therefore refers to it as the *projected arithmetic mean*. This estimator's appeal lies in its simplicity and statistically intuitive nature, though it has been noted that the estimator is not invariant under rigid transformations (see ?). However, this estimator does correspond to the maximum likelihood estimator of \mathbf{S} when the symmetrically distributed rotation errors in (??) follow a matrix Fisher distribution (?). ? also derived this estimator as the method of moment estimator under a Cayley distribution, and ? showed that the projected arithmetic mean corresponds to the maximum quasi-likelihood estimator for orientation data with rotation errors arising from the circular-von Mises-based distribution.

? and ? independently offered algorithms to find this matrix. ? refined their solutions and also considered special cases such as $\det(\bar{\mathbf{R}}) = 0$.

3.3 The projected median

A modification of the estimator from Section ?? is obtained by replacing the squared distances in (??) with pure distances, leading to a median-type estimator defined as

$$\tilde{\mathbf{S}}_E = \arg \min_{\mathbf{S} \in SO(3)} \sum_{i=1}^n d_E(\mathbf{R}_i, \mathbf{S}). \quad (6)$$

For rotation data, we refer to this estimator of \mathbf{S} as the *projected median*. The analog of this estimator for spherical data has been considered by ? (i.e., the so-called normalized spatial median of ?) for estimating the central direction of data points on the sphere. Those

authors found the estimator to be preferable for spherical data following the Fisher model on the sphere. For additional information on (??) and related estimators for directional data (e.g. the mediancentre (?) or the Weber point (?)), see ?.

An algorithm to compute this estimator hasn't been proposed as of this writing, so we detail our method here. Our method is based on the Weiszfeld algorithm originally given by ?. The algorithm requires an initial value that does not equal any sample point. For the purpose of speeding up computing time we use $\hat{\mathbf{S}}_E$ as the starting point. Note that the solution is generally not sensitive to the choice of starting points unless the data exhibit extreme spread.

1. Set $\hat{\mathbf{S}} = \hat{\mathbf{S}}_E$ and choose an arbitrarily small stopping rule ε .

2. For $i = 1, \dots, n$ compute $\mathbf{s}_i = \mathbf{R}_i - \hat{\mathbf{S}}$.

3. Calculate

$$\bar{\mathbf{R}}_W = \frac{\sum_{i=1}^n \mathbf{R}_i / \|\mathbf{s}_i\|_F}{\sum_{i=1}^n 1 / \|\mathbf{s}_i\|_F}$$

which we call the weighted mean with respect to $\hat{\mathbf{S}}$.

4. Define $\hat{\mathbf{S}}_{\text{new}}$ to be the $\mathcal{M}(3)$ projection of $\bar{\mathbf{R}}_W$.

5. If $\varepsilon > \|\hat{\mathbf{S}} - \hat{\mathbf{S}}_{\text{new}}\|_F$ return $\tilde{\mathbf{S}}_E = \hat{\mathbf{S}}_{\text{new}}$; otherwise set $\hat{\mathbf{S}} = \hat{\mathbf{S}}_{\text{new}}$ and return to step 2.

3.4 The geometric mean

As sketched in Section ??, the Lie group property of $SO(3)$ provides us with a convenient transform from $SO(3)$ into the tangent space $\mathfrak{so}(3)$ that is closed under addition and scalar multiplication. Obtaining the median or mean in this transformed space and projecting the result back to $SO(3)$ corresponds to the rotation that minimizes the first and second order Riemannian distances, respectively (????). ? made use of Riemannian manifolds to compute what is often called the Riemannian center of mass. ? applied Karcher's ideas to rotation matrices and defined

$$\hat{\mathbf{S}}_R = \arg \min_{\mathbf{S} \in SO(3)} \sum_{i=1}^n d_R^2(\mathbf{R}_i, \mathbf{S}). \quad (7)$$

which was termed as the *geometric mean*. Note that the solution to (??) may not be unique. Uniqueness is tied to the property of geodesic convexity of the objective function in (??). For more information, we refer to ?. Additionally, (??) generally does not have a closed-form solution making this estimator much more computationally intensive than its Euclidean counterpart (the projected arithmetic mean of Section ??). The algorithm proposed by ? was used in our simulation study.

3.5 The geometric median

The median-type counterpart to the geometric mean was defined first in the context of spherical data by ? as the point on the sphere that minimizes the sum of the arc lengths to all observations in the sample. For this type of data, the resulting estimator is known as the spherical median, which is a special case of the generalized median in \mathbb{R}^d proposed by ?. For spherical data, an alternative formulation to the spherical median has been given by ? in the framework of data depth leading, however, to the same solution.

We give an adaptation of the spherical median to rotation matrices. Recall that the shortest geodesic path between two rotations $\mathbf{R}_1, \mathbf{R}_2$ is given by the Riemannian distance $d_R(\mathbf{R}_1, \mathbf{R}_2)$. Thus the rotation matrix analog of the ? spherical median can be defined as

$$\tilde{\mathbf{S}}_R = \arg \min_{\mathbf{S} \in SO(3)} \sum_{i=1}^n d_R(\mathbf{R}_i, \mathbf{S}); \quad (8)$$

see also ??. We refer to this estimator of \mathbf{S} as the *geometric median*. ? offers an algorithm to find the geometric median in $SO(3)$.

4 Simulation Study

Section ?? outlines the simulation design. Section ?? briefly describes the parametric distributional models used in the study for describing symmetric rotation errors (cf. (??)) with differing variability.

Table 1: A summary of the estimators presented and their properties.

name	symbol	distance metric	cost function
Projected Arithmetic Mean	$\widehat{\mathbf{S}}_E$	Euclidean	$\sum_{i=1}^n d_E^2$
Projected Median	$\widetilde{\mathbf{S}}_E$	Euclidean	$\sum_{i=1}^n d_E$
Geometric Mean	$\widehat{\mathbf{S}}_R$	Riemannian	$\sum_{i=1}^n d_R^2$
0 Geometric Median	$\widetilde{\mathbf{S}}_R$	Riemannian	$\sum_{i=1}^n d_R$

4.1 Design of Simulation Study

To compare the performance of the proposed location estimators for determining the central direction \mathbf{S} given a sample of size n , we generated random rotation error samples $\mathbf{E}_1, \dots, \mathbf{E}_n$ in model (??) with sizes $n = 10, 50, 100$ and 300 . In (??), without loss of generality, we set the location parameter $\mathbf{S} = \mathbf{I}_{3 \times 3}$ (the identity matrix) and considered circular variances(denoted ν) of $\nu = 0.25, 0.50$ and 0.75 .

For each distribution and variability parameter, the density $f(\mathbf{E}_i|\nu)$ of a random rotation (with respect to the Haar measure) is plotted in Figure ??; the Haar measure (or uniform distribution on $SO(3)$) acts as the dominating measure for rotations and the symmetric nature of the random rotation $\mathbf{E}_i = \mathbf{E}_i(\mathbf{U}, r)$ means that its density $f(\mathbf{E}_i|\nu) = f(r|\nu)$ can be plotted in terms of the misorientation angle r of \mathbf{E}_i in (??), which is common in materials science (??).

For each combination of sample size, spread ν and choice of distribution, we generated 1,000 samples and for each sample estimated the central direction $\mathbf{S} = \mathbf{I}_{3 \times 3}$ using the four estimators on each sample. The next section continues with an introduction to the distributions under consideration.

4.2 Generating Random Rotations in the Location Model

As mentioned in the introduction, we wish to compare estimators of the (fixed) location parameter $\mathbf{S} \in SO(3)$ under three common distributional models for describing symmetric




Figure 2: Density functions for the three rotational distributions with respect to the Haar measure. The solid line corresponds to the density of matrix Fisher distribution, the dashed line the density of the circular-von Mises-based distribution and the dotted line to the density of the Cayley distribution.

rotation errors $\mathbf{E} \in SO(3)$ in a data model $\mathbf{R} = \mathbf{S}\mathbf{E}$ (cf. (??)): the symmetric matrix Fisher (????), the symmetric Cayley (??) and the circular-von Mises-based distribution (?). A general construction approach exists for random rotations that are symmetrically distributed around the identity matrix $\mathbf{I}_{3 \times 3}$; see ?? and ?. To this end, let $\mathbf{U} \in \mathbb{R}^3$ represent a point chosen uniformly on the unit sphere and, independently, generate a random angle r according to some circular density $C(r|\kappa)$ on $(-\pi, \pi]$, which is symmetric around 0 and where κ denotes a concentration parameter governing the spread of the circular distribution. Then, define a random rotation as $\mathbf{E} = \mathbf{E}(\mathbf{U}, r)$ using the constructive definition (??) (i.e., \mathbf{E} represents the position of $\mathbf{I}_{3 \times 3}$ upon rotating the standard coordinate frame in \mathbb{R}^3 about the random axis \mathbf{U} by the random angle r). The resulting rotation \mathbf{E} will be symmetrically distributed and its distributional type (i.e., matrix Fisher, Cayley or circular-von Mises-based) is determined by the form of the circular density $C(r|\kappa)$ for the (misorientation) angle r . These circular densities are given in Table ??, where $I_p(\cdot)$ denotes the Bessel function of order p defined as $I_p(\kappa) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \cos(pr) e^{\kappa \cos r} dr$. To ease the interpretation of the simulation results in Section ??, we choose not to present results in terms of κ but instead use the circular variance defined as $\nu = 1 - \rho$ as a measure of spread, where $\rho = E[\cos(r)]$. This allows us to compare the performance of the estimators for densities exhibiting the same spread. Note that $\rho = E[\cos(r)]$ is also known as the mean resultant

Table 2: Circular densities with respect to the Lebesgue measure and circular variance ν .

name	density	circular variance
Cayley	$C_C(r \kappa) = \frac{1}{\sqrt{\pi}} \frac{\Gamma(\kappa+2)}{\Gamma(\kappa+1/2)} 2^{-(\kappa+1)} (1 + \cos r)^\kappa (1 - \cos r)$	$\nu = \frac{3}{\kappa+2}$
matrix Fisher	$C_F(r \kappa) = \frac{1}{2\pi[I_0(2\kappa)-I_1(2\kappa)]} e^{2\kappa \cos(r)} [1 - \cos(r)]$	$\nu = \frac{3I_0(2\kappa)-4I_1(2\kappa)+I_2(2\kappa)}{2[I_0(2\kappa)-I_1(2\kappa)]}$
circular-von Mises	$C_M(r \kappa) = \frac{1}{2\pi I_0(\kappa)} e^{\kappa \cos(r)}$	$\nu = \frac{I_0(\kappa)-I_1(\kappa)}{I_0(\kappa)}$

length. The values of κ corresponding to the chosen circular variances are given in Table ??.

Table 3: Values of κ for each rotational distribution corresponding to the circular variances.

distribution	circular variance		
	$\nu = 0.25$	$\nu = 0.50$	$\nu = 0.75$
Cayley	10.00	4.00	2.00
matrix Fisher	3.17	1.71	1.15
circular-von Mises	2.40	1.16	0.52

In the simulations to follow, for generating random rotation errors based on the construction above, we used different samplers to randomly generate angles $r \in (-\pi, \pi]$ from a given circular density, recalling that the form of $C(r|\kappa)$ depends on the intended symmetric distribution for the rotation errors \mathbf{E} . We defer these details to the Appendix.

5 Results

In this section we summarize and present the main findings of the simulation study for estimating the central direction $\mathbf{S} = \mathbf{I}_{3 \times 3}$ with the four proposed estimators of Section ?. We quantify the estimation error between the true location $\mathbf{S} = \mathbf{I}_{3 \times 3}$ and an estimate $\hat{\mathbf{S}}$ using the geodesic distance, i.e.

$$d_R(\mathbf{S}, \hat{\mathbf{S}}) = \frac{1}{\sqrt{2}} \|\text{Log}(\mathbf{S}^\top \hat{\mathbf{S}})\|_F, \quad \text{where } \hat{\mathbf{S}} = \hat{\mathbf{S}}_E, \hat{\mathbf{S}}_R, \tilde{\mathbf{S}}_E \text{ or } \tilde{\mathbf{S}}_R. \quad (9)$$

In the appendix we show that for any two rotations \mathbf{R}_1 and \mathbf{R}_2 , $d_E(\mathbf{R}_1, \mathbf{R}_2) = 2\sqrt{2} \sin[d_R(\mathbf{R}_1, \mathbf{R}_2)/2]$. Hence, our results using d_E would prove equivalent, albeit on a smaller scale.

Figure ?? displays boxplots of the estimation errors for each of the considered rotation distributions and choice of circular spread ν for a sample of size $n = 100$. Additionally, Table ?? provides the root mean square error (RMSE) as a measure of precision as well as the *mean estimation error* for $\nu = 0.25$ in Figure ??, i.e. the top three plots. Despite skewness in some of the error distributions the *median estimation error* was quantitatively similar to the mean estimation error and is therefore not reported in Table ?? or the following.




Figure 3: Box-plots of the estimation error for each rotation distribution and level of ν , $n = 100$.

First and foremost depending on the type of distribution for the rotation errors in

(??), different location estimators emerge as preferable. For the circular-von Mises-based distribution both median-type estimators ($\tilde{\mathbf{S}}_E$ and $\tilde{\mathbf{S}}_R$) are superior with respect to the estimation error while for the Cayley and matrix Fisher models the mean-type estimators ($\hat{\mathbf{S}}_E$ and $\hat{\mathbf{S}}_R$) perform better though on a much less pronounced scale. While preferences within the median- and mean-type estimators are visible, these generally disappear as the variability in the data decreases. For the Cayley and the matrix Fisher distribution the overall pattern of estimation is very similar. $\hat{\mathbf{S}}_E$ and $\hat{\mathbf{S}}_R$ typically exhibit a little less spread and a slightly lower average estimation error than $\tilde{\mathbf{S}}_E$ and $\tilde{\mathbf{S}}_R$ but differences between all four estimators lessen as ν becomes smaller. Figure ?? further shows that the estimation error is a function of the circular spread ν ; as ν decreases the range of the observed estimation errors decreases within each rotation model and for each of the four estimators. The same is true for the mean estimation error, and RMSE in Table ??.

Table 4: Numerical summaries of estimation error for all rotational distributions, $n = 100$, $\nu = 0.25$.

estimator	Cayley		matrix Fisher		circular-von Mises	
	mean error	RMSE	mean error	RMSE	mean error	RMSE
$\hat{\mathbf{S}}_R$	0.069	0.075	0.070	0.076	0.074	0.081
$\hat{\mathbf{S}}_E$	0.070	0.076	0.070	0.076	0.062	0.067
$\tilde{\mathbf{S}}_R$	0.077	0.083	0.075	0.081	0.027	0.031
$\tilde{\mathbf{S}}_E$	0.079	0.086	0.077	0.083	0.026	0.030

Figure ?? illustrates the behavior of the estimators as a function of the sample size. Results are displayed for each level of n and $\nu = 0.75$. As to be expected, the estimation error decreases as the sample size increases for all three distributions. For small samples, $n = 10$, the estimator exhibiting the largest amount of variability is the geodesic mean $\hat{\mathbf{S}}_R$. This behavior is consistent for all three distributions. While the estimator's variability lessens considerably for the Cayley and matrix Fisher distribution as n increases, the estimator remains the most variable estimator for the circular-von Mises-based distribution. For a tabular display of the results in Figure ?? for the circular-von Mises based distribution we refer to Table ??.

Figure ?? may provide an explanation for why the circular-von Mises-based distribution




Figure 4: Box-plots of the estimation error for each rotation distribution and level of n , $\nu = 0.75$.

clearly distinguishes between the mean- and median-type estimators. Upon closer examination of the tail of its rotation density (expressed in terms of the misorientation angle r as in Figure ??) it has the heaviest tail with respect to the Haar measure, see Figure ??(b). This makes extreme observations much more likely in the circular-von Mises-based samples compared to the Cayley and matrix Fisher models.

To see if this theory holds any water in terms of the present simulation study, consider Figure ?. There we plot the difference in errors for the mean- and median-type projected estimators against the tail weight for each sample. There seems to be a positive relationship indicating as tail weight increases then the error in the mean estimator increases relative to the median estimator. Thereby confirming, atleast empirically, our theory above.

Table 5: Numerical summaries of the estimation error for all levels of n for the circular-von Mises-based distribution, $\nu = 0.75$.

n	estimator	mean error	RMSE	n	estimator	mean error	RMSE
10	$\hat{\mathbf{S}}_R$	0.652	0.789	100	$\hat{\mathbf{S}}_R$	0.204	0.222
	$\hat{\mathbf{S}}_E$	0.442	0.518		$\hat{\mathbf{S}}_E$	0.128	0.139
	$\tilde{\mathbf{S}}_R$	0.366	0.459		$\tilde{\mathbf{S}}_R$	0.069	0.079
	$\tilde{\mathbf{S}}_E$	0.326	0.450		$\tilde{\mathbf{S}}_E$	0.055	0.063
50	$\hat{\mathbf{S}}_R$	0.280	0.309	300	$\hat{\mathbf{S}}_R$	0.119	0.130
	$\hat{\mathbf{S}}_E$	0.185	0.202		$\hat{\mathbf{S}}_E$	0.075	0.081
	$\tilde{\mathbf{S}}_R$	0.109	0.130		$\tilde{\mathbf{S}}_R$	0.034	0.039
	$\tilde{\mathbf{S}}_E$	0.088	0.105		$\tilde{\mathbf{S}}_E$	0.027	0.031




Figure 5: Angular densities under consideration; $\nu = 0.75$ (a) and tail behavior (b)

We next explore the effect of the choice of geometry, Riemannian vs. Euclidean, on the estimation error for both types of loss functions. To provide more insight into possible differences we plotted the estimation error resulting from d_E versus the estimation error resulting from d_R for each loss function separately; see Figures ???. We begin with exploring both median-type estimators. For $n = 100$ and $\nu = .25$, $\nu = .75$, respectively, Figure ?? shows a scatter-plot of the projected median $\tilde{\mathbf{S}}_E$ (x -axis) versus the geometric median $\tilde{\mathbf{S}}_R$




Figure 6: A plot of the tail weight for our simulated samples versus the difference in estimator errors for each distribution.

(y -axis) for each of the three distributions. If the estimation error for a given sample agrees under both estimators the corresponding point for this sample will fall on the identity line (solid and diagonal); points below the identity line indicate less error associated with the geometric median and also show by how much the respective estimation error is smaller. For example, in Figure ??, $\tilde{\mathbf{S}}_R$ tends to yield less estimation error than $\tilde{\mathbf{S}}_E$ for the Cayley distribution as most of the points fall below the identity line while the Riemannian distance-based $\tilde{\mathbf{S}}_R$ results in greater errors for \mathbf{S} for the circular-von Mises-based distribution. This result supports results about $\tilde{\mathbf{S}}_E$ and $\tilde{\mathbf{S}}_R$ in Figure ??.

In Table ?? we support Figure ?? with an exact count (expressed as a percentage) of how often d_R resulted in a smaller estimation error than d_E . Additionally, we show the average amount by which the d_R - and d_E -based estimates deviate from one another. We denote the latter quantity by $\bar{\delta}$ in Table ?? where $\delta = d_R(\tilde{\mathbf{S}}_E, \mathbf{S}) - d_R(\tilde{\mathbf{S}}_R, \mathbf{S})$. Our previous results suggested the use of median-type estimators for the circular-von Mises-based distribution which favors $\hat{\mathbf{S}}_E$ over $\tilde{\mathbf{S}}_R$ as $d_R(\tilde{\mathbf{S}}_E, \mathbf{S}) < d_R(\tilde{\mathbf{S}}_R, \mathbf{S})$ most of the time. For the Cayley and the matrix Fisher model the preference is reversed, typically $\tilde{\mathbf{S}}_E$ exhibits a larger spread (cf. Figure ??). Note however, that both models show a marginal preference for the mean-type estimators which we will compare next in Figure ?? and Table ??.

In Figure ?? we plotted the estimation error when using $\hat{\mathbf{S}}_E$ versus the estimation error

Table 6: Average reduction in estimation error by using $\tilde{\mathbf{S}}_R$ instead of $\tilde{\mathbf{S}}_E$, $\delta = d_R(\tilde{\mathbf{S}}_E, \mathbf{S}) - d_R(\tilde{\mathbf{S}}_R, \mathbf{S})$ and percentage of samples for which $d_R(\tilde{\mathbf{S}}_R, \mathbf{S}) < d_R(\tilde{\mathbf{S}}_E, \mathbf{S})$.

		Cayley		matrix Fisher		circular- von Mises	
		$\bar{\delta}$	%	$\bar{\delta}$	%	$\bar{\delta}$	%
$\nu = 0.25$	10	0.008	0.743	0.006	0.725	-0.005	0.328
	50	0.003	0.783	0.002	0.697	-0.002	0.327
	100	0.002	0.789	0.002	0.712	-0.001	0.308
	300	0.001	0.781	0.001	0.711	-0.001	0.284
$\nu = 0.50$	10	0.031	0.772	0.017	0.662	-0.019	0.335
	50	0.013	0.811	0.005	0.620	-0.008	0.282
	100	0.009	0.809	0.005	0.660	-0.005	0.302
	300	0.005	0.804	0.002	0.658	-0.003	0.255
$\nu = 0.75$	10	0.089	0.821	0.034	0.633	-0.040	0.322
	50	0.037	0.866	0.009	0.597	-0.021	0.238
	100	0.025	0.858	0.007	0.603	-0.014	0.240
	300	0.014	0.850	0.003	0.589	-0.007	0.218

under $\hat{\mathbf{S}}_R$. Table ?? displays again the exact percentage of samples that resulted in a smaller estimation error for $\hat{\mathbf{S}}_R$ in comparison to $\hat{\mathbf{S}}_E$ as well as the average difference in the estimators' estimation errors.

Similarly to the median-type estimators, d_G is the preferred metric for the Cayley distribution especially when ν is large. For the matrix Fisher distribution the preference is less clear, especially for less variable data, but as ν increases the Euclidean-based mean yields generally a smaller estimation error more often.

In summary,

- the choice of location estimator can depend on the rotation error distribution in the location model (1). For the matrix Fisher and the Cayley distribution the projected arithmetic mean $\hat{\mathbf{S}}_E$ and the geometric mean $\hat{\mathbf{S}}_R$ are, respectively, preferable though $\tilde{\mathbf{S}}_E$ and $\tilde{\mathbf{S}}_R$ are not far behind especially when the circular spread is smaller. For the circular-von Mises-based distribution the projected median $\tilde{\mathbf{S}}_E$ should be used.
- However, a significant finding of these simulation results is that the (Euclidean-based)

Table 7: Average reduction in estimation error by using $\hat{\mathbf{S}}_R$ instead of $\hat{\mathbf{S}}_E$, $\delta = d_R(\hat{\mathbf{S}}_E, \mathbf{S}) - d_R(\hat{\mathbf{S}}_R, \mathbf{S})$ and percentage of samples for which $d_R(\hat{\mathbf{S}}_R, \mathbf{S}) < d_R(\hat{\mathbf{S}}_E, \mathbf{S})$.

Cayley				matrix Fisher		circular- von Mises	
	n	$\bar{\delta}$	%	$\bar{\delta}$	%	$\bar{\delta}$	%
$\nu = 0.25$	10	0.001	0.521	-0.002	0.450	-0.034	0.128
	50	0.001	0.531	-0.001	0.435	-0.016	0.209
	100	0.001	0.565	-0.000	0.469	-0.013	0.201
	300	0.000	0.588	-0.000	0.486	-0.007	0.239
$\nu = 0.50$	10	0.010	0.592	-0.018	0.434	-0.101	0.157
	50	0.007	0.645	-0.011	0.392	-0.055	0.145
	100	0.004	0.642	-0.007	0.393	-0.038	0.162
	300	0.003	0.642	-0.004	0.393	-0.023	0.157
$\nu = 0.75$	10	0.016	0.668	-0.096	0.357	-0.210	0.171
	50	0.026	0.741	-0.036	0.338	-0.096	0.150
	100	0.017	0.735	-0.024	0.346	-0.076	0.111
	300	0.009	0.728	-0.012	0.344	-0.045	0.127

projected median $\tilde{\mathbf{S}}_E$ is a generally good location estimator across rotation error models. For the circular-von Mises-based estimation, this generally has the best performance, while for the Cayley or matrix Fisher distributions, this estimator is often quite comparable to the best estimator. In other words, an estimator $\tilde{\mathbf{S}}_E$ not previously considered for rotation matrices in the literature appears to be generally suggestible, particularly in small samples and without knowledge of the underlying rotation error distribution.

6 Data Application

The data in our example represent the orientations of cubic crystals on a Nickel surface measured by electron backscatter diffraction. The data were obtained using a 14-fold technical replicate in each location. A central interest in EBSD data is the identification of so-called *grain maps* – grains are defined as regions on the surface with nearly identical main direction \mathbf{S} . This makes the estimation of \mathbf{S} crucial to the field.

Figure ?? shows sphere plots of data and the estimates for seven randomly chosen

locations within a single grain. Because rotation matrices are orthogonal, each of their columns (and rows) represents elements on the unit sphere. The sphere plots indicate some clustering in the data suggesting the presence of *several* main directions.

The projected median $\tilde{\mathbf{S}}_E$ is the only estimator that is not affected by this clustering in the sample. $\tilde{\mathbf{S}}_E$ identifies the main direction of the largest group within the sample.

Histograms of mis-orientation angles between each rotation in the sample and each of the four estimates of \mathbf{S} are shown in Figure ???. The distribution of the mis-orientation angles depends greatly on the choice of estimator. The only estimator with a large group of small mis-orientation angles is $\tilde{\mathbf{S}}_E$, the projected median. However, the multiple spikes in the histograms of mis-orientation angles for all of the estimators indicate a problem in estimating a single main direction.


7 Recommendations and Conclusions

To estimate the central orientation \mathbf{S} given a random sample of three-dimensional orientations from (??) the scientific literature suggests a variety of approaches. These approaches differ largely with respect to the geometry (Riemannian vs. Euclidean) in which the estimation is done, assumptions about the underlying data-generating mechanism and the choice of loss function when defining suitable estimators. The main goal of this paper was to explore the extent to which the above mentioned differences affect the estimation of \mathbf{S} . Our simulation study showed that the choice of loss function depends on the underlying data-generating mechanism. For the circular-von Mises-based model median-type estimators are recommended while for the Cayley and matrix Fisher model the mean-type estimators show less estimation error and variability. As noted in Section ?? the applied sciences generally pursue estimation of \mathbf{S} without considering the distributional underpinnings. This can be a pitfall. Restricting ourselves to the three rotation distributions under consideration, if indeed nothing is known about the underlying data-generating mechanism we suggest to use either median-type estimator, i.e. $\tilde{\mathbf{S}}_E$ or $\tilde{\mathbf{S}}_R$. The overall estimation error, even under mis-specification will be much less than the potential estimation error resulting from either mean-type estimator. The correct choice of geometry also depends on the underlying distributional model, however, the effects are not as large. The Riemannian


distance metric d_G is generally not to recommend with a mean-type estimator due to the resulting increase in variability in the estimation errors. This fact holds across all three distributions and is much expressed than for any of the other three estimators. Lastly, the extent to which all four estimators disagree relative to each other depend on the circular variance ν ; the estimators differ more when ν is large and tend to agree more and more as the data become more concentrated.

The scope of this study can be extended to other frequently encountered distributional model, especially because we restricted the simulation to symmetric perturbations around \mathbf{S} in (??). At least as important, if not even more is the extension of the studied point estimators to interval estimators. The latter requires a significant improvement in computing time before it can be practically implemented. Both of these tasks will be pursued by the authors in the near future.

Acknowledgements: We authors wish to thank ... for collecting and providing the EBSD data.



(a) $\nu = 0.25$



(b) $\nu = 0.75$

Figure 7: Comparison of the estimation errors resulting from the embedding (x-axis) and intrinsic (y-axis) approaches, $n = 100$. The mean-type estimators are in black dots and the median-type estimator is in light gray.




Figure 8: Sphere plots of EBSD measurements. The sample ($n = 98$) is taken from seven locations from within what was thought to be a single grain, but the data suggest the existence of several – very stable – main directions, as can be seen by the distinct grouping. The projected median $\tilde{\mathbf{S}}_E$ is located in the center of the largest cluster of points.




Figure 9: Histograms of the mis-orientation angles between rotational sample and estimates for a sample of size $n = 98$, combining data from seven locations. Angles are highly influenced by the choice of metric and method.

References

- K. Arun, T. Huang, and S. Blostein. Least-squares fitting of two 3-d point sets. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 9(5):698–700, 1987.
- C. Bajaj. The algebraic degree of geometric optimization problems. *Discrete & Computational Geometry*, 3(1):177–191, 1988.
- D. Best and N. Fisher. Efficient simulation of the von mises distribution. *Applied Statistics*, 28(2):152–157, 1979. ISSN 0035-9254.
- M. Bingham, D. Nordman, and S. Vardeman. Modeling and inference for measured crystal orientations and a tractable class of symmetric distributions for rotations in three dimensions. *Journal of the American Statistical Association*, 104(488):1385–1397, 2009.
- M. Bingham, D. Nordman, and S. Vardeman. Finite-sample investigation of likelihood and bayes inference for the symmetric von mises-fisher distribution. *Computational Statistics & Data Analysis*, 54(5):1317–1327, 2010.
- H. Bunge. *Texture Analysis in Material Science*. Butterworth, London, 1982.
- Y. Chan and X. He. On median-type estimators of direction for the von mises-fisher distribution. *Biometrika*, 80(4):869–875, 1993.
- Y. Chikuse. *Statistics on Special Manifolds*. Springer Verlag, 2003. ISBN 0387001603.
- Y. Dai, J. Trumpf, H. Li, N. Barnes, and R. Hartley. Rotation averaging with application to camera-rig calibration. *Computer Vision–ACCV 2009*, pages 335–346, 2010.
- T. Downs. Orientation statistics. *Biometrika*, 59(3):665–676, 1972.
- G. Ducharme and P. Milasevic. Spatial median and directional data. *Biometrika*, 74(1):212–215, 1987.
- S. Durocher and D. Kirkpatrick. The projection median of a set of points. *Computational Geometry*, 42(5):364–375, 2009.

- N. Fisher. Spherical Medians. *Journal of the Royal Statistical Society. Series B (Methodological)*, pages 342–348, 1985.
- R. Fisher. Dispersion on a Sphere. *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, pages 295–305, 1953. ISSN 0080-4630.
- P. Fletcher, S. Venkatasubramanian, and S. Joshi. Robust Statistics on Riemannian Manifolds via the Geometric Median. In *2008 IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, pages 1–8. IEEE, 2008.
- P. Fletcher, S. Venkatasubramanian, and S. Joshi. The geometric median on riemannian manifolds with application to robust atlas estimation. *NeuroImage*, 45(1):S143–S152, 2009.
- J. Gower. The mediancentre. *Applied Statistics*, 23(3):466–470, 1974.
- R. Hadani and A. Singer. Representation theoretic patterns in three-dimensional cryo-electron microscopy ii – the class averaging problem. *Foundations of Computational Mathematics*, pages 1–28, 2011.
- R. Hartley, K. Aftab, and J. Trumpf. L1 rotation averaging using the weiszfeld algorithm. In *2011 IEEE Conference on Computer Vision and Pattern Recognition (CVPR)*, pages 3041–3048. IEEE, 2011.
- R. Hielscher, H. Schaeben, and S. H. Orientation distribution within a single hematite crystal. *Mathematical Geosciences*, 42:359–375, 2010.
- B. Horn, H. Hilden, and S. Negahdaripour. Closed-form solution of absolute orientation using orthonormal matrices. *Journal of the Optical Society of America*, 5(7):1127–1135, 1988.
- P. Jupp and K. Mardia. Maximum likelihood estimators for the matrix von Mises-Fisher and Bingham distributions. *The Annals of Statistics*, 7(3):599–606, 1979. ISSN 0090-5364.

- P. Jupp and K. Mardia. A unified view of the theory of directional statistics. *International Statistical Review*, 57(3):261–294, 1989.
- H. Karcher. Riemannian center of mass and mollifier smoothing. *Communications on pure and applied mathematics*, 30(5):509–541, 1977.
- C. Khatri and K. Mardia. The von mises-fisher matrix distribution in orientation statistics. *Journal of the Royal Statistical Society. Series B (Methodological)*, 39(1):95–106, 1977. ISSN 0035-9246.
- P. Langevin. Magnetism and the theory of the electron. *Annales de chimie et de physique*, 5:70, 1905.
- C. León, J. Massé, and L. Rivest. A statistical model for random rotations. *Journal of Multivariate Analysis*, 97(2):412–430, 2006.
- R. Liu and K. Singh. Ordering directional data: concepts of data depth on circles and spheres. *The Annals of Statistics*, pages 1468–1484, 1992. ISSN 0090-5364.
- J. Manton. A globally convergent numerical algorithm for computing the centre of mass on compact lie groups. In *8th Conference on Control, Automation, Robotics and Vision, (ICARCV)*, volume 3, pages 2211–2216. IEEE, 2004.
- K. Mardia and P. Jupp. *Directional Statistics*. Wiley Chichester, 2000.
- S. Matthies, J. Muller, and G. Vinel. On the Normal Distribution in the Orientation Space. *Textures and Microstructures*, 10:77–96, 1988.
- M. Moakher. Means and averaging in the group of rotations. *SIAM Journal on Matrix Analysis and Applications*, 24(1):1–16, 2002.
- M. Pierrynowski and K. Ball. Oppugning the assumptions of spatial averaging of segment and joint orientations. *Journal of biomechanics*, 42(3):375–378, 2009.
- D. Rancourt, L. Rivest, and J. Asselin. Using orientation statistics to investigate variations in human kinematics. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 49(1):81–94, 2000. ISSN 1467-9876.

- V. Randle. *Microtexture Determination and its Applications*. London: Maney for The Institute of Materials, Minerals and Mining, 2003.
- T. Savyolova and D. Nikolayev. Normal Distribution on the Rotation Group $SO(3)$. *Textures and Microstructures*, 29:201–233, 1995.
- H. Schaeben. A simple standard orientation density function: The hyperspherical de la vallée poussin kernel. *Phys. Stat. Sol. (B)*, 200:367–376, 1997.
- M. Schwartz and A. Rozumalski. A new method for estimating joint parameters from motion data. *Journal of Biomechanics*, 38(1):107–116, 2005.
- S. Umeyama. Least squares estimation of transformation parameters between two point patterns. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 13:376–380, 1991.
- G. Watson. Statistics on spheres, university of arkansas lecture notes in the mathematical sciences, vol. 6, 1983.
- E. Weiszfeld. Sur le point pour lequel la somme des distances de n points donnés est minimum. *Tohoku Mathematics Journal*, 43:355–386, 1937.

A Sampling Process

A.1 circular-von Mises-based distribution

To simulate a set of random rotations from the circular-von Mises-based distribution we follow the algorithm proposed by ?. The algorithm is available in the IMSL Library (1991) and is implemented as follows. Let $\mu = 0$ denote the mean of the target angular distribution and κ its concentration parameter. We define constants a, b and d as $a \equiv 1 + \sqrt{1 + 4\kappa^2}$, $b \equiv (a - \sqrt{2a})$, $d \equiv (1 + b^2)/2b$. In steps one, two and four we generate three new observations u_1 , u_2 and u_3 , each from a uniform distribution defined over the interval $(0, 1)$.

1. Set $z = \cos(\pi u_1)$, $f = (1 + dz)/(z + d)$ and $c = \kappa(d - f)$.
2. If $c(2 - c) - u_2 > 0$ go to step 4.
3. If $\log(c/u_2) + 1 - c < 0$ return to step 1.
4. Set $r = \text{sign}(u_3 - 0.5) \cos^{-1}(f)$
5. Then r follows a von Mises (κ) distribution.

A.2 Cayley distribution

To simulate rotation matrices from a Cayley distribution we make use of a result given in ?. If the angle r follows a Cayley distribution it holds that $\frac{1+\cos r}{2} \sim \text{Beta}(\kappa + 1/2, 3/2)$. Hence, angles according to a Cayley distribution can be simulated through composition: we simulate a Bernoulli trial Y with outcomes -1 and 1 having probability 0.5 and an observation X from a $\text{Beta}(\kappa + 1/2, 3/2)$ distribution and then set $r = \frac{Y}{2} \cos^{-1}(2X - 1)$.

A.3 matrix Fisher distribution

Simulation from the matrix Fisher distribution is achieved through a rejection algorithm. Let $C_F(r|\kappa)$ denote the matrix von Mises-Fisher density as given in Table ?? and $Y \sim \text{Uniform}(-\pi, \pi]$.

1. Define $M = \frac{1}{2\kappa} e^{2\kappa-1} \frac{1}{\mathbf{I}_0(2\kappa) - \mathbf{I}_1(2\kappa)}$
2. Generate $U \sim \text{Uniform}(0, 1)$ and $Y \sim \text{Uniform}(-\pi, \pi]$, where U and Y are independent.
3. If $U < \frac{1}{M} C_F(Y|\kappa)$, accept Y ; otherwise return to step (2)

Given a set of randomly generated angles r_1, \dots, r_n we can now generate the corresponding set of rotation matrices as follows:

1. Generate a point uniformly on the unit sphere


$$\mathbf{U} = (u_1, u_2, u_3)^\top = (\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)^\top$$

where $0 \leq \theta \leq \pi$ and $0 \leq \phi \leq 2\pi$.

2. Given an angle of rotation, r_i , generated as described above from an angular distribution symmetric about 0 and with concentration κ rotate \mathbf{I} about \mathbf{U} by r_i radians.

A.4 Comparison of Euclidean and Riemannian Metrics When Measuring Distance to the Central Direction

The results section of this paper only compared the distances of each estimator from the central direction in terms of Riemannian distance. This section is intended to show that a comparison based on the Euclidean distance will not lead to different results. Recall that the Riemannian distance between two rotations is the shortest geodesic curve that connects those two rotations and the Euclidean distance is the shortest cord that connects those two rotations. Refer to the picture below for a two-dimensional simplification of the idea.



Let $\mathbf{R}_1\mathbf{v}$ and $\mathbf{R}_2\mathbf{v}$ be two observations in $SO(2)$. The Riemannian distance between these two observations is given by α and is indicated with the thick black arc. The Euclidean distance is given by β and is indicated by the gray line running through the circle. Because this is the unit circle, the Riemannian distance is also the angle in the center of the circle thus, using basic geometry its clear that half of the Euclidean distance is the sine of half of the Riemannian distance, i.e.

$$\beta = 2 \sin \left(\frac{\alpha}{2} \right).$$

Extending this to $SO(3)$ gives the following proposition: for all \mathbf{R}_1 and \mathbf{R}_2 in $SO(3)$

$$d_E(\mathbf{R}_1, \mathbf{R}_2) = 2^{3/2} \sin \left(\frac{d_R(\mathbf{R}_1, \mathbf{R}_2)}{2} \right). \quad (10)$$

Proof:

Let $\mathbf{o}_1, \mathbf{R}_2 \in SO(3)$ be given and define $\text{tr}(\mathbf{R}_1^\top \mathbf{R}_2) = 1 + 2 \cos(\theta)$ then $|\theta| = d_R(\mathbf{R}_1, \mathbf{R}_2)$. Notice that (??) holds trivially for $d_E(\mathbf{R}_1, \mathbf{R}_2) = d_R(\mathbf{R}_1, \mathbf{R}_2) = 0$ so consider the case

$|\theta| > 0$. By definition of $d_E(\mathbf{R}_1, \mathbf{R}_2)$ we have the following:

$$\begin{aligned}
d_E(\mathbf{R}_1, \mathbf{R}_2)^2 &= \|\mathbf{R}_1 - \mathbf{R}_2\|_F^2 \\
&= \text{tr}[(\mathbf{R}_1 - \mathbf{R}_2)^\top (\mathbf{R}_1 - \mathbf{R}_2)] = \text{tr}[(\mathbf{R}_1^\top - \mathbf{R}_2^\top)(\mathbf{R}_1 - \mathbf{R}_2)] \\
&= \text{tr}[\mathbf{R}_1^\top \mathbf{R}_1 + \mathbf{R}_2^\top \mathbf{R}_2 - \mathbf{R}_2^\top \mathbf{R}_1 - \mathbf{R}_1^\top \mathbf{R}_2] \\
&= \text{tr}[2\mathbf{I} - \mathbf{R}_2^\top \mathbf{R}_1 - \mathbf{R}_1^\top \mathbf{R}_2] \\
&= 2\text{tr}(\mathbf{I}) - \text{tr}(\mathbf{R}_2^\top \mathbf{R}_1) - \text{tr}(\mathbf{R}_1^\top \mathbf{R}_2) \\
&= 6 - 2\text{tr}(\mathbf{R}_1^\top \mathbf{R}_2) = 6 - 2(1 + 2\cos(\theta)) \\
&= 6 - 2 - 4\cos(\theta) = 8\left(\frac{1 - \cos(\theta)}{2}\right) \\
&= 8\sin^2\left(\frac{\theta}{2}\right) \\
&= \left[2^{3/2}\sin\left(\frac{|\theta|}{2}\right)\right]^2 \\
&= \left[2^{3/2}\sin\left(\frac{d_R(\mathbf{R}_1, \mathbf{R}_2)}{2}\right)\right]^2
\end{aligned}$$

Taking square roots on both sides and noticing that $d_E(\mathbf{R}_1, \mathbf{R}_2) \geq 0$ and $d_R(\mathbf{R}_1, \mathbf{R}_2) \geq 0$ gives (??).

□