

Duality

Lecture 4

October 1, 2025

Motivation

Consider an optimization problem

$$\begin{aligned} &\text{minimize } c^T x \\ &\text{such that } Ax \leq b. \end{aligned}$$

Motivation

Consider an optimization problem

$$\begin{aligned} &\text{minimize } c^T x \\ &\text{such that } Ax \leq b. \end{aligned}$$

1. *Given a feasible x , how can we know “how good” it is?*

Formally, how to quantify the gap $c^T x - p^*$ where p^* is the optimal value?

Motivation

Consider an optimization problem

$$\begin{aligned} &\text{minimize } c^T x \\ &\text{such that } Ax \leq b. \end{aligned}$$

1. *Given a feasible x , how can we know “how good” it is?*

Formally, how to quantify the gap $c^T x - p^*$ where p^* is the optimal value?

2. *Without a feasible x , how to **certify** that $\{x : Ax \leq b\}$ is empty?*

Motivation

Consider an optimization problem

$$\begin{aligned} &\text{minimize } c^T x \\ &\text{such that } Ax \leq b. \end{aligned}$$

1. *Given a feasible x , how can we know “how good” it is?*

Formally, how to quantify the gap $c^T x - p^*$ where p^* is the optimal value?

2. *Without a feasible x , how to **certify** that $\{x : Ax \leq b\}$ is empty?*

3. *Suppose one constraint is: $a_i^T x \leq 0$ where $a_i \in \mathcal{A}$ are unknown parameters. How to find an x that is feasible **for any** $a_i \in \mathcal{A}$?*

Motivation

Consider an optimization problem

$$\begin{aligned} &\text{minimize } c^T x \\ &\text{such that } Ax \leq b. \end{aligned}$$

1. *Given a feasible x , how can we know “how good” it is?*

Formally, how to quantify the gap $c^T x - p^*$ where p^* is the optimal value?

2. *Without a feasible x , how to **certify** that $\{x : Ax \leq b\}$ is empty?*

3. Suppose one constraint is: $a_i^T x \leq 0$ where $a_i \in \mathcal{A}$ are unknown parameters. *How to find an x that is feasible **for any** $a_i \in \mathcal{A}$?*

4. You are offered a bit more of b_i , for a “suitable price”. *Is the deal worthwhile?*

Duality theory will provide answers to these questions (and more)

Outline

- Consider a **primal** optimization problem:

$$\begin{aligned} (\mathcal{P}) \quad & \text{minimize } c^T x \\ & \text{such that } Ax \leq b. \end{aligned}$$

Outline

- Consider a **primal** optimization problem:

$$\begin{aligned} (\mathcal{P}) \quad & \text{minimize } c^T x \\ & \text{such that } Ax \leq b. \end{aligned}$$

- We will form a **dual** problem; also a linear program (LP):

$$\begin{aligned} (\mathcal{D}) \quad & \text{maximize } \tilde{r}^T y \\ & \text{such that } \tilde{A}y \leq \tilde{b}. \end{aligned}$$

Outline

- Consider a **primal** optimization problem:

$$\begin{aligned} (\mathcal{P}) \quad & \text{minimize } c^T x \\ & \text{such that } Ax \leq b. \end{aligned}$$

- We will form a **dual** problem; also a linear program (LP):

$$\begin{aligned} (\mathcal{D}) \quad & \text{maximize } \tilde{r}^T y \\ & \text{such that } \tilde{A}y \leq \tilde{b}. \end{aligned}$$

- We will show that the dual provides lower bounds for the primal:

$$\tilde{r}^T y \leq c^T x \quad \text{for any } x \text{ feasible for } (\mathcal{P}) \text{ and } y \text{ feasible for } (\mathcal{D})$$

Outline

- Consider a **primal** optimization problem:

$$\begin{aligned} (\mathcal{P}) \quad & \text{minimize } c^T x \\ & \text{such that } Ax \leq b. \end{aligned}$$

- We will form a **dual** problem; also a linear program (LP):

$$\begin{aligned} (\mathcal{D}) \quad & \text{maximize } \tilde{r}^T y \\ & \text{such that } \tilde{A}y \leq \tilde{b}. \end{aligned}$$

- We will show that the dual provides lower bounds for the primal:

$$\tilde{r}^T y \leq c^T x \quad \text{for any } x \text{ feasible for } (\mathcal{P}) \text{ and } y \text{ feasible for } (\mathcal{D})$$

- If (\mathcal{P}) has optimal solution x^* , then (\mathcal{D}) has optimal solution y^* and

$$c^T x^* = \tilde{r}^T y^* \quad \textbf{(strong duality)}$$

Outline

- Consider a **primal** optimization problem:

$$\begin{aligned} (\mathcal{P}) \quad & \text{minimize } c^T x \\ & \text{such that } Ax \leq b. \end{aligned}$$

- We will form a **dual** problem; also a linear program (LP):

$$\begin{aligned} (\mathcal{D}) \quad & \text{maximize } \tilde{r}^T y \\ & \text{such that } \tilde{A}y \leq \tilde{b}. \end{aligned}$$

- We will show that the dual provides lower bounds for the primal:

$$\tilde{r}^T y \leq c^T x \quad \text{for any } x \text{ feasible for } (\mathcal{P}) \text{ and } y \text{ feasible for } (\mathcal{D})$$

- If (\mathcal{P}) has optimal solution x^* , then (\mathcal{D}) has optimal solution y^* and

$$c^T x^* = \tilde{r}^T y^* \quad \text{(strong duality)}$$

- In the process, will uncover some **fundamental ideas in optimization**:

separation of convex sets \implies Farkas Lemma \implies strong duality

Deriving Lower Bounds

Consider a linear optimization problem in the most general form possible:

Primal Problem

$$\begin{array}{ll} (\mathcal{P}) \text{ minimize}_x & c^T x \\ \text{such that} & a_i^T x \geq b_i, \quad \forall i \in I_{ge}, \\ & a_i^T x \leq b_i, \quad \forall i \in I_{le}, \\ & a_i^T x = b_i, \quad \forall i \in I_{eq}, \\ & x_j \geq 0, \quad \forall j \in J_p, \\ & x_j \leq 0, \quad \forall j \in J_n, \\ & x_j \text{ free}, \quad \forall j \in J_f \\ \text{variable} & x \in \mathbb{R}^n. \end{array} \tag{1}$$

Note the mnemonic encoding...

Deriving Lower Bounds

Consider a linear optimization problem in the most general form possible:

Primal Problem

$$\begin{array}{ll} (\mathcal{P}) \text{ minimize}_x & c^T x \\ \text{such that} & a_i^T x \geq b_i, \quad \forall i \in I_{ge}, \\ & a_i^T x \leq b_i, \quad \forall i \in I_{le}, \\ & a_i^T x = b_i, \quad \forall i \in I_{eq}, \\ & x_j \geq 0, \quad \forall j \in J_p, \\ & x_j \leq 0, \quad \forall j \in J_n, \\ & x_j \text{ free}, \quad \forall j \in J_f \\ \text{variable} & x \in \mathbb{R}^n. \end{array} \quad \begin{array}{l} A = \begin{bmatrix} A^{le} \\ A^{ge} \\ A^{eq} \end{bmatrix} \in \mathbb{R}^{m \times n} \\ b = \begin{bmatrix} b^{le} \\ b^{ge} \\ b^{eq} \end{bmatrix} \in \mathbb{R}^m \end{array} \quad (1)$$

Note the mnemonic encoding...

Deriving Lower Bounds

Consider a linear optimization problem in the most general form possible:

Primal Problem

$$\begin{aligned} (\mathcal{P}) \quad & \text{minimize}_x \quad c^T x \\ & \text{such that} \quad \begin{aligned} a_i^T x &\geq b_i, & \forall i \in I_{ge}, \\ a_i^T x &\leq b_i, & \forall i \in I_{le}, \\ a_i^T x &= b_i, & \forall i \in I_{eq}, \\ x_j &\geq 0, & \forall j \in J_p, \\ x_j &\leq 0, & \forall j \in J_n, \\ x_j &\text{ free}, & \forall j \in J_f \end{aligned} \\ & \text{variable} \quad x \in \mathbb{R}^n. \end{aligned} \quad \begin{aligned} A &= \begin{bmatrix} A^{le} \\ A^{ge} \\ A^{eq} \end{bmatrix} \in \mathbb{R}^{m \times n} \\ b &= \begin{bmatrix} b^{le} \\ b^{ge} \\ b^{eq} \end{bmatrix} \in \mathbb{R}^m \end{aligned} \quad (1)$$

Note the mnemonic encoding...

Definition

We will refer to this as the **primal problem** or problem (\mathcal{P}) .

Let P denote its feasible set (a polyhedron), and p^* denote its optimal value.

Deriving Lower Bounds

Consider the primal problem:

$$\begin{array}{ll} (\mathcal{P}) \text{ minimize}_x & c^T x \\ \text{such that} & a_i^T x \geq b_i, \quad \forall i \in I_{\text{ge}}, \\ & a_i^T x \leq b_i, \quad \forall i \in I_{\text{le}}, \\ & a_i^T x = b_i, \quad \forall i \in I_{\text{eq}}, \\ & x_j \geq 0, \quad \forall j \in J_p, \\ & x_j \leq 0, \quad \forall j \in J_n, \\ & x_j \text{ free}, \quad \forall j \in J_f \\ \text{variable} & x \in \mathbb{R}^n. \end{array}$$

(\mathcal{P}) is a minimization; we seek **valid lower bounds** on (\mathcal{P}) . *Any ideas?*

Deriving Lower Bounds

Consider the primal problem:

$$\begin{array}{ll} (\mathcal{P}) \text{ minimize}_x & c^T x \\ \text{such that} & a_i^T x \geq b_i, \quad \forall i \in I_{\text{ge}}, \\ & a_i^T x \leq b_i, \quad \forall i \in I_{\text{le}}, \\ & a_i^T x = b_i, \quad \forall i \in I_{\text{eq}}, \\ & x_j \geq 0, \quad \forall j \in J_p, \\ & x_j \leq 0, \quad \forall j \in J_n, \\ & x_j \text{ free}, \quad \forall j \in J_f \\ \text{variable} & x \in \mathbb{R}^n. \end{array}$$

(\mathcal{P}) is a minimization; we seek **valid lower bounds** on (\mathcal{P}) . *Any ideas?*

Can **remove** constraints! Drastic, and could end up with a bound of $-\infty$!

Deriving Lower Bounds

Consider the primal problem:

$$\begin{array}{ll} (\mathcal{P}) \text{ minimize}_x & c^T x \\ \text{such that} & a_i^T x \geq b_i, \quad \forall i \in I_{ge}, \\ & a_i^T x \leq b_i, \quad \forall i \in I_{le}, \\ & a_i^T x = b_i, \quad \forall i \in I_{eq}, \\ & x_j \geq 0, \quad \forall j \in J_p, \\ & x_j \leq 0, \quad \forall j \in J_n, \\ & x_j \text{ free}, \quad \forall j \in J_f \\ \text{variable} & x \in \mathbb{R}^n. \end{array}$$

(\mathcal{P}) is a minimization; we seek **valid lower bounds** on (\mathcal{P}) . *Any ideas?*

Can **remove** constraints! Drastic, and could end up with a bound of $-\infty$!

Let's **relax** some constraints and penalize ourselves for the relaxation! *Which / how?*

Deriving Lower Bounds

Consider the primal problem:

$$\begin{array}{ll} (\mathcal{P}) \text{ minimize}_x & c^T x \\ \text{such that} & a_i^T x \geq b_i, \quad \forall i \in I_{ge}, \\ & a_i^T x \leq b_i, \quad \forall i \in I_{le}, \\ & a_i^T x = b_i, \quad \forall i \in I_{eq}, \\ & x_j \geq 0, \quad \forall j \in J_p, \\ & x_j \leq 0, \quad \forall j \in J_n, \\ & x_j \text{ free}, \quad \forall j \in J_f \\ \text{variable} & x \in \mathbb{R}^n. \end{array}$$

(\mathcal{P}) is a minimization; we seek **valid lower bounds** on (\mathcal{P}) . *Any ideas?*

Can **remove** constraints! Drastic, and could end up with a bound of $-\infty$!

Let's **relax** some constraints and penalize ourselves for the relaxation! *Which / how?*

General principle: (i) relax “complicating” constraints; (ii) try “simple” penalty

Deriving Lower Bounds

Consider the primal problem:

$$\begin{array}{ll} (\mathcal{P}) \text{ minimize}_x & c^T x \\ & (\lambda_i \rightarrow) \quad a_i^T x \geq b_i, \quad \forall i \in I_{\text{ge}}, \\ & (\lambda_i \rightarrow) \quad a_i^T x \leq b_i, \quad \forall i \in I_{\text{le}}, \\ & (\lambda_i \rightarrow) \quad a_i^T x = b_i, \quad \forall i \in I_{\text{eq}}, \\ & x_j \geq 0, \quad \forall j \in J_p, \\ & x_j \leq 0, \quad \forall j \in J_n, \\ & x_j \text{ free}, \quad \forall j \in J_f. \end{array}$$

For every constraint i , have a **penalty** λ_i

Construct the **lower bound** as the **Lagrangian**:

$$\mathcal{L}(x, \lambda) = c^T x - \sum_{i=1}^m \lambda_i (a_i^T x - b_i) = c^T x - \lambda^T (Ax - b)$$

Deriving Lower Bounds

Consider the primal problem:

$$\begin{aligned} (\mathcal{P}) \text{ minimize}_x \quad & c^T x \\ & (\lambda_i \rightarrow) \quad a_i^T x \geq b_i, \quad \forall i \in I_{ge}, \\ & (\lambda_i \rightarrow) \quad a_i^T x \leq b_i, \quad \forall i \in I_{le}, \\ & (\lambda_i \rightarrow) \quad a_i^T x = b_i, \quad \forall i \in I_{eq}, \\ & x_j \geq 0, \quad \forall j \in J_p, \\ & x_j \leq 0, \quad \forall j \in J_n, \\ & x_j \text{ free}, \quad \forall j \in J_f. \end{aligned}$$

For every constraint i , have a **penalty** λ_i

Construct the **lower bound** as the **Lagrangian**:

$$\mathcal{L}(x, \lambda) = c^T x - \sum_{i=1}^m \lambda_i (a_i^T x - b_i) = c^T x - \lambda^T (Ax - b)$$

Note: we relaxed the complicating constraints, $a_i^T x \text{ (?) } b_i$, and used a linear penalty

Not apriori clear that this will give us very good bounds...

Deriving Lower Bounds

Consider the primal problem:

$$\begin{array}{ll} (\mathcal{P}) \text{ minimize}_x & c^T x \\ & (\lambda_i \rightarrow) \quad a_i^T x \geq b_i, \quad \forall i \in I_{ge}, \\ & (\lambda_i \rightarrow) \quad a_i^T x \leq b_i, \quad \forall i \in I_{le}, \\ & (\lambda_i \rightarrow) \quad a_i^T x = b_i, \quad \forall i \in I_{eq}, \\ & x_j \geq 0, \quad \forall j \in J_p, \\ & x_j \leq 0, \quad \forall j \in J_n, \\ & x_j \text{ free}, \quad \forall j \in J_f. \end{array}$$

We want the Lagrangean to give us **a valid lower bound**:

$$\mathcal{L}(x, \lambda) = c^T x - \lambda^T (Ax - b) \leq c^T x, \quad \forall x \in P.$$

Deriving Lower Bounds

Consider the primal problem:

$$\begin{array}{ll} (\mathcal{P}) \text{ minimize}_x & c^T x \\ & (\lambda_i \rightarrow) \quad a_i^T x \geq b_i, \quad \forall i \in I_{ge}, \\ & (\lambda_i \rightarrow) \quad a_i^T x \leq b_i, \quad \forall i \in I_{le}, \\ & (\lambda_i \rightarrow) \quad a_i^T x = b_i, \quad \forall i \in I_{eq}, \\ & x_j \geq 0, \quad \forall j \in J_p, \\ & x_j \leq 0, \quad \forall j \in J_n, \\ & x_j \text{ free}, \quad \forall j \in J_f. \end{array}$$

We want the Lagrangean to give us **a valid lower bound**:

$$\mathcal{L}(x, \lambda) = c^T x - \lambda^T (Ax - b) \leq c^T x, \quad \forall x \in P.$$

We must impose constraints on λ :

Deriving Lower Bounds

Consider the primal problem:

$$\begin{aligned} (\mathcal{P}) \text{ minimize}_x \quad & c^T x \\ & (\lambda_i \rightarrow) \quad a_i^T x \geq b_i, \quad \forall i \in I_{ge}, \\ & (\lambda_i \rightarrow) \quad a_i^T x \leq b_i, \quad \forall i \in I_{le}, \\ & (\lambda_i \rightarrow) \quad a_i^T x = b_i, \quad \forall i \in I_{eq}, \\ & x_j \geq 0, \quad \forall j \in J_p, \\ & x_j \leq 0, \quad \forall j \in J_n, \\ & x_j \text{ free}, \quad \forall j \in J_f. \end{aligned}$$

We want the Lagrangean to give us **a valid lower bound**:

$$\mathcal{L}(x, \lambda) = c^T x - \lambda^T (Ax - b) \leq c^T x, \forall x \in P.$$

We must impose constraints on λ :

$$\left. \begin{aligned} \lambda_i &\geq 0, & \forall i \in I_{ge} \\ \lambda_i &\leq 0, & \forall i \in I_{le} \\ \lambda_i &\text{ free}, & \forall i \in I_{eq}. \end{aligned} \right\} \Leftrightarrow \lambda \in \Lambda \quad (2)$$

Deriving Lower Bounds

Summarizing... any $\lambda \in \Lambda$ produces a **valid lower bound**:

$$\mathcal{L}(x, \lambda) = c^T x - \lambda^T (Ax - b) \leq c^T x, \forall x \in P.$$

*How can we get a lower bound on the primal's **optimal value** p^* ?*

Deriving Lower Bounds

Summarizing... any $\lambda \in \Lambda$ produces a **valid lower bound**:

$$\mathcal{L}(x, \lambda) = c^T x - \lambda^T (Ax - b) \leq c^T x, \forall x \in P.$$

*How can we get a lower bound on the primal's **optimal value** p^* ?*

Claim

The function $g : \Lambda \rightarrow \mathbb{R}$ defined as:

$$\begin{aligned} g(\lambda) &:= \min_x \mathcal{L}(x, \lambda) \\ &\text{s.t. } x_j \geq 0, \forall j \in J_p \\ &\quad x_j \leq 0, \forall j \in J_n \\ &\quad x_j \text{ free}, \forall j \in J_f \end{aligned} \tag{3}$$

satisfies $g(\lambda) \leq p^$ for any $\lambda \in \Lambda$.*

Note: including the sign constraints on x in this optimization improves the lower bound!

Deriving Lower Bounds

Let us analyze this further:

$$\begin{aligned} g(\lambda) = \min_x \mathcal{L}(x, \lambda) &= \min_x [\lambda^T b + (c^T - \lambda^T A)x] \\ \text{s.t. } x_j &\geq 0, \forall j \in J_p, & \text{s.t. } x_j &\geq 0, \forall j \in J_p, \\ x_j &\leq 0, \forall j \in J_n, & x_j &\leq 0, \forall j \in J_n, \\ x_j &\text{ free}, \forall j \in J_f & x_j &\text{ free}, \forall j \in J_f \end{aligned}$$

Deriving Lower Bounds

Let us analyze this further:

$$\begin{aligned} g(\lambda) = \min_x \mathcal{L}(x, \lambda) &= \min_x [\lambda^T b + (c^T - \lambda^T A)x] \\ \text{s.t. } x_j &\geq 0, \forall j \in J_p, & \text{s.t. } x_j &\geq 0, \forall j \in J_p, \\ x_j &\leq 0, \forall j \in J_n, & x_j &\leq 0, \forall j \in J_n, \\ x_j &\text{ free}, \forall j \in J_f & x_j &\text{ free}, \forall j \in J_f \end{aligned}$$

$$g(\lambda) = \begin{cases} \lambda^T b, & \text{if } \lambda^T A_j \leq c_j, \forall j \in J_p \text{ and } \lambda^T A_j \geq c_j, \forall j \in J_n \text{ and } \lambda^T A_j = c_j, \forall j \in J_f \\ -\infty, & \text{otherwise.} \end{cases}$$

Deriving the Dual Problem

$$g(\lambda) = \begin{cases} \lambda^T b, & \text{if } \lambda^T A_j \leq c_j, \forall j \in J_p \text{ and } \lambda^T A_j \geq c_j, \forall j \in J_n \text{ and } \lambda^T A_j = c_j, \forall j \in J_f \\ -\infty, & \text{otherwise.} \end{cases}$$

is a **valid lower bound** on the primal **optimal value**: $g(\lambda) \leq p^*$ for any $\lambda \in \Lambda$.

*How can we get the **best** lower bound?*

Deriving the Dual Problem

$$g(\lambda) = \begin{cases} \lambda^T b, & \text{if } \lambda^T A_j \leq c_j, \forall j \in J_p \text{ and } \lambda^T A_j \geq c_j, \forall j \in J_n \text{ and } \lambda^T A_j = c_j, \forall j \in J_f \\ -\infty, & \text{otherwise.} \end{cases}$$

is a **valid lower bound** on the primal **optimal value**: $g(\lambda) \leq p^*$ for any $\lambda \in \Lambda$.

*How can we get the **best** lower bound?*

$$\underset{\lambda \in \Lambda}{\text{maximize}} \ g(\lambda) \tag{4}$$

This is equivalent to the following optimization problem:

Deriving the Dual Problem

$$g(\lambda) = \begin{cases} \lambda^T b, & \text{if } \lambda^T A_j \leq c_j, \forall j \in J_p \text{ and } \lambda^T A_j \geq c_j, \forall j \in J_n \text{ and } \lambda^T A_j = c_j, \forall j \in J_f \\ -\infty, & \text{otherwise.} \end{cases}$$

is a **valid lower bound** on the primal **optimal value**: $g(\lambda) \leq p^*$ for any $\lambda \in \Lambda$.

*How can we get the **best** lower bound?*

$$\underset{\lambda \in \Lambda}{\text{maximize}} \quad g(\lambda) \tag{4}$$

This is equivalent to the following optimization problem:

Dual Problem

$$\begin{aligned} & \text{maximize} && \lambda^T b \\ & \text{subject to} && \lambda_i \geq 0, && \forall i \in I_{ge}, \\ & && \lambda_i \leq 0, && \forall i \in I_{le}, \\ & && \lambda_i \text{ free}, && \forall i \in I_{eq}, \\ & && \lambda^T A_j \leq c_j, && \forall j \in J_p, \\ & && \lambda^T A_j \geq c_j, && \forall j \in J_n, \\ & && \lambda^T A_j = c_j, && \forall j \in J_f. \end{aligned} \tag{5}$$

Deriving the Dual Problem

Dual Problem

$$\begin{array}{lll} \text{maximize} & \lambda^T b & \\ \text{subject to} & \lambda_i \geq 0, & \forall i \in I_{\text{ge}}, \\ & \lambda_i \leq 0, & \forall i \in I_{\text{le}}, \\ & \lambda_i \text{ free}, & \forall i \in I_{\text{eq}}, \\ & \lambda^T A_j \leq c_j, & \forall j \in J_p, \\ & \lambda^T A_j \geq c_j, & \forall j \in J_n, \\ & \lambda^T A_j = c_j, & \forall j \in J_f. \end{array} \quad (6)$$

Deriving the Dual Problem

Dual Problem

$$\begin{array}{ll} \text{maximize} & \lambda^T b \\ \text{subject to} & \lambda_i \geq 0, \quad \forall i \in I_{\text{ge}}, \\ & \lambda_i \leq 0, \quad \forall i \in I_{\text{le}}, \\ & \lambda_i \text{ free}, \quad \forall i \in I_{\text{eq}}, \\ & \lambda^T A_j \leq c_j, \quad \forall j \in J_p, \\ & \lambda^T A_j \geq c_j, \quad \forall j \in J_n, \\ & \lambda^T A_j = c_j, \quad \forall j \in J_f. \end{array} \quad (6)$$

Definition

This is the **dual** of (\mathcal{P}) , which we will also refer to as (\mathcal{D}) . We denote its feasible set with D and its optimal value with d^* .

Note: The dual is also a linear optimization problem!

Primal-Dual Pair

Primal-Dual Pair of Problems

Primal (\mathcal{P})

minimize $c^T x$

$(\lambda_i \rightarrow) \quad a_i^T x \geq b_i, \quad \forall i \in I_{ge}$

$(\lambda_i \rightarrow) \quad a_i^T x \leq b_i, \quad \forall i \in I_{le}$

$(\lambda_i \rightarrow) \quad a_i^T x = b_i, \quad \forall i \in I_{eq}$

$x_j \geq 0, \quad \forall j \in J_p$

$x_j \leq 0, \quad \forall j \in J_n$

x_j free, $\forall j \in J_f$

variables $x \in \mathbb{R}^n$

Dual (\mathcal{D})

maximize $\lambda^T b$

$\lambda_i \geq 0, \quad \forall i \in I_{ge}$

$\lambda_i \leq 0, \quad \forall i \in I_{le}$

λ_i free, $\forall i \in I_{eq}$

$\lambda^T A_j \leq c_j, \quad \forall j \in J_p$

$\lambda^T A_j \geq c_j, \quad \forall j \in J_n$

$\lambda^T A_j = c_j, \quad \forall j \in J_f$

variables $\lambda \in \mathbb{R}^m$.

Primal-Dual Pair

Primal-Dual Pair of Problems

Primal (\mathcal{P})		Dual (\mathcal{D})	
minimize	$\underset{x}{c}^T x$	maximize	$\underset{\lambda}{\lambda}^T b$
$(\lambda_i \rightarrow)$	$a_i^T x \geq b_i, \quad \forall i \in I_{ge}$	$\lambda_i \geq 0,$	$\forall i \in I_{ge}$
$(\lambda_i \rightarrow)$	$a_i^T x \leq b_i, \quad \forall i \in I_{le}$	$\lambda_i \leq 0,$	$\forall i \in I_{le}$
$(\lambda_i \rightarrow)$	$a_i^T x = b_i, \quad \forall i \in I_{eq}$	λ_i free,	$\forall i \in I_{eq}$
	$x_j \geq 0, \quad \forall j \in J_p$	$\lambda^T A_j \leq c_j, \quad \forall j \in J_p$	
	$x_j \leq 0, \quad \forall j \in J_n$	$\lambda^T A_j \geq c_j, \quad \forall j \in J_n$	
	x_j free, $\forall j \in J_f$	$\lambda^T A_j = c_j, \quad \forall j \in J_f$	
variables	$x \in \mathbb{R}^n$	variables	$\lambda \in \mathbb{R}^m.$

Recall the procedure for deriving the dual:

- a dual decision variable λ_i for every primal constraint (except variable signs)
- constrain λ_i to ensure lower bound: $\lambda_i \text{ (?) } 0$
- for every primal decision x_j , add a dual constraint in the form $\lambda^T A_j \text{ (?) } c_j$
(involving the column A_j and the objective coefficient c_j corresponding to λ_i)

Primal-Dual Pair

Primal-Dual Pair of Problems

Primal (\mathcal{P})

minimize $\underset{x}{c^T x}$

$(\lambda_i \rightarrow) \quad a_i^T x \geq b_i, \quad \forall i \in I_{ge}$

$(\lambda_i \rightarrow) \quad a_i^T x \leq b_i, \quad \forall i \in I_{le}$

$(\lambda_i \rightarrow) \quad a_i^T x = b_i, \quad \forall i \in I_{eq}$

$x_j \geq 0, \quad \forall j \in J_p$

$x_j \leq 0, \quad \forall j \in J_n$

x_j free, $\forall j \in J_f$

variables $x \in \mathbb{R}^n$

Dual (\mathcal{D})

maximize $\underset{\lambda}{\lambda^T b}$

$\lambda_i \geq 0, \quad \forall i \in I_{ge}$

$\lambda_i \leq 0, \quad \forall i \in I_{le}$

λ_i free, $\forall i \in I_{eq}$

$\lambda^T A_j \leq c_j, \quad \forall j \in J_p$

$\lambda^T A_j \geq c_j, \quad \forall j \in J_n$

$\lambda^T A_j = c_j, \quad \forall j \in J_f$

variables $\lambda \in \mathbb{R}^m$.

Exercise

Rewrite the dual problem as a minimization problem and construct its dual.

Primal-Dual Pair

Primal-Dual Pair of Problems

Primal (\mathcal{P})

minimize $c^T x$

$$(\lambda_i \rightarrow) \quad a_i^T x \geq b_i, \quad \forall i \in I_{ge}$$

$$(\lambda_i \rightarrow) \quad a_i^T x \leq b_i, \quad \forall i \in I_{le}$$

$$(\lambda_i \rightarrow) \quad a_i^T x = b_i, \quad \forall i \in I_{eq}$$

$$x_j \geq 0, \quad \forall j \in J_p$$

$$x_j \leq 0, \quad \forall j \in J_n$$

$$x_j \text{ free}, \quad \forall j \in J_f$$

variables $x \in \mathbb{R}^n$

Dual (\mathcal{D})

maximize $\lambda^T b$

$$\lambda_i \geq 0, \quad \forall i \in I_{ge}$$

$$\lambda_i \leq 0, \quad \forall i \in I_{le}$$

$$\lambda_i \text{ free}, \quad \forall i \in I_{eq}$$

$$\lambda^T A_j \leq c_j, \quad \forall j \in J_p$$

$$\lambda^T A_j \geq c_j, \quad \forall j \in J_n$$

$$\lambda^T A_j = c_j, \quad \forall j \in J_f$$

variables $\lambda \in \mathbb{R}^m$.

Exercise

Rewrite the dual problem as a minimization problem and construct its dual.

Theorem (For LPs, the dual of the dual is the primal)

If we transform the dual of a linear optimization problem into an equivalent minimization problem and form its dual, we obtain a problem equivalent to the primal.

Rules for Constructing the Dual of Any LP

Consider any linear optimization problem (minimization/maximization):

$$\begin{array}{ll} \underset{x}{\text{minimize}} & / \quad \underset{x}{\text{maximize}} \quad c^T x \\ & (\lambda \rightarrow) \quad Ax \begin{array}{l} \leq \\ \geq \\ \leq \\ \geq \end{array} b \\ & \quad \quad \quad x \begin{array}{l} \geq \\ \leq \\ \geq \\ \leq \end{array} 0 \end{array} \quad (7)$$

Rules for Constructing the Dual of Any LP

Consider any linear optimization problem (minimization/maximization):

$$\begin{array}{ll} \underset{x}{\text{minimize}} & / \quad \underset{x}{\text{maximize}} \quad c^T x \\ & (\lambda \rightarrow) \quad Ax \begin{array}{l} \leq \\ \geq \\ = \end{array} b \\ & \quad \quad \quad x \begin{array}{l} \geq \\ \leq \\ = \end{array} 0 \end{array} \quad (7)$$

R1: A dual variable λ_i for every constraint, i.e., every row a_i^T of A .

λ_i **free** for **equality** constraints ($a_i^T x = b_i$). Otherwise: $\lambda_i \geq 0$.

Rules for Constructing the Dual of Any LP

Consider any linear optimization problem (minimization/maximization):

$$\begin{array}{ll} \underset{x}{\text{minimize}} & / \quad \underset{x}{\text{maximize}} \quad c^T x \\ & (\lambda \rightarrow) \quad Ax \begin{array}{l} \leq \\ \geq \\ = \end{array} b \\ & \quad \quad \quad x \begin{array}{l} \geq \\ \leq \\ = \end{array} 0 \end{array} \quad (7)$$

R1: A dual variable λ_i for every constraint, i.e., every row a_i^T of A .
 λ_i **free** for **equality** constraints ($a_i^T x = b_i$). Otherwise: $\lambda_i \geq 0$.

R2: In the dual, add a constraint for *every primal variable* x_j .
If x_j is **free**, write this as $\lambda^T A_j = c_j$. Otherwise: $\lambda^T A_j \leq c_j$.

Rules for Constructing the Dual of Any LP

Consider any linear optimization problem (minimization/maximization):

$$\begin{array}{ll} \underset{x}{\text{minimize}} & / \quad \underset{x}{\text{maximize}} \quad c^T x \\ & (\lambda \rightarrow) \quad Ax \begin{array}{l} \leq \\ \geq \\ = \end{array} b \\ & \quad \quad \quad x \begin{array}{l} \geq \\ \leq \\ = \end{array} 0 \end{array} \quad (7)$$

R1: A dual variable λ_i for every constraint, i.e., every row a_i^T of A .
 λ_i **free** for **equality** constraints ($a_i^T x = b_i$). Otherwise: $\lambda_i \text{ } \textcircled{?} 0$.

R2: In the dual, add a constraint for *every primal variable* x_j .
 If x_j is **free**, write this as $\lambda^T A_j = c_j$. Otherwise: $\lambda^T A_j \text{ } \textcircled{?} c_j$.

R3: To determine the signs $\textcircled{?}$, use this rule of thumb:

the dual variable λ_i is the (sub)gradient of the optimal objective value with respect to the constraint's right-hand-side b_i

Rules for Constructing the Dual of Any LP

Consider any linear optimization problem (minimization/maximization):

$$\begin{array}{ll} \underset{x}{\text{minimize}} & / \quad \underset{x}{\text{maximize}} \quad c^T x \\ & (\lambda \rightarrow) \quad Ax \begin{array}{l} \leq \\ \geq \\ = \end{array} b \\ & \quad \quad \quad x \begin{array}{l} \geq \\ \leq \\ = \end{array} 0 \end{array} \quad (7)$$

R1: A dual variable λ_i for every constraint, i.e., every row a_i^T of A .
 λ_i **free** for **equality** constraints ($a_i^T x = b_i$). Otherwise: $\lambda_i \geq 0$.

R2: In the dual, add a constraint for *every primal variable* x_j .
 If x_j is **free**, write this as $\lambda^T A_j = c_j$. Otherwise: $\lambda^T A_j \geq c_j$.

R3: To determine the signs $(?)$, use this rule of thumb:

the dual variable λ_i is the (sub)gradient of the optimal objective value with respect to the constraint's right-hand-side b_i

- in a minimization, for a " \leq " constraint, the dual variable is ≥ 0
- in a minimization, for a " \geq " constraint, the dual variable is ≤ 0
- in a maximization, for a " \leq " constraint, the dual variable is ≤ 0
- in a maximization, for a " \geq " constraint, the dual variable is ≥ 0 .

Example 1

$$\begin{aligned}(\mathcal{P}) \quad & \max 3x_1 + 2x_2 \\ & \text{s.t. } x_1 + 2x_2 \leq 4 \quad (1) \\ & \quad \quad 3x_1 + 2x_2 \geq 6 \quad (2) \\ & \quad \quad x_1 - x_2 = 1 \quad (3) \\ & \quad \quad x_1, x_2 \geq 0.\end{aligned}$$

Example 1

$$\begin{aligned}(\mathcal{P}) \quad & \max 3x_1 + 2x_2 \\ & \text{s.t. } x_1 + 2x_2 \leq 4 \quad (1) \\ & \quad \quad 3x_1 + 2x_2 \geq 6 \quad (2) \\ & \quad \quad x_1 - x_2 = 1 \quad (3) \\ & \quad \quad x_1, x_2 \geq 0.\end{aligned}$$

$$\begin{aligned}(\mathcal{D}) \quad & \min 4y_1 + 6y_2 + y_3 \\ & \text{s.t. } y_1 + 3y_2 + y_3 \geq 3, \\ & \quad \quad 2y_1 + 2y_2 - y_3 \geq 2, \\ & \quad \quad y_1 \geq 0, \quad y_2 \leq 0, \quad y_3 \text{ free.}\end{aligned}$$

Some Quick Results

Theorem ("Duals of equivalent primals")

If we transform a primal P_1 into an equivalent formulation P_2 by:

- *replacing a free variable x_i with $x_i = x_i^+ - x_i^-$,*
- *replacing an inequality with an equality by introducing a slack variable,*
- *removing linearly dependent rows a_i^T for a **feasible** LP in standard form,*

*then the duals of (P_1) and (P_2) are **equivalent**, i.e., they are either both infeasible or they have the same optimal objective.*

Weak duality

Primal (\mathcal{P})

minimize _{x} $c^T x$

$$(\lambda_i \rightarrow) \quad a_i^T x \geq b_i, \quad \forall i \in I_{\text{ge}},$$

$$(\lambda_i \rightarrow) \quad a_i^T x \leq b_i, \quad \forall i \in I_{\text{le}},$$

$$(\lambda_i \rightarrow) \quad a_i^T x = b_i, \quad \forall i \in I_{\text{eq}},$$

$$x_j \geq 0, \quad \forall j \in J_p,$$

$$x_j \leq 0, \quad \forall j \in J_n,$$

$$x_j \text{ free}, \quad \forall j \in J_f.$$

Dual (\mathcal{D})

maximize _{λ} $\lambda^T b$

$$\lambda_i \geq 0, \quad \forall i \in I_{\text{ge}},$$

$$\lambda_i \leq 0, \quad \forall i \in I_{\text{le}},$$

$$\lambda_i \text{ free}, \quad \forall i \in I_{\text{eq}},$$

$$(x_j \rightarrow) \quad \lambda^T A_j \leq c_j, \quad \forall j \in J_p,$$

$$(x_j \rightarrow) \quad \lambda^T A_j \geq c_j, \quad \forall j \in J_n,$$

$$(x_j \rightarrow) \quad \lambda^T A_j = c_j, \quad \forall j \in J_f.$$

Weak duality

Primal (\mathcal{P})			Dual (\mathcal{D})		
minimize _{x}	$c^T x$		maximize _{λ}	$\lambda^T b$	
$(\lambda_i \rightarrow)$	$a_i^T x \geq b_i,$	$\forall i \in I_{ge},$	$\lambda_i \geq 0,$		$\forall i \in I_{ge},$
$(\lambda_i \rightarrow)$	$a_i^T x \leq b_i,$	$\forall i \in I_{le},$	$\lambda_i \leq 0,$		$\forall i \in I_{le},$
$(\lambda_i \rightarrow)$	$a_i^T x = b_i,$	$\forall i \in I_{eq},$	λ_i free,		$\forall i \in I_{eq},$
	$x_j \geq 0,$	$\forall j \in J_p,$	$(x_j \rightarrow)$	$\lambda^T A_j \leq c_j,$	$\forall j \in J_p,$
	$x_j \leq 0,$	$\forall j \in J_n,$	$(x_j \rightarrow)$	$\lambda^T A_j \geq c_j,$	$\forall j \in J_n,$
	x_j free,	$\forall j \in J_f.$	$(x_j \rightarrow)$	$\lambda^T A_j = c_j,$	$\forall j \in J_f.$

Theorem (Weak duality)

If x is feasible for (\mathcal{P}) and λ is feasible for (\mathcal{D}), then $\lambda^T b \leq c^T x$.

Proof. Trivially true from our construction – omitted.

Implications of Weak Duality

Corollary

The following results hold:

(c) and (d) provide (sub)optimality certificates, but...

How do we know that the gaps in (c) are not very large?

How do we know that x and λ satisfying (d) even exist?

Implications of Weak Duality

Corollary

The following results hold:

(a) If the optimal objective in (\mathcal{P}) is $-\infty$, then (\mathcal{D}) must be infeasible.

(c) and (d) provide (sub)optimality certificates, but...

How do we know that the gaps in (c) are not very large?

How do we know that x and λ satisfying (d) even exist?

Implications of Weak Duality

Corollary

The following results hold:

- (a) If the optimal objective in (\mathcal{P}) is $-\infty$, then (\mathcal{D}) must be infeasible.*
- (b) If the optimal objective in (\mathcal{D}) is $+\infty$, then (\mathcal{P}) must be infeasible.*

(c) and (d) provide (sub)optimality certificates, but...

How do we know that the gaps in (c) are not very large?

How do we know that x and λ satisfying (d) even exist?

Implications of Weak Duality

Corollary

The following results hold:

(a) If the optimal objective in (\mathcal{P}) is $-\infty$, then (\mathcal{D}) must be infeasible.

(b) If the optimal objective in (\mathcal{D}) is $+\infty$, then (\mathcal{P}) must be infeasible.

(c) If $x \in P$ and $\lambda \in D$, then:

$$c^T x - p^* \leq c^T x - \lambda^T b \quad \textbf{and} \quad d^* - \lambda^T b \leq c^T x - \lambda^T b.$$

(c) and (d) provide (sub)optimality certificates, but...

How do we know that the gaps in (c) are not very large?

How do we know that x and λ satisfying (d) even exist?

Implications of Weak Duality

Corollary

The following results hold:

(a) *If the optimal objective in (\mathcal{P}) is $-\infty$, then (\mathcal{D}) must be infeasible.*

(b) *If the optimal objective in (\mathcal{D}) is $+\infty$, then (\mathcal{P}) must be infeasible.*

(c) *If $x \in P$ and $\lambda \in D$, then:*

$$c^T x - p^* \leq c^T x - \lambda^T b \quad \text{and} \quad d^* - \lambda^T b \leq c^T x - \lambda^T b.$$

(d) *If $x \in P$, $\lambda \in D$, and $\lambda^T b = c^T x$, then x **optimal** for (\mathcal{P}) and λ **optimal** for (\mathcal{D}) .*

(c) and (d) provide (sub)optimality certificates, but...

Implications of Weak Duality

Corollary

The following results hold:

(a) *If the optimal objective in (\mathcal{P}) is $-\infty$, then (\mathcal{D}) must be infeasible.*

(b) *If the optimal objective in (\mathcal{D}) is $+\infty$, then (\mathcal{P}) must be infeasible.*

(c) *If $x \in P$ and $\lambda \in D$, then:*

$$c^T x - p^* \leq c^T x - \lambda^T b \quad \text{and} \quad d^* - \lambda^T b \leq c^T x - \lambda^T b.$$

(d) *If $x \in P$, $\lambda \in D$, and $\lambda^T b = c^T x$, then x **optimal** for (\mathcal{P}) and λ **optimal** for (\mathcal{D}) .*

(c) and (d) provide (sub)optimality certificates, but...

How do we know that the gaps in (c) are not very large?

How do we know that x and λ satisfying (d) even exist?