

CME 307 / MS&E 311: Optimization

Optimality conditions and convexity

Professor Udell

Management Science and Engineering
Stanford

January 29, 2024

Solution of an optimization problem

$$\text{minimize } f(x)$$

for $f : \mathcal{D} \rightarrow \mathbf{R}$. x^* is a

- ▶ **global minimizer** if $f(x) \geq f(x^*)$ for all $x \in \mathcal{D}$.
- ▶ **local minimizer** if there is a neighborhood \mathcal{N} around x^* so that $f(x) \geq f(x^*)$ for all $x \in \mathcal{N}$.
- ▶ **isolated local minimizer** if the neighborhood \mathcal{N} contains no other local minimizers.
- ▶ **unique minimizer** if it is the only global minimizer.

Solution of an optimization problem

$$\text{minimize } f(x)$$

for $f : \mathcal{D} \rightarrow \mathbf{R}$. x^* is a

- ▶ **global minimizer** if $f(x) \geq f(x^*)$ for all $x \in \mathcal{D}$.
- ▶ **local minimizer** if there is a neighborhood \mathcal{N} around x^* so that $f(x) \geq f(x^*)$ for all $x \in \mathcal{N}$.
- ▶ **isolated local minimizer** if the neighborhood \mathcal{N} contains no other local minimizers.
- ▶ **unique minimizer** if it is the only global minimizer.

pictures!

First order optimality condition

Theorem

If $x^ \in \mathbf{R}^n$ is a local minimizer of a differentiable function $f : \mathbf{R}^n \rightarrow \mathbf{R}$, then $\nabla f(x^*) = 0$.*

First order optimality condition

Theorem

If $x^ \in \mathbf{R}^n$ is a local minimizer of a differentiable function $f : \mathbf{R}^n \rightarrow \mathbf{R}$, then $\nabla f(x^*) = 0$.*

proof: suppose by contradiction that $\nabla f(x^*) \neq 0$. consider points of the form $x_\alpha = x^* - \alpha \nabla f(x^*)$ for $\alpha > 0$. by definition of the gradient,

$$\lim_{\alpha \rightarrow 0} \frac{f(x_\alpha) - f(x^*)}{\alpha} = -\nabla f(x^*)^\top \nabla f(x^*) = -\|\nabla f(x^*)\|^2 < 0$$

so for any sufficiently small $\alpha > 0$, we have $f(x_\alpha) < f(x^*)$, which contradicts the fact that x^* is a local minimizer.

Second order optimality condition

Theorem

If $x^ \in \mathbf{R}^n$ is a local minimizer of a twice differentiable function $f : \mathbf{R}^n \rightarrow \mathbf{R}$, then $\nabla^2 f(x^*) \succeq 0$.*

Second order optimality condition

Theorem

If $x^ \in \mathbf{R}^n$ is a local minimizer of a twice differentiable function $f : \mathbf{R}^n \rightarrow \mathbf{R}$, then $\nabla^2 f(x^*) \succeq 0$.*

proof: similar to the previous proof. use the fact that the second order approximation

$$f(x_\alpha) \approx f(x^*) + \nabla f(x^*)^\top (x_\alpha - x^*) + \frac{1}{2} (x_\alpha - x^*)^\top \nabla^2 f(x^*) (x_\alpha - x^*)$$

is accurate locally to show a contradiction unless $\nabla^2 f(x^*) \succeq 0$: if not, there is a direction v such that $v^\top \nabla^2 f(x^*) v < 0$. then $f(x + \alpha v) < f(x^*)$ for α arbitrarily small, which contradicts the fact that x^* is a local minimizer.

Outline

Convex sets

Definition

A set $S \subseteq \mathbf{R}^n$ is convex if it contains every chord: for all $\theta \in [0, 1]$, $w, v \in S$,

$$\theta w + (1 - \theta)v \in S$$

Convex sets

Definition

A set $S \subseteq \mathbf{R}^n$ is convex if it contains every chord: for all $\theta \in [0, 1]$, $w, v \in S$,

$$\theta w + (1 - \theta)v \in S$$

Q: Which of these are convex?

ellipsoid, half moon

Convex functions

a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex iff

Convex functions

a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex iff

- **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

Convex functions

a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex iff

- ▶ **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.** $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$ is convex

Convex functions

a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex iff

- ▶ **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.** $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$ is convex
- ▶ **First order condition.** if f is differentiable,

$$f(v) - f(w) \geq \nabla f(w)^\top (v - w) \quad \forall w, v \in \mathbf{R}^n$$

Convex functions

a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex iff

- ▶ **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.** $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$ is convex
- ▶ **First order condition.** if f is differentiable,

$$f(v) - f(w) \geq \nabla f(w)^\top (v - w) \quad \forall w, v \in \mathbf{R}^n$$

- ▶ **Second order condition.** If f is twice differentiable, its Hessian is always psd:

$$\lambda_{\min}(\nabla^2 f(x)) \geq 0 \quad \text{for all } x \in \mathbf{R}^n$$

Convex functions

a function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is convex iff

- ▶ **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.** $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$ is convex
- ▶ **First order condition.** if f is differentiable,

$$f(v) - f(w) \geq \nabla f(w)^\top (v - w) \quad \forall w, v \in \mathbf{R}^n$$

- ▶ **Second order condition.** If f is twice differentiable, its Hessian is always psd:

$$\lambda_{\min}(\nabla^2 f(x)) \geq 0 \quad \text{for all } x \in \mathbf{R}^n$$

Q: Which of these are convex?

quadratic, abs, pwl, step, jump, logistic, logistic loss

Convex optimization

an optimization problem is convex if:

- ▶ **Geometrically:** the feasible set and the epigraph of the objective are convex
- ▶ **NLP:** the objective and inequality constraints are convex functions, and the equality constraints are affine

Convex optimization

an optimization problem is convex if:

- ▶ **Geometrically:** the feasible set and the epigraph of the objective are convex
- ▶ **NLP:** the objective and inequality constraints are convex functions, and the equality constraints are affine

why convex optimization?

- ▶ relatively complete theory
- ▶ efficient solvers
- ▶ conceptual tools that generalize

duality, stopping conditions, ...

Convex optimization

an optimization problem is convex if:

- ▶ **Geometrically:** the feasible set and the epigraph of the objective are convex
- ▶ **NLP:** the objective and inequality constraints are convex functions, and the equality constraints are affine

why convex optimization?

- ▶ relatively complete theory
- ▶ efficient solvers
- ▶ conceptual tools that generalize

duality, stopping conditions, ...

- ▶ a function f is concave if $-f$ is convex
- ▶ concave maximization results in a **convex** optimization problem

Local minima are global for convex functions

Theorem

If x^ is a local minimizer of a convex function f , then x^* is a global minimizer.*

Local minima are global for convex functions

Theorem

If x^ is a local minimizer of a convex function f , then x^* is a global minimizer.*

proof: suppose by contradiction that another point x' is a global minimizer, with $f(x') < f(x^*)$. draw the chord between x' and x^* . since the chord lies above f , every convex combination $x = \theta x^* + (1 - \theta)x'$ of x' and x^* for $\theta \in (0, 1)$ has a value $f(x) < f(x^*)$. this is true even for $x \rightarrow x^*$, contradicting our assumption that x^* is a local minimizer.

Corollary

Corollary

If f is convex and differentiable and $\nabla f(x^) = 0$, then x^* is a global minimizer.*

Corollary

Corollary

If f is convex and differentiable and $\nabla f(x^) = 0$, then x^* is a global minimizer.*

Q: Is a global minimizer of a convex function always unique?

Corollary

Corollary

If f is convex and differentiable and $\nabla f(x^) = 0$, then x^* is a global minimizer.*

Q: Is a global minimizer of a convex function always unique?

A: No. Picture.

First-order condition

Definition

$x^* \in \mathbf{R}^n$ is a **stationary point** of a differentiable function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ if $\nabla f(x^*) = 0$.

First-order condition

Definition

$x^* \in \mathbf{R}^n$ is a **stationary point** of a differentiable function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ if $\nabla f(x^*) = 0$.

Q: Can a global minimum have a non-zero gradient?

First-order condition

Definition

$x^* \in \mathbf{R}^n$ is a **stationary point** of a differentiable function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ if $\nabla f(x^*) = 0$.

Q: Can a global minimum have a non-zero gradient?

A: No.

First-order condition

Definition

$x^* \in \mathbf{R}^n$ is a **stationary point** of a differentiable function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ if $\nabla f(x^*) = 0$.

Q: Can a global minimum have a non-zero gradient?

A: No.

Q: Is a stationary point always a global minimum?

First-order condition

Definition

$x^* \in \mathbf{R}^n$ is a **stationary point** of a differentiable function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ if $\nabla f(x^*) = 0$.

Q: Can a global minimum have a non-zero gradient?

A: No.

Q: Is a stationary point always a global minimum?

A: No.

First-order condition

Definition

$x^* \in \mathbf{R}^n$ is a **stationary point** of a differentiable function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ if $\nabla f(x^*) = 0$.

Q: Can a global minimum have a non-zero gradient?

A: No.

Q: Is a stationary point always a global minimum?

A: No.

Q: ... for convex functions?

First-order condition

Definition

$x^* \in \mathbf{R}^n$ is a **stationary point** of a differentiable function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ if $\nabla f(x^*) = 0$.

Q: Can a global minimum have a non-zero gradient?

A: No.

Q: Is a stationary point always a global minimum?

A: No.

Q: ... for convex functions?

A: Yes.

First-order condition

Definition

$x^* \in \mathbf{R}^n$ is a **stationary point** of a differentiable function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ if $\nabla f(x^*) = 0$.

Q: Can a global minimum have a non-zero gradient?

A: No.

Q: Is a stationary point always a global minimum?

A: No.

Q: ... for convex functions?

A: Yes.

$\nabla f(x^*) = 0$ is the **first-order (necessary) condition** for optimality.

Invex function

Definition

A function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is **invex** if for some vector-valued function $\eta : \mathbf{R}^n \times \mathbf{R}^n \rightarrow \mathbf{R}^n$,

$$f(x) - f(u) \geq \eta(x, u)^\top \nabla f(u) \quad \forall u \in \mathbf{R}^n, x \in \text{dom } f$$

Theorem (Craven and Glover, Ben-Israel and Mond)

A function is invex iff every stationary point is a global minimum.