

# CME 307 / MS&E 311: Optimization

## Optimality conditions and convexity

Professor Udell

Management Science and Engineering  
Stanford

February 5, 2024

## Questions from last class

- ▶ clarify proof of strong duality
- ▶ how many iterations of branch and bound?
- ▶ how to use duality to solve a problem? when to stop?
- ▶ duality for problems with inequality constraints?

# Outline

Constrained and unconstrained optimization

Optimality conditions

Convex optimization

## Constrained vs unconstrained optimization

constrained optimization

- ▶ examples: scheduling, routing, packing, logistics, scheduling, control
- ▶ what's hard: finding a feasible point

unconstrained optimization

- ▶ examples: data fitting, statistical/machine learning
- ▶ what's hard: reducing the objective

both are necessary for real-world problems!

## Unconstrained smooth optimization

for  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  ctly differentiable,

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{variable} & x \in \mathbf{R}^n \end{array}$$

how to solve?

## Unconstrained smooth optimization

for  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  ctly differentiable,

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{variable} & x \in \mathbf{R}^n \end{array}$$

how to solve? approximate as a quadratic problem

$$f(x) \approx f(x_0) + \nabla f(x_0)^T (x - x_0) + \frac{1}{2}(x - x_0)^T H(x_0)(x - x_0)$$

and find solution  $x_{\text{quad}}$  to the quadratic problem.

then set  $x_0 \leftarrow x_{\text{quad}}$  and repeat.

## Nonlinear optimization

optimization problem: nonlinear form

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq b_i, \quad i = 1, \dots, m_1 \\ & h(x) = 0 \\ \text{variable} & x \in \mathbf{R}^n\end{array}$$

- ▶  $x = (x_1, \dots, x_n)$ : optimization variables
- ▶  $f_0 : \mathbf{R}^n \rightarrow \mathbf{R}$ : objective function
- ▶  $f_i : \mathbf{R}^n \rightarrow \mathbf{R}, i = 1, \dots, m$ : constraint functions

special case: **unconstrained optimization**

## Example: process control

You are the process engineer for a desalination plant that produces drinking water. The plant has a variety of knobs, collected in vector  $x$ , that you can turn to control the process. These control, e.g., how much water is pumped into the plant, how much pressure is used to force the water through filters, and how much of each chemical is added to the water.

- ▶  $f_0(x)$ : cost of water produced
- ▶  $f_i(x)$ : level of each measured impurity in the water
- ▶  $b_i$ : maximum allowable level of each impurity

Given a setting of the knobs, you can observe the cost of water produced and the levels of impurities.

**What is the optimal setting of the knobs?**



## Oracles

an optimization **oracle** is your interface for accessing the problem data:  
e.g., an oracle for  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  can evaluate for any  $x \in \mathbf{R}^n$ :

- ▶ **zero-order:**  $f_0(x)$
- ▶ **first-order:**  $f_0(x)$  and  $\nabla f_0(x)$
- ▶ **second-order:**  $f_0(x)$ ,  $\nabla f_0(x)$ , and  $\nabla^2 f_0(x)$

why oracles?

- ▶ can optimize real systems based on observed output (not just models)
- ▶ can use and extend old or complex but trusted code (e.g., NASA, PDE simulations, ...)
- ▶ can prove lower bounds on the oracle complexity of a problem class

source: Nesterov 2004 “Introductory Lectures on Convex Optimization”

## Nonlinear optimization: how to solve?

depends on the oracle:

- ▶ first- or second-order: approximate by a sequence of quadratic problems
- ▶ zero-order: harder, lots of methods
  - ▶ simulated annealing
  - ▶ Bayesian optimization
  - ▶ pseudo-higher-order methods, e.g., compute approximate gradient

# Outline

Constrained and unconstrained optimization

Optimality conditions

Convex optimization

## Solution of an optimization problem

$$\text{minimize } f(x)$$

for  $f : \mathcal{D} \rightarrow \mathbf{R}$ .  $x^*$  is a

- ▶ **global minimizer** if  $f(x) \geq f(x^*)$  for all  $x \in \mathcal{D}$ .
- ▶ **local minimizer** if there is a neighborhood  $\mathcal{N}$  around  $x^*$  so that  $f(x) \geq f(x^*)$  for all  $x \in \mathcal{N}$ .
- ▶ **isolated local minimizer** if the neighborhood  $\mathcal{N}$  contains no other local minimizers.
- ▶ **unique minimizer** if it is the only global minimizer.

## Solution of an optimization problem

$$\text{minimize } f(x)$$

for  $f : \mathcal{D} \rightarrow \mathbf{R}$ .  $x^*$  is a

- ▶ **global minimizer** if  $f(x) \geq f(x^*)$  for all  $x \in \mathcal{D}$ .
- ▶ **local minimizer** if there is a neighborhood  $\mathcal{N}$  around  $x^*$  so that  $f(x) \geq f(x^*)$  for all  $x \in \mathcal{N}$ .
- ▶ **isolated local minimizer** if the neighborhood  $\mathcal{N}$  contains no other local minimizers.
- ▶ **unique minimizer** if it is the only global minimizer.

pictures!

## First order optimality condition

### Theorem

*If  $x^* \in \mathbf{R}^n$  is a local minimizer of a differentiable function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$ , then  $\nabla f(x^*) = 0$ .*

## First order optimality condition

### Theorem

If  $x^* \in \mathbf{R}^n$  is a local minimizer of a differentiable function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$ , then  $\nabla f(x^*) = 0$ .

**proof:** suppose by contradiction that  $\nabla f(x^*) \neq 0$ . consider points of the form  $x_\alpha = x^* - \alpha \nabla f(x^*)$  for  $\alpha > 0$ . by definition of the gradient,

$$\lim_{\alpha \rightarrow 0} \frac{f(x_\alpha) - f(x^*)}{\alpha} = -\nabla f(x^*)^\top \nabla f(x^*) = -\|\nabla f(x^*)\|^2 < 0$$

so for any sufficiently small  $\alpha > 0$ , we have  $f(x_\alpha) < f(x^*)$ , which contradicts the fact that  $x^*$  is a local minimizer.

## Second order optimality condition

### Theorem

*If  $x^* \in \mathbf{R}^n$  is a local minimizer of a twice differentiable function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$ , then  $\nabla^2 f(x^*) \succeq 0$ .*



## Second order optimality condition

### Theorem

If  $x^* \in \mathbf{R}^n$  is a local minimizer of a twice differentiable function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$ , then  $\nabla^2 f(x^*) \succeq 0$ .

**proof:** similar to the previous proof. use the fact that the second order approximation

$$f(x_\alpha) \approx f(x^*) + \nabla f(x^*)^\top (x_\alpha - x^*) + \frac{1}{2}(x_\alpha - x^*)^\top \nabla^2 f(x^*)(x_\alpha - x^*)$$

is accurate locally to show a contradiction unless  $\nabla^2 f(x^*) \succeq 0$ : if not, there is a direction  $v$  such that  $v^\top \nabla^2 f(x^*) v < 0$ . then  $f(x + \alpha v) < f(x^*)$  for  $\alpha$  arbitrarily small, which contradicts the fact that  $x^*$  is a local minimizer.

# Outline

Constrained and unconstrained optimization

Optimality conditions

Convex optimization

## Convex sets

### Definition

A set  $S \subseteq \mathbf{R}^n$  is convex if it contains every chord: for all  $\theta \in [0, 1]$ ,  $w, v \in S$ ,

$$\theta w + (1 - \theta)v \in S$$

## Convex sets

### Definition

A set  $S \subseteq \mathbf{R}^n$  is convex if it contains every chord: for all  $\theta \in [0, 1]$ ,  $w, v \in S$ ,

$$\theta w + (1 - \theta)v \in S$$

**Q:** Which of these are convex?

ellipsoid, half moon

## Convex functions

a function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is convex iff

## Convex functions

a function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is convex iff

► **Chords.** it never lies above its chord:  $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

## Convex functions

a function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is convex iff

- ▶ **Chords.** it never lies above its chord:  $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.**  $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$  is convex

## Convex functions

a function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is convex iff

- ▶ **Chords.** it never lies above its chord:  $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.**  $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$  is convex
- ▶ **First order condition.** if  $f$  is differentiable,

$$f(v) \geq f(w) + \nabla f(w)^\top (v - w), \quad \forall w, v \in \mathbf{R}^n$$



## Convex functions

a function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is convex iff

- ▶ **Chords.** it never lies above its chord:  $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.**  $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$  is convex
- ▶ **First order condition.** if  $f$  is differentiable,

$$f(v) \geq f(w) + \nabla f(w)^\top (v - w), \quad \forall w, v \in \mathbf{R}^n$$

- ▶ **Second order condition.** If  $f$  is twice differentiable, its Hessian is always psd:

$$\lambda_{\min}(\nabla^2 f(x)) \geq 0, \quad \forall x \in \mathbf{R}^n$$

## Convex functions

a function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is convex iff

- ▶ **Chords.** it never lies above its chord:  $\forall \theta \in [0, 1], w, v \in \mathbf{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.**  $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$  is convex
- ▶ **First order condition.** if  $f$  is differentiable,

$$f(v) \geq f(w) + \nabla f(w)^\top (v - w), \quad \forall w, v \in \mathbf{R}^n$$

- ▶ **Second order condition.** If  $f$  is twice differentiable, its Hessian is always psd:

$$\lambda_{\min}(\nabla^2 f(x)) \geq 0, \quad \forall x \in \mathbf{R}^n$$

**Q:** Which of these are convex?

quadratic, abs, pwl, step, jump, logistic, logistic loss

## Convex optimization

an optimization problem is convex if:

- ▶ **Geometrically:** the feasible set and the epigraph of the objective are convex
- ▶ **NLP:** the objective and inequality constraints are convex functions, and the equality constraints are affine

## Convex optimization

an optimization problem is convex if:

- ▶ **Geometrically:** the feasible set and the epigraph of the objective are convex
- ▶ **NLP:** the objective and inequality constraints are convex functions, and the equality constraints are affine

why convex optimization?

- ▶ relatively complete theory
- ▶ efficient solvers
- ▶ conceptual tools that generalize

duality, stopping conditions, ...

## Convex optimization

an optimization problem is convex if:

- ▶ **Geometrically:** the feasible set and the epigraph of the objective are convex
- ▶ **NLP:** the objective and inequality constraints are convex functions, and the equality constraints are affine

why convex optimization?

- ▶ relatively complete theory
- ▶ efficient solvers
- ▶ conceptual tools that generalize

duality, stopping conditions, ...

- ▶ a function  $f$  is concave if  $-f$  is convex
- ▶ concave maximization  $\implies$  a convex optimization problem

## Local minima are global for convex functions

### Theorem

*If  $x^*$  is a local minimizer of a convex function  $f$ , then  $x^*$  is a global minimizer.*

## Local minima are global for convex functions

### Theorem

*If  $x^*$  is a local minimizer of a convex function  $f$ , then  $x^*$  is a global minimizer.*

**proof?**

## Local minima are global for convex functions

### Theorem

*If  $x^*$  is a local minimizer of a convex function  $f$ , then  $x^*$  is a global minimizer.*

**proof?** suppose by contradiction that another point  $x'$  is a global minimizer, with  $f(x') < f(x^*)$ . draw the chord between  $x'$  and  $x^*$ . since the chord lies above  $f$ , every convex combination  $x = \theta x^* + (1 - \theta)x'$  of  $x'$  and  $x^*$  for  $\theta \in (0, 1)$  has a value  $f(x) < f(x^*)$ . this is true even for  $x \rightarrow x^*$ , contradicting our assumption that  $x^*$  is a local minimizer.



## Corollary

### Corollary

*If  $f$  is convex and differentiable and  $\nabla f(x^*) = 0$ , then  $x^*$  is a global minimizer.*

## Corollary

### Corollary

*If  $f$  is convex and differentiable and  $\nabla f(x^*) = 0$ , then  $x^*$  is a global minimizer.*

**Q:** Is a global minimizer of a convex function always unique?

## Corollary

### Corollary

*If  $f$  is convex and differentiable and  $\nabla f(x^*) = 0$ , then  $x^*$  is a global minimizer.*

**Q:** Is a global minimizer of a convex function always unique?

**A:** No. Picture.

## First-order condition

### Definition

$x^* \in \mathbf{R}^n$  is a **stationary point** of a differentiable function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  if  $\nabla f(x^*) = 0$ .

## First-order condition

### Definition

$x^* \in \mathbf{R}^n$  is a **stationary point** of a differentiable function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  if  $\nabla f(x^*) = 0$ .

**Q:** Can a global minimum have a non-zero gradient?

## First-order condition

### Definition

$x^* \in \mathbf{R}^n$  is a **stationary point** of a differentiable function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  if  $\nabla f(x^*) = 0$ .

**Q:** Can a global minimum have a non-zero gradient?

**A:** No.

## First-order condition

### Definition

$x^* \in \mathbf{R}^n$  is a **stationary point** of a differentiable function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  if  $\nabla f(x^*) = 0$ .

**Q:** Can a global minimum have a non-zero gradient?

**A:** No.

**Q:** Is a stationary point always a global minimum?

## First-order condition

### Definition

$x^* \in \mathbf{R}^n$  is a **stationary point** of a differentiable function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  if  $\nabla f(x^*) = 0$ .

**Q:** Can a global minimum have a non-zero gradient?

**A:** No.

**Q:** Is a stationary point always a global minimum?

**A:** No.



## First-order condition

### Definition

$x^* \in \mathbf{R}^n$  is a **stationary point** of a differentiable function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  if  $\nabla f(x^*) = 0$ .

**Q:** Can a global minimum have a non-zero gradient?

**A:** No.

**Q:** Is a stationary point always a global minimum?

**A:** No.

**Q:** ... for convex functions?

## First-order condition

### Definition

$x^* \in \mathbf{R}^n$  is a **stationary point** of a differentiable function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  if  $\nabla f(x^*) = 0$ .

**Q:** Can a global minimum have a non-zero gradient?

**A:** No.

**Q:** Is a stationary point always a global minimum?

**A:** No.

**Q:** ... for convex functions?

**A:** Yes.

## First-order condition

### Definition

$x^* \in \mathbf{R}^n$  is a **stationary point** of a differentiable function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  if  $\nabla f(x^*) = 0$ .

**Q:** Can a global minimum have a non-zero gradient?

**A:** No.

**Q:** Is a stationary point always a global minimum?

**A:** No.

**Q:** ... for convex functions?

**A:** Yes.

$\nabla f(x^*) = 0$  is the **first-order (necessary) condition** for optimality.

## Invex function

### Definition

A function  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is **invex** if for some vector-valued function  $\eta : \mathbf{R}^n \times \mathbf{R}^n \rightarrow \mathbf{R}^n$ ,

$$f(x) - f(u) \geq \eta(x, u)^\top \nabla f(u) \quad \forall u \in \mathbf{R}^n, x \in \text{dom } f$$

### Theorem (Craven and Glover, Ben-Israel and Mond)

*A function is invex iff every stationary point is a global minimum.*