

CME 307 / MS&E 311 / OIT 676: Optimization

LP geometry, modeling and solution techniques

Professor Udell

Management Science and Engineering
Stanford

September 29, 2025

Course survey

you're interested in:

- ▶ modeling real-world problems, from finance and economics to energy systems and trajectory planning
- ▶ robustness and modeling under uncertainty
- ▶ understanding core optimization concepts like duality
- ▶ ...

questions:

- ▶ what readings are required?
- ▶ what projects are allowed?
- ▶ Friday section?
- ▶ programming requirements?

Quiz Wednesday

- ▶ Take paper when you arrive
- ▶ Write your SUNetID¹ and your name.
- ▶ We will project the question.
- ▶ You will have 5 minutes to write your answer on the paper.
- ▶ When time is up, put your pen or pencil down and stop writing.

¹Your SUNetID is your Stanford University Network ID, and consists of letters, **not** numbers. Your email is yourSUNetID@stanford.edu.

Quiz Wednesday

- ▶ Take paper when you arrive
- ▶ Write your SUNetID¹ and your name.
- ▶ We will project the question.
- ▶ You will have 5 minutes to write your answer on the paper.
- ▶ When time is up, put your pen or pencil down and stop writing.

since we are calibrating what is possible with a 5 minute quiz, we will grade on a **check** basis.

- ▶ **no check**: no serious attempt
- ▶ **check**: a serious attempt
- ▶ **check+**: correct or nearly correct

¹Your SUNetID is your Stanford University Network ID, and consists of letters, **not** numbers. Your email is yourSUNetID@stanford.edu.

Exit ticket

Please fill out the exit ticket (linked from course schedule spreadsheet) before you leave:

- ▶ Helps us improve the course

We will not grade this, but we will read it!

Course overview

the practice of optimization consists of:

- ▶ modeling (real-world problems \rightarrow optimization problems)
- ▶ analysis (properties of optimization problems)
- ▶ algorithms (how to solve optimization problems)

all three are necessary for success in optimization!

Course overview

the practice of optimization consists of:

- ▶ modeling (real-world problems \rightarrow optimization problems)
- ▶ analysis (properties of optimization problems)
- ▶ algorithms (how to solve optimization problems)

all three are necessary for success in optimization!

- ▶ last Monday: mostly modeling
- ▶ today: mostly analysis
- ▶ next Monday: algorithms and some more modeling

Outline

LP standard form

LP inequality form

What kinds of points can be optimal?

Solving LPs

Modeling

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $\pm\infty$

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $\pm\infty$

Q: if $p^* = -\infty$, does a solution exist?

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $\pm\infty$

Q: if $p^* = -\infty$, does a solution exist?

Q: if $p^* = \infty$, does a solution exist?

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $\pm\infty$

Q: if $p^* = -\infty$, does a solution exist?

Q: if $p^* = \infty$, does a solution exist?

henceforth assume $A \in \mathbb{R}^{m \times n}$ has full row rank m

Q: why? how to check?

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $\pm\infty$

Q: if $p^* = -\infty$, does a solution exist?

Q: if $p^* = \infty$, does a solution exist?

henceforth assume $A \in \mathbb{R}^{m \times n}$ has full row rank m

Q: why? how to check?

A: otherwise infeasible or redundant rows; use gaussian elimination to check and remove

LP example: diet problem

We are planning a backpacking trip, and want to minimize the total weight of the food packed subject to nutritional requirements. We have a list of essential nutrients and how much an active person needs of each. We also know the weight of each food, and how much of each nutrient is in each food.

- ▶ x_j servings of food j , $j = 1, \dots, n$
- ▶ c_j weight per serving
- ▶ a_{ij} amount of nutrient i in food j
- ▶ b_i required amount of nutrient i , $i = 1, \dots, m$

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

Geometry of LP

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

Geometry of LP

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

interpretation: conic hull

- ▶ define the **cone** generated by $A = [A_1, \dots, A_n]$:

$$\{Ax \mid x \geq 0\} = \left\{ \sum_{i=1}^n A_i x_i \mid x \geq 0 \right\} = \mathbf{cone}(A_1, \dots, A_n)$$

Geometry of LP

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

interpretation: conic hull

- ▶ define the **cone** generated by $A = [A_1, \dots, A_n]$:

$$\{Ax \mid x \geq 0\} = \left\{ \sum_{i=1}^n A_i x_i \mid x \geq 0 \right\} = \mathbf{cone}(A_1, \dots, A_n)$$

- ▶ LP is feasible if $b \in \mathbf{cone}(A_1, \dots, A_n)$

Geometry of LP

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

interpretation: conic hull

- ▶ define the **cone** generated by $A = [A_1, \dots, A_n]$:

$$\{Ax \mid x \geq 0\} = \left\{ \sum_{i=1}^n A_i x_i \mid x \geq 0 \right\} = \mathbf{cone}(A_1, \dots, A_n)$$

- ▶ LP is feasible if $b \in \mathbf{cone}(A_1, \dots, A_n)$

interpretation: intersection of hyperplane and halfspaces

- ▶ define a **hyperplane** $\{x \mid Ax = b\}$

Geometry of LP

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

interpretation: conic hull

- ▶ define the **cone** generated by $A = [A_1, \dots, A_n]$:

$$\{Ax \mid x \geq 0\} = \left\{ \sum_{i=1}^n A_i x_i \mid x \geq 0 \right\} = \mathbf{cone}(A_1, \dots, A_n)$$

- ▶ LP is feasible if $b \in \mathbf{cone}(A_1, \dots, A_n)$

interpretation: intersection of hyperplane and halfspaces

- ▶ define a **hyperplane** $\{x \mid Ax = b\}$ (dimension?)

Geometry of LP

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

interpretation: conic hull

- ▶ define the **cone** generated by $A = [A_1, \dots, A_n]$:

$$\{Ax \mid x \geq 0\} = \left\{ \sum_{i=1}^n A_i x_i \mid x \geq 0 \right\} = \mathbf{cone}(A_1, \dots, A_n)$$

- ▶ LP is feasible if $b \in \mathbf{cone}(A_1, \dots, A_n)$

interpretation: intersection of hyperplane and halfspaces

- ▶ define a **hyperplane** $\{x \mid Ax = b\}$ (dimension?)
- ▶ define a **halfspace** $\{x \mid a^T x \geq b\}$

Geometry of LP

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

interpretation: conic hull

- ▶ define the **cone** generated by $A = [A_1, \dots, A_n]$:

$$\{Ax \mid x \geq 0\} = \left\{ \sum_{i=1}^n A_i x_i \mid x \geq 0 \right\} = \mathbf{cone}(A_1, \dots, A_n)$$

- ▶ LP is feasible if $b \in \mathbf{cone}(A_1, \dots, A_n)$

interpretation: intersection of hyperplane and halfspaces

- ▶ define a **hyperplane** $\{x \mid Ax = b\}$ (dimension?)
- ▶ define a **halfspace** $\{x \mid a^T x \geq b\}$
- ▶ the **positive orthant** $x \geq 0$ is an intersection of halfspaces

Geometry of LP

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

interpretation: conic hull

- ▶ define the **cone** generated by $A = [A_1, \dots, A_n]$:

$$\{Ax \mid x \geq 0\} = \left\{ \sum_{i=1}^n A_i x_i \mid x \geq 0 \right\} = \mathbf{cone}(A_1, \dots, A_n)$$

- ▶ LP is feasible if $b \in \mathbf{cone}(A_1, \dots, A_n)$

interpretation: intersection of hyperplane and halfspaces

- ▶ define a **hyperplane** $\{x \mid Ax = b\}$ (dimension?)
- ▶ define a **halfspace** $\{x \mid a^T x \geq b\}$
- ▶ the **positive orthant** $x \geq 0$ is an intersection of halfspaces
- ▶ LP is feasible if hyperplane $\{x \mid Ax = b\}$ intersects the positive orthant

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ define the **convex hull** of a set S :

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1 \right\}$$

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ define the **convex hull** of a set S :

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1 \right\}$$

- ▶ define **polytope**: the convex hull of a finite set: $\text{conv}(\{x_1, \dots, x_k\})$

some useful convex sets:

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ define the **convex hull** of a set S :

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1 \right\}$$

- ▶ define **polytope**: the convex hull of a finite set: $\text{conv}(\{x_1, \dots, x_k\})$

some useful convex sets:

- ▶ a hyperplane is convex

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ define the **convex hull** of a set S :

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1 \right\}$$

- ▶ define **polytope**: the convex hull of a finite set: $\text{conv}(\{x_1, \dots, x_k\})$

some useful convex sets:

- ▶ a hyperplane is convex
- ▶ a halfspace is convex

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ define the **convex hull** of a set S :

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1 \right\}$$

- ▶ define **polytope**: the convex hull of a finite set: $\text{conv}(\{x_1, \dots, x_k\})$

some useful convex sets:

- ▶ a hyperplane is convex
- ▶ a halfspace is convex
- ▶ the intersection of convex sets is convex

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ define the **convex hull** of a set S :

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1 \right\}$$

- ▶ define **polytope**: the convex hull of a finite set: $\text{conv}(\{x_1, \dots, x_k\})$

some useful convex sets:

- ▶ a hyperplane is convex
- ▶ a halfspace is convex
- ▶ the intersection of convex sets is convex
- ▶ the feasible set $\{x : Ax = b, x \geq 0\}$ is convex

Outline

LP standard form

LP inequality form

What kinds of points can be optimal?

Solving LPs

Modeling

LP inequality form

another useful form for LP is **inequality form**

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

LP inequality form

another useful form for LP is **inequality form**

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

interpretation: halfspaces

- ▶ $a_i^T x \leq b_i$ defines a **halfspace**
- ▶ $Ax \leq b$ defines a **polyhedron**: intersection of halfspaces
- ▶ LP is feasible if polyhedron $\{x \mid Ax \leq b\}$ is nonempty

LP example: production planning

- ▶ x_i units of product i
- ▶ c_i cost per unit
- ▶ a_{ij} amount of resource j used by product i
- ▶ b_j amount of resource j available
- ▶ d_i demand for product i

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & 0 \leq x \leq d\end{array}$$

LP example: production planning

- ▶ x_i units of product i
- ▶ c_i cost per unit
- ▶ a_{ij} amount of resource j used by product i
- ▶ b_j amount of resource j available
- ▶ d_i demand for product i

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & 0 \leq x \leq d\end{array}$$

extensions:

- ▶ fixed cost for producing product i at all?

LP example: production planning

- ▶ x_i units of product i
- ▶ c_i cost per unit
- ▶ a_{ij} amount of resource j used by product i
- ▶ b_j amount of resource j available
- ▶ d_i demand for product i

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & 0 \leq x \leq d\end{array}$$

extensions:

- ▶ fixed cost for producing product i at all?
 $c^T x + f^T z$, $z_i \in \{0, 1\}$, $x_i \leq Mz_i$ for M large

LP inequality form to standard form

standard form to inequality form

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad \rightarrow$$

LP inequality form to standard form

standard form to inequality form

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad \rightarrow \quad \begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & Ax \geq b \\ & -x \leq 0 \end{array}$$

LP inequality form to standard form

standard form to inequality form

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad \rightarrow \quad \begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & Ax \geq b \\ & -x \leq 0 \end{array}$$

inequality form to standard form

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \end{array} \quad \rightarrow$$

LP inequality form to standard form

standard form to inequality form

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad \rightarrow \quad \begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & Ax \geq b \\ & -x \leq 0 \end{array}$$

inequality form to standard form

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \end{array} \quad \rightarrow \quad \begin{array}{ll} \text{minimize} & c^T (x_+ - x_-) \\ \text{subject to} & A(x_+ - x_-) + s = b \\ & s, x_+, x_- \geq 0 \end{array}$$

so both forms have the same expressive power, and feasible sets are polyhedra

Active constraints and variables

active constraints. for constraint set $Ax \leq b$, an **active constraint** i at x is one that holds with equality:

$$a_i^T x = b_i$$

Active constraints and variables

active constraints. for constraint set $Ax \leq b$, an **active constraint** i at x is one that holds with equality:

$$a_i^T x = b_i$$

- ▶ the **active set** $S(x)$ at x is the set of indices of active constraints

$$S(x) = \{i \mid a_i^T x = b_i\}$$

Active constraints and variables

active constraints. for constraint set $Ax \leq b$, an **active constraint** i at x is one that holds with equality:

$$a_i^T x = b_i$$

- ▶ the **active set** $S(x)$ at x is the set of indices of active constraints

$$S(x) = \{i \mid a_i^T x = b_i\}$$

active variables. for nonnegative variable $x \geq 0$, variable i is **active** if $x_i > 0$

example: active slack variables are dual to active constraints

$$Ax \leq b \iff Ax + s = b, s \geq 0$$

$$a_i^T x = b_i \iff s_i = 0$$

constraint i is active \iff slack variable s_i is inactive

Outline

LP standard form

LP inequality form

What kinds of points can be optimal?

Solving LPs

Modeling

Extreme points

define **extreme point**: $x \in \mathbb{R}^n$ is extreme in $C \subset \mathbb{R}^n$ if it cannot be written as a convex combination of other points in C : for $\theta \in [0, 1]$,

$$x \in C \quad \text{and} \quad x = \theta y + (1 - \theta)z \quad \implies \quad x = y = z$$

Extreme points

define **extreme point**: $x \in \mathbb{R}^n$ is extreme in $C \subset \mathbb{R}^n$ if it cannot be written as a convex combination of other points in C : for $\theta \in [0, 1]$,

$$x \in C \quad \text{and} \quad x = \theta y + (1 - \theta)z \quad \implies \quad x = y = z$$

fact: if x^* is the unique optimal solution of $\text{minimize}_{x \in S} c^T x$, then x^* is extreme in the set S .

Extreme points

define **extreme point**: $x \in \mathbb{R}^n$ is extreme in $C \subset \mathbb{R}^n$ if it cannot be written as a convex combination of other points in C : for $\theta \in [0, 1]$,

$$x \in C \quad \text{and} \quad x = \theta y + (1 - \theta)z \quad \implies \quad x = y = z$$

fact: if x^* is the unique optimal solution of $\text{minimize}_{x \in S} c^T x$, then x^* is extreme in the set S .

proof: suppose by way of contradiction that x^* is not extreme in S :

$$\begin{aligned} x^* &= \theta y + (1 - \theta)z \quad \text{for } y, z \in S, \theta \in (0, 1) \\ p^* := c^T x^* &= \theta c^T y + (1 - \theta)c^T z > \theta p^* + (1 - \theta)p^* = p^* \end{aligned}$$

where the inequality follows from the (unique) optimality of x^* . Contradiction!

Extreme points

define **extreme point**: $x \in \mathbb{R}^n$ is extreme in $C \subset \mathbb{R}^n$ if it cannot be written as a convex combination of other points in C : for $\theta \in [0, 1]$,

$$x \in C \quad \text{and} \quad x = \theta y + (1 - \theta)z \quad \implies \quad x = y = z$$

fact: if x^* is the unique optimal solution of $\text{minimize}_{x \in S} c^T x$,
then x^* is extreme in the set S .

proof: suppose by way of contradiction that x^* is not extreme in S :

$$\begin{aligned} x^* &= \theta y + (1 - \theta)z && \text{for } y, z \in S, \theta \in (0, 1) \\ p^* := c^T x^* &= \theta c^T y + (1 - \theta)c^T z > \theta p^* + (1 - \theta)p^* = p^* \end{aligned}$$

where the inequality follows from the (unique) optimality of x^* . Contradiction!

Q: Example of an LP with a non-extreme solution?

Extreme points

define **extreme point**: $x \in \mathbb{R}^n$ is extreme in $C \subset \mathbb{R}^n$ if it cannot be written as a convex combination of other points in C : for $\theta \in [0, 1]$,

$$x \in C \quad \text{and} \quad x = \theta y + (1 - \theta)z \quad \implies \quad x = y = z$$

fact: if x^* is the unique optimal solution of $\text{minimize}_{x \in S} c^T x$, then x^* is extreme in the set S .

proof: suppose by way of contradiction that x^* is not extreme in S :

$$\begin{aligned} x^* &= \theta y + (1 - \theta)z && \text{for } y, z \in S, \theta \in (0, 1) \\ p^* := c^T x^* &= \theta c^T y + (1 - \theta)c^T z > \theta p^* + (1 - \theta)p^* = p^* \end{aligned}$$

where the inequality follows from the (unique) optimality of x^* . Contradiction!

Q: Example of an LP with a non-extreme solution?

Q: Does there always exist an extreme solution?

Vertices

define **vertex**: $x \in \mathbb{R}^n$ is a vertex of set $S \subset \mathbb{R}^n$ if for some vector $c \in \mathbb{R}^n$,

$$c^T x < c^T y \quad \forall y \in S \setminus \{x\}$$

Vertices

define **vertex**: $x \in \mathbb{R}^n$ is a vertex of set $S \subset \mathbb{R}^n$ if for some vector $c \in \mathbb{R}^n$,

$$c^T x < c^T y \quad \forall y \in S \setminus \{x\}$$

interpretation: $\{z : c^T z = c^T x\}$ is a hyperplane that intersects S only at x .
we say this hyperplane **supports** S at x

Vertices

define **vertex**: $x \in \mathbb{R}^n$ is a vertex of set $S \subset \mathbb{R}^n$ if for some vector $c \in \mathbb{R}^n$,

$$c^T x < c^T y \quad \forall y \in S \setminus \{x\}$$

interpretation: $\{z : c^T z = c^T x\}$ is a hyperplane that intersects S only at x .
we say this hyperplane **supports** S at x

fact: x is a vertex of $S \implies x$ is an extreme point of S

Vertices

define **vertex**: $x \in \mathbb{R}^n$ is a vertex of set $S \subset \mathbb{R}^n$ if for some vector $c \in \mathbb{R}^n$,

$$c^T x < c^T y \quad \forall y \in S \setminus \{x\}$$

interpretation: $\{z : c^T z = c^T x\}$ is a hyperplane that intersects S only at x .
we say this hyperplane **supports** S at x

fact: x is a vertex of $S \implies x$ is an extreme point of S

proof:

Vertices

define **vertex**: $x \in \mathbb{R}^n$ is a vertex of set $S \subset \mathbb{R}^n$ if for some vector $c \in \mathbb{R}^n$,

$$c^T x < c^T y \quad \forall y \in S \setminus \{x\}$$

interpretation: $\{z : c^T z = c^T x\}$ is a hyperplane that intersects S only at x .
we say this hyperplane **supports** S at x

fact: x is a vertex of $S \implies x$ is an extreme point of S

proof: x is a vertex of S . suppose its defining vector is c and consider the optimization problem

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & x \in S \end{array}$$

Vertices

define **vertex**: $x \in \mathbb{R}^n$ is a vertex of set $S \subset \mathbb{R}^n$ if for some vector $c \in \mathbb{R}^n$,

$$c^T x < c^T y \quad \forall y \in S \setminus \{x\}$$

interpretation: $\{z : c^T z = c^T x\}$ is a hyperplane that intersects S only at x .
we say this hyperplane **supports** S at x

fact: x is a vertex of $S \implies x$ is an extreme point of S

proof: x is a vertex of S . suppose its defining vector is c and consider the optimization problem

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & x \in S \end{array}$$

x is the unique optimum of this problem, so the proof of this statement follows from the previous proof.

Basic feasible solution

recall the standard form LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad (\text{LP})$$

Basic feasible solution

recall the standard form LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad (\text{LP})$$

define: $x \in \mathbb{R}^n$ is a **basic feasible solution** (BFS) of (LP) if there is a set $S \subset \{1, \dots, n\}$ of m columns so that $A_S \in \mathbb{R}^{m \times m}$ is invertible and

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0, \quad x \geq 0.$$

- ▶ $A_S = \{A_{S_1}, \dots, A_{S_m}\} \in \mathbb{R}^{m \times m}$ is submatrix of A with columns in S
- ▶ \bar{S} is the complement of S in $\{1, \dots, n\}$

Basic feasible solution

recall the standard form LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad (\text{LP})$$

define: $x \in \mathbb{R}^n$ is a **basic feasible solution** (BFS) of (LP) if there is a set $S \subset \{1, \dots, n\}$ of m columns so that $A_S \in \mathbb{R}^{m \times m}$ is invertible and

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0, \quad x \geq 0.$$

- ▶ $A_S = \{A_{S_1}, \dots, A_{S_m}\} \in \mathbb{R}^{m \times m}$ is submatrix of A with columns in S
- ▶ \bar{S} is the complement of S in $\{1, \dots, n\}$
- ▶ two BFS with S, S' are neighbors if they share all but one columns:
 $|S \cap S'| = m - 1$

Basic feasible solution

recall the standard form LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad (\text{LP})$$

define: $x \in \mathbb{R}^n$ is a **basic feasible solution** (BFS) of (LP) if there is a set $S \subset \{1, \dots, n\}$ of m columns so that $A_S \in \mathbb{R}^{m \times m}$ is invertible and

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0, \quad x \geq 0.$$

- ▶ $A_S = \{A_{S_1}, \dots, A_{S_m}\} \in \mathbb{R}^{m \times m}$ is submatrix of A with columns in S
- ▶ \bar{S} is the complement of S in $\{1, \dots, n\}$
- ▶ two BFS with S, S' are neighbors if they share all but one columns:
 $|S \cap S'| = m - 1$

Basic feasible solution

recall the standard form LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad (\text{LP})$$

define: $x \in \mathbb{R}^n$ is a **basic feasible solution** (BFS) of (LP) if there is a set $S \subset \{1, \dots, n\}$ of m columns so that $A_S \in \mathbb{R}^{m \times m}$ is invertible and

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0, \quad x \geq 0.$$

- ▶ $A_S = \{A_{S_1}, \dots, A_{S_m}\} \in \mathbb{R}^{m \times m}$ is submatrix of A with columns in S
- ▶ \bar{S} is the complement of S in $\{1, \dots, n\}$
- ▶ two BFS with S, S' are neighbors if they share all but one columns:
 $|S \cap S'| = m - 1$

Q: how to find a BFS?

Basic feasible solution

recall the standard form LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad (\text{LP})$$

define: $x \in \mathbb{R}^n$ is a **basic feasible solution** (BFS) of (LP) if there is a set $S \subset \{1, \dots, n\}$ of m columns so that $A_S \in \mathbb{R}^{m \times m}$ is invertible and

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0, \quad x \geq 0.$$

- ▶ $A_S = \{A_{S_1}, \dots, A_{S_m}\} \in \mathbb{R}^{m \times m}$ is submatrix of A with columns in S
- ▶ \bar{S} is the complement of S in $\{1, \dots, n\}$
- ▶ two BFS with S, S' are neighbors if they share all but one columns:
 $|S \cap S'| = m - 1$

Q: how to find a BFS?

A: choose m linearly independent columns of A and set $x = A_S^{-1}b$; check $x \geq 0$.

Extreme point \iff vertex \iff BFS

fact. consider the feasible set $F = \{x \mid Ax = b, x \geq 0\}$ in \mathbb{R}^n . the following are equivalent:

- ▶ x is an extreme point of F
- ▶ x is a vertex of F
- ▶ x is a BFS of F

Extreme point \iff vertex \iff BFS

fact. consider the feasible set $F = \{x \mid Ax = b, x \geq 0\}$ in \mathbb{R}^n . the following are equivalent:

- ▶ x is an extreme point of F
- ▶ x is a vertex of F
- ▶ x is a BFS of F

implications: since any polyhedron $Ax \leq b$ can be written as $Ax = b, x \geq 0$,

- ▶ (BFS \implies) a polyhedron has a finite number of extreme points
- ▶ (extreme point \implies) BFS are independent of the representation of the feasible set

Extreme point \iff vertex \iff BFS

fact. consider the feasible set $F = \{x \mid Ax = b, x \geq 0\}$ in \mathbb{R}^n . the following are equivalent:

- ▶ x is an extreme point of F
- ▶ x is a vertex of F
- ▶ x is a BFS of F

implications: since any polyhedron $Ax \leq b$ can be written as $Ax = b, x \geq 0$,

- ▶ (BFS \implies) a polyhedron has a finite number of extreme points
- ▶ (extreme point \implies) BFS are independent of the representation of the feasible set

we have already shown that vertex \implies extreme point. need to show

- ▶ extreme point \implies BFS
- ▶ BFS \implies vertex

Extreme point \implies BFS

we will show the contrapositive: x is not a BFS $\implies x$ is not an extreme point

Extreme point \implies BFS

we will show the contrapositive: x is not a BFS $\implies x$ is not an extreme point

suppose that $\hat{x} \in F$ is feasible ($A\hat{x} = b$ and $\hat{x} \geq 0$) but is not a BFS:

there is no $S \subseteq [n]$ so that A_S is invertible, $\hat{x}_S = A_S^{-1}b$, and $\hat{x}_{\bar{S}} = 0$.

Extreme point \implies BFS

we will show the contrapositive: x is not a BFS $\implies x$ is not an extreme point

suppose that $\hat{x} \in F$ is feasible ($A\hat{x} = b$ and $\hat{x} \geq 0$) but is not a BFS:

there is no $S \subseteq [n]$ so that A_S is invertible, $\hat{x}_S = A_S^{-1}b$, and $\hat{x}_{\bar{S}} = 0$.

consider $\hat{S} = \{i : x_i^* > 0\}$, the active set of variables in \hat{x} .

- ▶ if $A_{\hat{S}}$ were full rank $|\hat{S}|$, either it would be invertible or we could complete $A_{\hat{S}}$ to an invertible A_S with $\hat{S} \subseteq S$, contradicting that \hat{x} is not a BFS,
- ▶ so $A_{\hat{S}}$ has a nontrivial nullspace: there is some $d \in \text{nullspace}(A_{\hat{S}})$, $d_{\hat{S}} \neq 0$.

Extreme point \implies BFS

we will show the contrapositive: x is not a BFS $\implies x$ is not an extreme point

suppose that $\hat{x} \in F$ is feasible ($A\hat{x} = b$ and $\hat{x} \geq 0$) but is not a BFS:

there is no $S \subseteq [n]$ so that A_S is invertible, $\hat{x}_S = A_S^{-1}b$, and $\hat{x}_{\bar{S}} = 0$.

consider $\hat{S} = \{i : x_i^* > 0\}$, the active set of variables in \hat{x} .

- ▶ if $A_{\hat{S}}$ were full rank $|\hat{S}|$, either it would be invertible or we could complete $A_{\hat{S}}$ to an invertible A_S with $\hat{S} \subseteq S$, contradicting that \hat{x} is not a BFS,
- ▶ so $A_{\hat{S}}$ has a nontrivial nullspace: there is some $d \in \text{nullspace}(A_{\hat{S}})$, $d_{\hat{S}} \neq 0$.

extend this vector to $d \in \mathbb{R}^n$ by appending zeros, so $Ad = A_{\hat{S}}d_{\hat{S}} = 0$.

now for $\epsilon \leq \min_{i \in \hat{S}} \hat{x}_i / \max_{i \in \hat{S}} |d_i|$, define $x^+, x^- \in \mathbb{R}^n$ as

$$x^+ = \hat{x} + \epsilon d, \quad x^- = \hat{x} - \epsilon d.$$

these are feasible:

- ▶ $x^+, x^- \geq 0$ by our choice of ϵ ,
- ▶ $Ax^+ = Ax^- = b$ since $Ad = 0$.

Extreme point \implies BFS

we will show the contrapositive: x is not a BFS $\implies x$ is not an extreme point

suppose that $\hat{x} \in F$ is feasible ($A\hat{x} = b$ and $\hat{x} \geq 0$) but is not a BFS:

there is no $S \subseteq [n]$ so that A_S is invertible, $\hat{x}_S = A_S^{-1}b$, and $\hat{x}_{\bar{S}} = 0$.

consider $\hat{S} = \{i : x_i^* > 0\}$, the active set of variables in \hat{x} .

- ▶ if $A_{\hat{S}}$ were full rank $|\hat{S}|$, either it would be invertible or we could complete $A_{\hat{S}}$ to an invertible A_S with $\hat{S} \subseteq S$, contradicting that \hat{x} is not a BFS,
- ▶ so $A_{\hat{S}}$ has a nontrivial nullspace: there is some $d \in \text{nullspace}(A_{\hat{S}})$, $d_{\hat{S}} \neq 0$.

extend this vector to $d \in \mathbb{R}^n$ by appending zeros, so $Ad = A_{\hat{S}}d_{\hat{S}} = 0$.

now for $\epsilon \leq \min_{i \in \hat{S}} \hat{x}_i / \max_{i \in \hat{S}} |d_i|$, define $x^+, x^- \in \mathbb{R}^n$ as

$$x^+ = \hat{x} + \epsilon d, \quad x^- = \hat{x} - \epsilon d.$$

these are feasible:

- ▶ $x^+, x^- \geq 0$ by our choice of ϵ ,
- ▶ $Ax^+ = Ax^- = b$ since $Ad = 0$.

so $\hat{x} = \frac{1}{2}x^+ + \frac{1}{2}x^-$ is not extreme in F

BFS \Rightarrow vertex

suppose x^* is a BFS of F with active set S and A_S invertible. define $c \in \mathbb{R}^n$ as

$$c_i = \begin{cases} 0 & \text{if } i \in S \\ 1 & \text{otherwise} \end{cases}$$

so $c^T x^* = 0$.

BFS \implies vertex

suppose x^* is a BFS of F with active set S and A_S invertible. define $c \in \mathbb{R}^n$ as

$$c_i = \begin{cases} 0 & \text{if } i \in S \\ 1 & \text{otherwise} \end{cases}$$

so $c^T x^* = 0$.

- ▶ x^* is the only point in F supported on S , as $\text{nullspace}(A_S) = 0$,
- ▶ so any other feasible point $x \in F$ has a positive objective value $c^T x > 0$.

hence x^* is a vertex of F with defining vector c .

Outline

LP standard form

LP inequality form

What kinds of points can be optimal?

Solving LPs

Modeling

Solving LPs

algorithms:

- ▶ enumerate all vertices and check
- ▶ fourier-motzkin elimination
- ▶ simplex method
- ▶ ellipsoid method
- ▶ interior point methods
- ▶ first-order methods
- ▶ ...

Solving LPs

algorithms:

- ▶ enumerate all vertices and check
- ▶ fourier-motzkin elimination
- ▶ simplex method
- ▶ ellipsoid method
- ▶ interior point methods
- ▶ first-order methods
- ▶ ...

remarks:

- ▶ enumeration and elimination are simple but not practical
- ▶ simplex was the first practical algorithm; still used today
- ▶ ellipsoid method is the first polynomial-time algorithm; not practical
- ▶ interior point methods are polynomial-time and practical
- ▶ first-order methods are practical and scale to large problems

Example of Fourier-Motzkin elimination

consider the system of inequalities

$$x_1 + 2x_2 \leq 4$$

$$-x_1 + x_2 \leq 1$$

$$x_1, x_2 \geq 0$$

Example of Fourier-Motzkin elimination

consider the system of inequalities

$$x_1 + 2x_2 \leq 4$$

$$-x_1 + x_2 \leq 1$$

$$x_1, x_2 \geq 0$$

we can collect inequalities on x_1 into those bounding it above and below:

$$\{0, x_2 - 1\} \leq x_1 \leq 4 - 2x_2$$

Example of Fourier-Motzkin elimination

consider the system of inequalities

$$x_1 + 2x_2 \leq 4$$

$$-x_1 + x_2 \leq 1$$

$$x_1, x_2 \geq 0$$

we can collect inequalities on x_1 into those bounding it above and below:

$$\{0, x_2 - 1\} \leq x_1 \leq 4 - 2x_2$$

by appending all pairwise inequalities to existing inequalities on x_2 , we recover the feasible set for x_2 :

$$0 \leq 4 - 2x_2$$

$$x_2 - 1 \leq 4 - 2x_2$$

$$x_2 \geq 0$$

$$\implies x_2 \in [0, 5/3].$$

Example of Fourier-Motzkin elimination

consider the system of inequalities

$$x_1 + 2x_2 \leq 4$$

$$-x_1 + x_2 \leq 1$$

$$x_1, x_2 \geq 0$$

we can collect inequalities on x_1 into those bounding it above and below:

$$\{0, x_2 - 1\} \leq x_1 \leq 4 - 2x_2$$

by appending all pairwise inequalities to existing inequalities on x_2 , we recover the feasible set for x_2 :

$$0 \leq 4 - 2x_2$$

$$x_2 - 1 \leq 4 - 2x_2$$

$$x_2 \geq 0$$

$$\implies x_2 \in [0, 5/3].$$

elimination method also shows projection of a polyhedron is a (closed) polyhedron

Enumerate vertices of LP

can generate all extreme points of LP: for each $S \subseteq \{1, \dots, n\}$ with $|S| = m$,

- ▶ $A_S \in \mathbb{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ solve $A_S x_S = b$ for x_S and set $x_{\bar{S}} = 0$
- ▶ if $x_S \geq 0$, then x is a BFS
- ▶ evaluate objective $c^T x$

the best BFS is optimal!

Enumerate vertices of LP

can generate all extreme points of LP: for each $S \subseteq \{1, \dots, n\}$ with $|S| = m$,

- ▶ $A_S \in \mathbb{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ solve $A_S x_S = b$ for x_S and set $x_{\bar{S}} = 0$
- ▶ if $x_S \geq 0$, then x is a BFS
- ▶ evaluate objective $c^T x$

the best BFS is optimal!

problem: how many BFSs are there?

Enumerate vertices of LP

can generate all extreme points of LP: for each $S \subseteq \{1, \dots, n\}$ with $|S| = m$,

- ▶ $A_S \in \mathbb{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ solve $A_S x_S = b$ for x_S and set $x_{\bar{S}} = 0$
- ▶ if $x_S \geq 0$, then x is a BFS
- ▶ evaluate objective $c^T x$

the best BFS is optimal!

problem: how many BFSs are there?

n choose m is $\binom{n}{m} = \frac{n!}{m!(n-m)!}$ (“exponentially many”)

Simplex algorithm

basic idea: local search on the vertices of the feasible set

- ▶ start at BFS x and evaluate objective $c^T x$
- ▶ move to a neighboring BFS x' with better objective $c^T x'$
- ▶ repeat until no improvement possible

Simplex algorithm

basic idea: local search on the vertices of the feasible set

- ▶ start at BFS x and evaluate objective $c^T x$
- ▶ move to a neighboring BFS x' with better objective $c^T x'$
- ▶ repeat until no improvement possible

discuss in groups:

- ▶ how to find an initial BFS?
- ▶ how to find a neighboring BFS with better objective?
- ▶ how to prove optimality?

Finding an initial BFS

solve an auxiliary problem for which a BFS is known:

Finding an initial BFS

solve an auxiliary problem for which a BFS is known:

$$\begin{array}{ll}\text{minimize} & \sum_{i=1}^m z_i \\ \text{subject to} & Ax + Dz = b \\ & x, z \geq 0\end{array}$$

where $D \in \mathbb{R}^{m \times m}$ is a diagonal matrix with $D_{ii} = \mathbf{sign}(b_i)$ for $i = 1, \dots, m$.

Finding an initial BFS

solve an auxiliary problem for which a BFS is known:

$$\begin{array}{ll}\text{minimize} & \sum_{i=1}^m z_i \\ \text{subject to} & Ax + Dz = b \\ & x, z \geq 0\end{array}$$

where $D \in \mathbb{R}^{m \times m}$ is a diagonal matrix with $D_{ii} = \mathbf{sign}(b_i)$ for $i = 1, \dots, m$.

► $x = 0, z = |b|$ is a BFS of this problem

Finding an initial BFS

solve an auxiliary problem for which a BFS is known:

$$\begin{array}{ll}\text{minimize} & \sum_{i=1}^m z_i \\ \text{subject to} & Ax + Dz = b \\ & x, z \geq 0\end{array}$$

where $D \in \mathbb{R}^{m \times m}$ is a diagonal matrix with $D_{ii} = \mathbf{sign}(b_i)$ for $i = 1, \dots, m$.

- ▶ $x = 0, z = |b|$ is a BFS of this problem
- ▶ $(x, z) = (x, 0)$ is a BFS of this problem $\iff x$ is a BFS of the original problem

Find a better neighboring BFS

start with BFS x with active set S , $x_S > 0$. (called a **non-degenerate** BFS.)
construct the **j th basic direction** d^j by turning on variable $j \notin S$

$$x^+ \leftarrow x + \theta d^j, \quad \theta > 0$$

where $d_j^j = 1$ and $d_i^j = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S^j .

Find a better neighboring BFS

start with BFS x with active set S , $x_S > 0$. (called a **non-degenerate** BFS.)
construct the **j th basic direction** d^j by turning on variable $j \notin S$

$$x^+ \leftarrow x + \theta d^j, \quad \theta > 0$$

where $d_j^j = 1$ and $d_i^j = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S^j .

► need to stay feasible wrt equality constraints, so need

$$0 = Ad^j = A_S d_S^j + A_j \implies d_S^j = -A_S^{-1} A_j$$

Find a better neighboring BFS

start with BFS x with active set S , $x_S > 0$. (called a **non-degenerate** BFS.)
construct the **j th basic direction** d^j by turning on variable $j \notin S$

$$x^+ \leftarrow x + \theta d^j, \quad \theta > 0$$

where $d_j^j = 1$ and $d_i^j = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S^j .

- ▶ need to stay feasible wrt equality constraints, so need

$$0 = A d^j = A_S d_S^j + A_j \implies d_S^j = -A_S^{-1} A_j$$

- ▶ as $x_S > 0$ is non-degenerate, $\exists \theta > 0$ st $x^+ \geq 0$

Find a better neighboring BFS

start with BFS x with active set S , $x_S > 0$. (called a **non-degenerate** BFS.)
construct the **j th basic direction** d^j by turning on variable $j \notin S$

$$x^+ \leftarrow x + \theta d^j, \quad \theta > 0$$

where $d_j^j = 1$ and $d_i^j = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S^j .

- ▶ need to stay feasible wrt equality constraints, so need

$$0 = A d^j = A_S d_S^j + A_j \implies d_S^j = -A_S^{-1} A_j$$

- ▶ as $x_S > 0$ is non-degenerate, $\exists \theta > 0$ st $x^+ \geq 0$
- ▶ how does objective change if we move to $x^+ = x + \theta d^j$?

$$c^T x^+ - c^T x = \theta c^T d^j = \theta c_j - \theta c_S^T A_S^{-1} A_j$$

Reduced cost

define **reduced cost** $\bar{c}_j = c_j - c_S^T A_S^{-1} A_j, j \notin S$

Reduced cost

define **reduced cost** $\bar{c}_j = c_j - c_S^T A_S^{-1} A_j, j \notin S$

we will show:

- ▶ if $\bar{c} \geq 0$, x is optimal
- ▶ if x is optimal and nondegenerate ($x_S > 0$), then $\bar{c} \geq 0$

(handling degenerate x is more complicated, we will skip it)

if $\bar{c} \geq 0$, x is optimal

three steps to the proof:

- ▶ every feasible direction at x is contained in $\text{cone}(\{d_j \mid j \notin S\})$

if $\bar{c} \geq 0$, x is optimal

three steps to the proof:

- ▶ every feasible direction at x is contained in **cone**($\{d_j \mid j \notin S\}$)
feasible directions d must satisfy, for some $\theta \geq 0$,

$$A(x + \theta d) = b, \quad x + \theta d \geq 0$$

- ▶ nonnegativity requires $d_j \geq 0$ for $j \notin S$
- ▶ feasibility requires $0 = Ad = A(d_S + \sum_{j \notin S} \alpha_j e_j)$ for some $\alpha \geq 0$
- ▶ solve: $d_S = -A_S^{-1} \sum_{j \notin S} \alpha_j A_j = \sum_{j \notin S} \alpha_j (-A_S^{-1} A_j) = \sum_{j \notin S} \alpha_j d_S^j$
- ▶ so $d = \sum_{j \notin S} \alpha_j (d_S^j + e_j) = \sum_{j \notin S} \alpha_j d^j$

if $\bar{c} \geq 0$, x is optimal

three steps to the proof:

- ▶ every feasible direction at x is contained in $\text{cone}(\{d_j \mid j \notin S\})$
feasible directions d must satisfy, for some $\theta \geq 0$,

$$A(x + \theta d) = b, \quad x + \theta d \geq 0$$

- ▶ nonnegativity requires $d_j \geq 0$ for $j \notin S$
- ▶ feasibility requires $0 = Ad = A(d_S + \sum_{j \notin S} \alpha_j e_j)$ for some $\alpha \geq 0$
- ▶ solve: $d_S = -A_S^{-1} \sum_{j \notin S} \alpha_j A_j = \sum_{j \notin S} \alpha_j (-A_S^{-1} A_j) = \sum_{j \notin S} \alpha_j d_S^j$
- ▶ so $d = \sum_{j \notin S} \alpha_j (d_S^j + e_j) = \sum_{j \notin S} \alpha_j d^j$
- ▶ the feasible set $F = \{x \mid Ax = b, x \geq 0\} \subseteq x + \text{cone}(\{d_j \mid j \notin S\})$

if $\bar{c} \geq 0$, x is optimal

three steps to the proof:

- ▶ every feasible direction at x is contained in $\text{cone}(\{d_j \mid j \notin S\})$
feasible directions d must satisfy, for some $\theta \geq 0$,

$$A(x + \theta d) = b, \quad x + \theta d \geq 0$$

- ▶ nonnegativity requires $d_j \geq 0$ for $j \notin S$
- ▶ feasibility requires $0 = Ad = A(d_S + \sum_{j \notin S} \alpha_j e_j)$ for some $\alpha \geq 0$
- ▶ solve: $d_S = -A_S^{-1} \sum_{j \notin S} \alpha_j A_j = \sum_{j \notin S} \alpha_j (-A_S^{-1} A_j) = \sum_{j \notin S} \alpha_j d_S^j$
- ▶ so $d = \sum_{j \notin S} \alpha_j (d_S^j + e_j) = \sum_{j \notin S} \alpha_j d^j$
- ▶ the feasible set $F = \{x \mid Ax = b, x \geq 0\} \subseteq x + \text{cone}(\{d_j \mid j \notin S\})$
by convexity (consider segment from x to $x' \in F$)

if $\bar{c} \geq 0$, x is optimal

three steps to the proof:

- ▶ every feasible direction at x is contained in $\text{cone}(\{d_j \mid j \notin S\})$
feasible directions d must satisfy, for some $\theta \geq 0$,

$$A(x + \theta d) = b, \quad x + \theta d \geq 0$$

- ▶ nonnegativity requires $d_j \geq 0$ for $j \notin S$
- ▶ feasibility requires $0 = Ad = A(d_S + \sum_{j \notin S} \alpha_j e_j)$ for some $\alpha \geq 0$
- ▶ solve: $d_S = -A_S^{-1} \sum_{j \notin S} \alpha_j A_j = \sum_{j \notin S} \alpha_j (-A_S^{-1} A_j) = \sum_{j \notin S} \alpha_j d_S^j$
- ▶ so $d = \sum_{j \notin S} \alpha_j (d_S^j + e_j) = \sum_{j \notin S} \alpha_j d^j$
- ▶ the feasible set $F = \{x \mid Ax = b, x \geq 0\} \subseteq x + \text{cone}(\{d_j \mid j \notin S\})$
by convexity (consider segment from x to $x' \in F$)
- ▶ so

$$\begin{aligned} p^* = \min_{x' \in F} c^T x' &\geq \min_{\alpha \geq 0} c^T (x + \sum_{j \notin S} \alpha_j d_j) \\ &= c^T x + \min \sum \alpha_j \bar{c}_j = c^T x \end{aligned}$$

Outline

LP standard form

LP inequality form

What kinds of points can be optimal?

Solving LPs

Modeling

LP solvers

practical solvers for MILP:

- ▶ Gurobi and COPT are state-of-the-art commercial solvers
- ▶ GLPK and SCIP are free solvers that are not as fast

LP solvers

practical solvers for MILP:

- ▶ Gurobi and COPT are state-of-the-art commercial solvers
- ▶ GLPK and SCIP are free solvers that are not as fast
- ▶ gurobipy is a python interface to Gurobi
- ▶ CVX* (including CVXPY in python) are modeling languages that call solvers with good support for convex problems

LP solvers

practical solvers for MILP:

- ▶ Gurobi and COPT are state-of-the-art commercial solvers
- ▶ GLPK and SCIP are free solvers that are not as fast
- ▶ gurobipy is a python interface to Gurobi
- ▶ CVX* (including CVXPY in python) are modeling languages that call solvers with good support for convex problems
- ▶ OptiMUS is a LLM-based modeling tool for MILP that produces gurobipy code
<https://optimus-solver.com/dashboard>

LP solvers

practical solvers for MILP:

- ▶ Gurobi and COPT are state-of-the-art commercial solvers
- ▶ GLPK and SCIP are free solvers that are not as fast
- ▶ gurobipy is a python interface to Gurobi
- ▶ CVX* (including CVXPY in python) are modeling languages that call solvers with good support for convex problems
- ▶ OptiMUS is a LLM-based modeling tool for MILP that produces gurobipy code
<https://optimus-solver.com/dashboard>
- ▶ JuliaOpt/JuMP is a modeling language in Julia that calls solvers and is super speedy for MILP applications

LP solvers

practical solvers for MILP:

- ▶ Gurobi and COPT are state-of-the-art commercial solvers
- ▶ GLPK and SCIP are free solvers that are not as fast
- ▶ gurobipy is a python interface to Gurobi
- ▶ CVX* (including CVXPY in python) are modeling languages that call solvers with good support for convex problems
- ▶ OptiMUS is a LLM-based modeling tool for MILP that produces gurobipy code
<https://optimus-solver.com/dashboard>
- ▶ JuliaOpt/JuMP is a modeling language in Julia that calls solvers and is super speedy for MILP applications demos:
 - ▶ power systems
https://jump.dev/JuMP.jl/stable/tutorials/applications/power_systems/

Oro Verde case + tutorial

<https://github.com/stanford-cme-307/demos/tree/main/gurobipy>

Modeling challenges

model the following as standard form LPs:

1. **inequality constraints.** $Ax \leq b$
2. **free variable.** $x \in \mathbb{R}$
3. **absolute value.** constraint $|x| \leq 10$
4. **piecewise linear.** objective $\max(x_1, x_2)$
5. **assignment.** e.g., every class is assigned exactly one classroom
6. **logic.** e.g., class enrollment \leq capacity of assigned room
7. **(big-M).** $Ax \leq b$ if $x \geq 10$
8. **flow.** e.g., the least cost way to ship an item from s to t

Modeling challenges

model the following as standard form LPs:

1. **inequality constraints.** $Ax \leq b$
2. **free variable.** $x \in \mathbb{R}$
3. **absolute value.** constraint $|x| \leq 10$
4. **piecewise linear.** objective $\max(x_1, x_2)$
5. **assignment.** e.g., every class is assigned exactly one classroom
6. **logic.** e.g., class enrollment \leq capacity of assigned room
7. **(big-M).** $Ax \leq b$ if $x \geq 10$
8. **flow.** e.g., the least cost way to ship an item from s to t

(see chapter 1 of Bertsimas and Tsitsiklis for more details on 1–6. see

[https://github.com/stanford-cme-307/demos/blob/main/](https://github.com/stanford-cme-307/demos/blob/main/Multicast_Routing_Demonstration.ipynb)

[Multicast_Routing_Demonstration.ipynb](https://github.com/stanford-cme-307/demos/blob/main/Multicast_Routing_Demonstration.ipynb) for a detailed treatment of a flow problem.)

Use slack variables to represent inequality constraints

to represent the following problem in standard form,

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & x \geq 0\end{array}$$

Use slack variables to represent inequality constraints

to represent the following problem in standard form,

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & x \geq 0\end{array}$$

introduce slack variable $s \in \mathbb{R}^m$: $Ax + s = b, s \geq 0 \iff Ax \leq b$

$$\begin{array}{ll}\text{minimize} & c^T x + 0^T s \\ \text{subject to} & Ax + s = b \\ & x, s \geq 0\end{array}$$

Split variable into parts to represent free variables

to represent the following problem in standard form,

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b\end{array}$$

Split variable into parts to represent free variables

to represent the following problem in standard form,

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b\end{array}$$

introduce positive variables x_+, x_- so $x = x_+ - x_-$:

$$\begin{array}{ll}\text{minimize} & c^T x_+ - c^T x_- \\ \text{subject to} & Ax_+ - Ax_- = b \\ & x_+, x_- \geq 0\end{array}$$

Use epigraph variables to handle absolute value

to represent the following problem in standard form,

$$\begin{array}{ll}\text{minimize} & \|x\|_1 = \sum_{i=1}^n |x_i| \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

Use epigraph variables to handle absolute value

to represent the following problem in standard form,

$$\begin{array}{ll}\text{minimize} & \|x\|_1 = \sum_{i=1}^n |x_i| \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

introduce epigraph variable $t \in \mathbb{R}^n$ so $|x_i| \leq t_i$:

$$\begin{array}{ll}\text{minimize} & \mathbf{1}^T t \\ \text{subject to} & Ax = b \\ & -t \leq x \leq t \\ & x, t \geq 0\end{array}$$

verify these constraints ensure $|x_i| \leq t_i$.

Use epigraph variables to handle absolute value

to represent the following problem in standard form,

$$\begin{array}{ll}\text{minimize} & \|x\|_1 = \sum_{i=1}^n |x_i| \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

introduce epigraph variable $t \in \mathbb{R}^n$ so $|x_i| \leq t_i$:

$$\begin{array}{ll}\text{minimize} & \mathbf{1}^T t \\ \text{subject to} & Ax = b \\ & -t \leq x \leq t \\ & x, t \geq 0\end{array}$$

verify these constraints ensure $|x_i| \leq t_i$.

Q: Why does this work? For what kinds of functions can we use this trick?

Use binary variables to handle assignment

every class is assigned exactly one classroom:

define variable $X_{ij} \in \{0, 1\}$ for each class $i = 1, \dots, n$ and room $j = 1, \dots, m$

$$X_{ij} = \begin{cases} 1 & \text{class } i \text{ is assigned to room } j \\ 0 & \text{otherwise} \end{cases}$$

Use binary variables to handle assignment

every class is assigned exactly one classroom:

define variable $X_{ij} \in \{0, 1\}$ for each class $i = 1, \dots, n$ and room $j = 1, \dots, m$

$$X_{ij} = \begin{cases} 1 & \text{class } i \text{ is assigned to room } j \\ 0 & \text{otherwise} \end{cases}$$

now solve the problem

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^n \sum_{j=1}^m C_{ij} X_{ij} \\ \text{subject to} & \sum_{j=1}^m X_{ij} = 1, \forall i \quad (\text{every class assigned one room}) \\ & \sum_{i=1}^n X_{ij} \leq 1, \forall j \quad (\text{no more than one class per room}) \\ & X_{ij} \in \{0, 1\} \quad (\text{binary variables}) \end{array}$$

where C_{ij} is the cost of assigning class i to room j .

Use binary variables to handle logic

model class enrollment $p_i \leq$ capacity c_j of assigned room:

define variable $X_{ij} \in \{0, 1\}$ for each class $i = 1, \dots, n$ and room $j = 1, \dots, m$

$$X_{ij} = \begin{cases} 1 & \text{class } i \text{ is assigned to room } j \\ 0 & \text{otherwise} \end{cases}$$

Use binary variables to handle logic

model class enrollment $p_i \leq$ capacity c_j of assigned room:

define variable $X_{ij} \in \{0, 1\}$ for each class $i = 1, \dots, n$ and room $j = 1, \dots, m$

$$X_{ij} = \begin{cases} 1 & \text{class } i \text{ is assigned to room } j \\ 0 & \text{otherwise} \end{cases}$$

solve the problem

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^n \sum_{j=1}^m C_{ij} X_{ij} \\ \text{subject to} & \sum_{j=1}^m X_{ij} = 1, \forall i \quad (\text{every class assigned one room}) \\ & \sum_{i=1}^n X_{ij} \leq 1, \forall j \quad (\text{no more than one class per room}) \\ & \sum_{i=1}^n p_i X_{ij} \leq c_j, \forall j \quad (\text{capacity constraint}) \\ & X_{ij} \in \{0, 1\} \quad (\text{binary variables}) \end{array}$$

where C_{ij} is the cost of assigning class i to room j .

Use binary variables to handle logic

model class enrollment $p_i \leq$ capacity c_j of assigned room:

define variable $X_{ij} \in \{0, 1\}$ for each class $i = 1, \dots, n$ and room $j = 1, \dots, m$

$$X_{ij} = \begin{cases} 1 & \text{class } i \text{ is assigned to room } j \\ 0 & \text{otherwise} \end{cases}$$

solve the problem

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^n \sum_{j=1}^m C_{ij} X_{ij} \\ \text{subject to} & \sum_{j=1}^m X_{ij} = 1, \forall i \quad (\text{every class assigned one room}) \\ & \sum_{i=1}^n X_{ij} \leq 1, \forall j \quad (\text{no more than one class per room}) \\ & \sum_{i=1}^n p_i X_{ij} \leq c_j, \forall j \quad (\text{capacity constraint}) \\ & X_{ij} \in \{0, 1\} \quad (\text{binary variables}) \end{array}$$

where C_{ij} is the cost of assigning class i to room j .

what if we want enrollment p to be a variable, too?

...or use a big-M relaxation!

model class enrollment $p_i \leq$ capacity c_j of assigned room:

define variable $X_{ij} \in \{0, 1\}$ for each class $i = 1, \dots, n$ and room $j = 1, \dots, m$

$$X_{ij} = \begin{cases} 1 & \text{class } i \text{ is assigned to room } j \\ 0 & \text{otherwise} \end{cases}$$

suppose M is a very large number.

...or use a big-M relaxation!

model class enrollment $p_i \leq$ capacity c_j of assigned room:

define variable $X_{ij} \in \{0, 1\}$ for each class $i = 1, \dots, n$ and room $j = 1, \dots, m$

$$X_{ij} = \begin{cases} 1 & \text{class } i \text{ is assigned to room } j \\ 0 & \text{otherwise} \end{cases}$$

suppose M is a very large number. solve the problem

$$\begin{array}{ll} \text{minimize} & \sum_{i=1}^n \sum_{j=1}^m C_{ij} X_{ij} \\ \text{subject to} & \sum_{i=1}^n X_{ij} = 1, \forall j \quad (\text{every class assigned one room}) \\ & \sum_{j=1}^m X_{ij} = 1, \forall i \quad (\text{no more than one class per room}) \\ & p_i \leq c_j + (1 - X_{ij})M, \forall i, j \quad (\text{capacity constraint}) \\ & X_{ij} \in \{0, 1\} \quad (\text{binary variables}) \end{array}$$

where C_{ij} is the cost of assigning class i to room j .