

# CME 307 / MS&E 311: Optimization

## Interior Point Methods

Professor Udell

Management Science and Engineering  
Stanford

November 18, 2024

slides developed with Prof. Luiz-Rafael Santos, UFSC <https://lrsantos11.github.io/>

## Convex optimization

### Convex optimization problem

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & g(x) \leq 0 \\ & Ax = b\end{array}$$

where  $f : \mathbf{R}^n \rightarrow \mathbf{R}$ ,  $g : \mathbf{R}^n \rightarrow \mathbf{R}^p$  are smooth and convex,  $A \in \mathbf{R}^{m \times n}$  is full rank.

## Convex optimization

### Convex optimization problem

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & g(x) \leq 0 \\ & Ax = b\end{array}$$

where  $f : \mathbf{R}^n \rightarrow \mathbf{R}$ ,  $g : \mathbf{R}^n \rightarrow \mathbf{R}^p$  are smooth and convex,  $A \in \mathbf{R}^{m \times n}$  is full rank.

KKT conditions:

## Convex optimization

### Convex optimization problem

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & g(x) \leq 0 \quad s \in \mathbf{R}^p \\ & Ax = b \quad y \in \mathbf{R}^m\end{array}$$

where  $f : \mathbf{R}^n \rightarrow \mathbf{R}$ ,  $g : \mathbf{R}^n \rightarrow \mathbf{R}^p$  are smooth and convex,  $A \in \mathbf{R}^{m \times n}$  is full rank.

KKT conditions:

## Convex optimization

### Convex optimization problem

$$\begin{array}{ll}\text{minimize} & f(x) \\ \text{subject to} & g(x) \leq 0 \quad s \in \mathbf{R}^p \\ & Ax = b \quad y \in \mathbf{R}^m\end{array}$$

where  $f : \mathbf{R}^n \rightarrow \mathbf{R}$ ,  $g : \mathbf{R}^n \rightarrow \mathbf{R}^p$  are smooth and convex,  $A \in \mathbf{R}^{m \times n}$  is full rank.

KKT conditions:

$$\begin{aligned}\nabla f(x) + A^T y + (\nabla g(x))^T s &= 0 \\ Ax &= b \\ g(x) &\leq 0 \\ s &\geq 0 \\ s_j g_j(x) &= 0, \quad j = 1, \dots, p\end{aligned}$$

# Outline

IPM for linear and quadratic programs

IPM for convex nonlinear programming

IPM for conic optimization

## IPM for linear and quadratic programs

### Linear/Quadratic Program

$$\begin{array}{ll}\text{minimize} & c^\top x + \frac{1}{2}x^\top Qx \\ \text{subject to} & Ax = b, \\ & x \geq 0,\end{array}$$

where  $Q \in \mathbf{S}_+^n$ , and  $A \in \mathbf{R}^{m \times n}$  is full-rank.

## IPM for linear and quadratic programs

### Linear/Quadratic Program

$$\begin{array}{ll}\text{minimize} & c^\top x + \frac{1}{2}x^\top Qx \\ \text{subject to} & Ax = b, \\ & x \geq 0,\end{array}$$

where  $Q \in \mathbf{S}_+^n$ , and  $A \in \mathbf{R}^{m \times n}$  is full-rank.

- $\mathcal{P} := \{x \in \mathbf{R}^n \mid Ax = b, x \geq 0\}$  is a polyhedron.



## IPM for linear and quadratic programs

### Linear/Quadratic Program

$$\begin{array}{ll}\text{minimize} & c^\top x + \frac{1}{2}x^\top Qx \\ \text{subject to} & Ax = b, \\ & x \geq 0,\end{array}$$

where  $Q \in \mathbf{S}_+^n$ , and  $A \in \mathbf{R}^{m \times n}$  is full-rank.

- ▶  $\mathcal{P} := \{x \in \mathbf{R}^n \mid Ax = b, x \geq 0\}$  is a polyhedron.
- ▶ If  $Q = 0$ , problem is a linear program.

## IPM for linear and quadratic programs

### Linear/Quadratic Program

$$\begin{array}{ll}\text{minimize} & c^\top x + \frac{1}{2}x^\top Qx \\ \text{subject to} & Ax = b, \\ & x \geq 0,\end{array}$$

where  $Q \in \mathbf{S}_+^n$ , and  $A \in \mathbf{R}^{m \times n}$  is full-rank.

- ▶  $\mathcal{P} := \{x \in \mathbf{R}^n \mid Ax = b, x \geq 0\}$  is a polyhedron.
- ▶ If  $Q = 0$ , problem is a linear program.

**How to solve LP/QP problems?**

# IPM for linear and quadratic programs

## Linear/Quadratic Program

$$\begin{array}{ll}\text{minimize} & c^\top x + \frac{1}{2}x^\top Qx \\ \text{subject to} & Ax = b, \\ & x \geq 0,\end{array}$$

where  $Q \in \mathbf{S}_+^n$ , and  $A \in \mathbf{R}^{m \times n}$  is full-rank.

- ▶  $\mathcal{P} := \{x \in \mathbf{R}^n \mid Ax = b, x \geq 0\}$  is a polyhedron.
- ▶ If  $Q = 0$ , problem is a linear program.

**How to solve LP/QP problems?**

Simplex: vertex to vertex  
IPM: go through the middle!



# IPM for linear and quadratic programs

## Linear/Quadratic Program

$$\begin{array}{ll}\text{minimize} & c^\top x + \frac{1}{2}x^\top Qx \\ \text{subject to} & Ax = b, \\ & x \geq 0,\end{array}$$

where  $Q \in \mathbf{S}_+^n$ , and  $A \in \mathbf{R}^{m \times n}$  is full-rank.

- ▶  $\mathcal{P} := \{x \in \mathbf{R}^n \mid Ax = b, x \geq 0\}$  is a polyhedron.
- ▶ If  $Q = 0$ , problem is a linear program.

## How to solve LP/QP problems?

Advantages of vertex solution vs interior solution?

Simplex: vertex to vertex  
IPM: go through the middle!



## Building blocks of IPM

### Ingredients for Interior Point Method

- ▶ Duality theory: Lagrangian function; KKT (first order optimality) condition.
- ▶ Barrier function: logarithmic barrier.
- ▶ Newton's method (and a good linear solver)

## Building blocks of IPM

### Ingredients for Interior Point Method

- ▶ Duality theory: Lagrangian function; KKT (first order optimality) condition.
- ▶ Barrier function: logarithmic barrier.
- ▶ Newton's method (and a good linear solver)

### The reward: fantastic convergence properties!

- ▶ Theoretical:  $O(\sqrt{n} \log(1/\varepsilon))$  iterations
- ▶ Practical:  $O(\log n \log(1/\varepsilon))$  iterations

(but the per-iteration cost may be high due to the Newton solve: often  $O(n^3)$ )

## IPM: algorithmic template

### IPM procedure

- ▶ replace inequalities with log barriers;
- ▶ form the Lagrangian;
- ▶ write down the KKT conditions of the perturbed problem;
- ▶ find one (or more) directions using [Newton's method](#) on the KKT system;
- ▶ (decide how to combine the directions and) compute a stepsize.

## Duality and KKT conditions

### Primal-dual QPs

#### Primal problem

$$\begin{array}{ll}\text{minimize} & c^\top x + \frac{1}{2}x^\top Qx \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

#### Dual problem

$$\begin{array}{ll}\text{maximize} & b^\top y - \frac{1}{2}x^\top Qx \\ \text{subject to} & A^\top y + s - Qx = c \\ & s \geq 0\end{array}$$



## Duality and KKT conditions

### Primal-dual QPs

#### Primal problem

$$\begin{aligned} \text{minimize} \quad & c^\top x + \frac{1}{2} x^\top Q x \\ \text{subject to} \quad & Ax = b \\ & x \geq 0 \end{aligned}$$

#### Dual problem

$$\begin{aligned} \text{maximize} \quad & b^\top y - \frac{1}{2} x^\top Q x \\ \text{subject to} \quad & A^\top y + s - Qx = c \\ & s \geq 0 \end{aligned}$$


### KKT conditions

$$\begin{aligned} Ax &= b && \triangleright (\text{primal feasibility}) \\ A^\top y + s - Qx &= c && \triangleright (\text{dual feasibility}) \\ XSe &= 0 && \triangleright (\text{complementarity: } x_i s_i = 0, i = 1, \dots, n) \\ (x, s) &\geq 0 \end{aligned}$$

where  $X = \mathbf{diag}(x_1, \dots, x_n)$ ,  $S = \mathbf{diag}(s_1, \dots, s_n) \in \mathbf{R}^{n \times n}$ , and  $e = (1, \dots, 1) \in \mathbf{R}^n$ .


## Logarithmic barrier

$-\ln x_j$   
replaces the inequality  
 $x_j \geq 0$



## Logarithmic barrier

$-\ln x_j$   
replaces the inequality  
 $x_j \geq 0$



$$\text{minimize} \quad -\sum_{j=1}^n \ln x_j \quad \Longleftrightarrow \quad \text{maximize} \quad \prod_{1 \leq j \leq n} x_j$$

$\Rightarrow$  keeps every entry of  $x$  away from 0.

## Barrier primal QP

Step 1: replace inequality constraints by barrier

Replace the primal QP

$$\begin{array}{ll}\text{minimize} & c^T x + \frac{1}{2} x^T Q x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

with the barrier primal QP

$$\begin{array}{ll}\text{minimize} & c^T x + \frac{1}{2} x^T Q x - \mu \sum_{j=1}^n \ln x_j \\ \text{subject to} & Ax = b\end{array}$$

## Logarithmic barrier and stationarity

Step 2: remove equality constraints using Lagrangian

$$\mathcal{L}(x, y, \mu) = c^\top x + \frac{1}{2} x^\top Q x - y^\top (Ax - b) - \mu \sum_{j=1}^n \ln x_j$$

## Logarithmic barrier and stationarity

Step 2: remove equality constraints using Lagrangian

$$\mathcal{L}(x, y, \mu) = c^\top x + \frac{1}{2} x^\top Q x - y^\top (Ax - b) - \mu \sum_{j=1}^n \ln x_j$$

A stationary point  $(x, y, \mu)$  of the Lagrangian satisfies

$$\nabla_x \mathcal{L}(x, y, \mu) = 0 \qquad = c + Qx - A^\top y - \mu X^{-1} e$$

with  $X^{-1} = \mathbf{diag}(x_1^{-1}, \dots, x_n^{-1}) \in \mathbf{R}^{n \times n}, (x_j > 0)$ .

## KKT conditions for barrier problem

- Define  $s := \mu X^{-1}e$ , which implies  $XSe = \mu e$ , to get

KKT <sub>$\mu$</sub>

$$Ax = b$$

$$A^\top y + s - Qx = c$$

$$XSe = \mu e$$

$$(x, s) > 0$$

## KKT conditions for barrier problem

- Define  $s := \mu X^{-1}e$ , which implies  $XSe = \mu e$ , to get

KKT <sub>$\mu$</sub>

$$\begin{aligned}Ax &= b \\ A^\top y + s - Qx &= c \\ XSe &= \mu e \\ (x, s) &> 0\end{aligned}$$

KKT

$$\begin{aligned}Ax &= b \\ A^\top y + s - Qx &= c \\ XSe &= 0 \\ (x, s) &\geq 0\end{aligned}$$



## KKT conditions for barrier problem

- Define  $s := \mu X^{-1}e$ , which implies  $XSe = \mu e$ , to get

KKT <sub>$\mu$</sub>

$$\begin{aligned}Ax &= b \\ A^\top y + s - Qx &= c \\ XSe &= \mu e \\ (x, s) &> 0\end{aligned}$$

KKT

$$\begin{aligned}Ax &= b \\ A^\top y + s - Qx &= c \\ XSe &= 0 \\ (x, s) &\geq 0\end{aligned}$$

KKT <sub>$\mu$</sub>   $\rightarrow$  KKT as  $\mu \rightarrow 0$ .

## Central path (LP case)

- ▶ Parameter  $\mu$  controls the distance to optimality

$$c^\top x - b^\top y = c^\top x - x^\top A^\top y = x^\top s = n\mu$$

## Central path (LP case)

- ▶ Parameter  $\mu$  controls the distance to optimality

$$c^\top x - b^\top y = c^\top x - x^\top A^\top y = x^\top s = n\mu$$

- ▶ Analytic center ( $\mu$ -center): unique point

$$(x(\mu), y(\mu), s(\mu)), \quad x(\mu) > 0, \quad s(\mu) > 0$$

that satisfies the  $\text{KKT}_\mu$  conditions.

## Central path (LP case)

- ▶ Parameter  $\mu$  controls the distance to optimality

$$c^\top x - b^\top y = c^\top x - x^\top A^\top y = x^\top s = n\mu$$

- ▶ Analytic center ( $\mu$ -center): unique point


$$(x(\mu), y(\mu), s(\mu)), \quad x(\mu) > 0, \quad s(\mu) > 0$$

that satisfies the  $\text{KKT}_\mu$  conditions.


- ▶ The curve

$$\mathcal{C}_\mu = \{(x(\mu), y(\mu), s(\mu)) \mid \mu > 0\}$$


is called the primal-dual central path.




(a)  $\mu=\infty$



(b)  $\mu=1$



(c)  $\mu=0.01$



(d) central path

## Recall Newton's method for nonlinear equation

- For  $F : \mathbf{R}^n \rightarrow \mathbf{R}^n$  smooth, solve  $F(x) = 0$ .

## Recall Newton's method for nonlinear equation


- ▶ For  $F : \mathbf{R}^n \rightarrow \mathbf{R}^n$  smooth, solve  $F(x) = 0$ .
- ▶ Newton's method: define Jacobian  $J_F(x)$  so  $J_F(x)_{ij} = \frac{\partial F_i}{\partial x_j}$ , and iterate

$$x^{k+1} = x^k - \alpha_k J_F(x^k)^{-1} F(x^k)$$

## Recall Newton's method for nonlinear equation

- ▶ For  $F : \mathbf{R}^n \rightarrow \mathbf{R}^n$  smooth, solve  $F(x) = 0$ .
- ▶ Newton's method: define Jacobian  $J_F(x)$  so  $J_F(x)_{ij} = \frac{\partial F_i}{\partial x_j}$ , and iterate

$$x^{k+1} = x^k - \alpha_k J_F(x^k)^{-1} F(x^k)$$





## Apply Newton Method to $\text{KKT}_\mu$

The first order optimality conditions for the barrier problem form a large system of nonlinear equations:

$$F(x, y, s) = 0,$$

where  $F : \mathbf{R}^{2n+m} \mapsto \mathbf{R}^{2n+m}$  is defined as

$$F(x, y, s) = \begin{bmatrix} Ax & -b \\ A^\top y + s - Qx & -c \\ XSe & -\mu e \end{bmatrix}$$

## Apply Newton Method to $\text{KKT}_\mu$

The first order optimality conditions for the barrier problem form a large system of nonlinear equations:

$$F(x, y, s) = 0,$$

where  $F : \mathbf{R}^{2n+m} \mapsto \mathbf{R}^{2n+m}$  is defined as

$$F(x, y, s) = \begin{bmatrix} Ax & -b \\ A^\top y + s - Qx & -c \\ XSe & -\mu e \end{bmatrix}$$

- ▶ The first two blocks are **linear**.
- ▶ The last block, corresponding to the complementarity condition, is **nonlinear**.
- ▶ Jacobian is

$$J_F(x, y, s) = \begin{bmatrix} A & 0 & 0 \\ -Q & A^\top & I \\ S & 0 & X \end{bmatrix}$$

## Interior-point QP Algorithm

### IPM Framework

Fix the barrier parameter  $\mu$  and make *one* (damped) Newton step towards the solution of  $\text{KKT}_\mu$ . Then reduce the barrier parameter  $\mu$  and repeat.

## Interior-point QP Algorithm

### IPM Framework

Fix the barrier parameter  $\mu$  and make *one* (damped) Newton step towards the solution of  $\text{KKT}_\mu$ . Then reduce the barrier parameter  $\mu$  and repeat.

- ▶ Given  $(x_0, y_0, s_0)$  feasible,  $\mu_0 = \frac{1}{n}(x^0)^\top s^0$

## Interior-point QP Algorithm

### IPM Framework

Fix the barrier parameter  $\mu$  and make *one* (damped) Newton step towards the solution of  $\text{KKT}_\mu$ . Then reduce the barrier parameter  $\mu$  and repeat.

- ▶ Given  $(x_0, y_0, s_0)$  feasible,  $\mu_0 = \frac{1}{n}(x^0)^\top s^0$
- ▶ For  $k = 1, 2, \dots$

## Interior-point QP Algorithm

### IPM Framework

Fix the barrier parameter  $\mu$  and make *one* (damped) Newton step towards the solution of  $\text{KKT}_\mu$ . Then reduce the barrier parameter  $\mu$  and repeat.

- ▶ Given  $(x_0, y_0, s_0)$  feasible,  $\mu_0 = \frac{1}{n}(x^0)^\top s^0$
- ▶ For  $k = 1, 2, \dots$ 
  - ▶  $\mu_k = \sigma \mu_{k-1}$ , where  $\sigma \in (0, 1)$

## Interior-point QP Algorithm

### IPM Framework

Fix the barrier parameter  $\mu$  and make *one* (damped) Newton step towards the solution of  $\text{KKT}_\mu$ . Then reduce the barrier parameter  $\mu$  and repeat.

- ▶ Given  $(x_0, y_0, s_0)$  feasible,  $\mu_0 = \frac{1}{n}(x^0)^\top s^0$
- ▶ For  $k = 1, 2, \dots$ 
  - ▶  $\mu_k = \sigma \mu_{k-1}$ , where  $\sigma \in (0, 1)$
  - ▶ Find Newton direction  $(\Delta x^k, \Delta y^k, \Delta s^k)$  by solving

$$\begin{bmatrix} A & 0 & 0 \\ -Q & A^\top & I \\ S^k & 0 & X^k \end{bmatrix} \begin{bmatrix} \Delta x^k \\ \Delta y^k \\ \Delta s^k \end{bmatrix} = \begin{bmatrix} b - Ax^k \\ c - A^\top y^k - s^k + Qx^k \\ \mu_k e - X^k S^k e \end{bmatrix}$$

## Interior-point QP Algorithm

### IPM Framework

Fix the barrier parameter  $\mu$  and make *one* (damped) Newton step towards the solution of  $\text{KKT}_\mu$ . Then reduce the barrier parameter  $\mu$  and repeat.

- ▶ Given  $(x_0, y_0, s_0)$  feasible,  $\mu_0 = \frac{1}{n}(x^0)^\top s^0$
- ▶ For  $k = 1, 2, \dots$ 
  - ▶  $\mu_k = \sigma \mu_{k-1}$ , where  $\sigma \in (0, 1)$
  - ▶ Find Newton direction  $(\Delta x^k, \Delta y^k, \Delta s^k)$  by solving

$$\begin{bmatrix} A & 0 & 0 \\ -Q & A^\top & I \\ S^k & 0 & X^k \end{bmatrix} \begin{bmatrix} \Delta x^k \\ \Delta y^k \\ \Delta s^k \end{bmatrix} = \begin{bmatrix} b - Ax^k \\ c - A^\top y^k - s^k + Qx^k \\ \mu_k e - X^k S^k e \end{bmatrix}$$

- ▶ Find step length  $\alpha_k$  so  $(x^k + \alpha_k \Delta x^k, y^k + \alpha_k \Delta y^k, s^k + \alpha_k \Delta s^k)$  is feasible.



## Interior-point QP Algorithm

### IPM Framework

Fix the barrier parameter  $\mu$  and make *one* (damped) Newton step towards the solution of  $\text{KKT}_\mu$ . Then reduce the barrier parameter  $\mu$  and repeat.

- ▶ Given  $(x_0, y_0, s_0)$  feasible,  $\mu_0 = \frac{1}{n}(x^0)^\top s^0$
- ▶ For  $k = 1, 2, \dots$ 
  - ▶  $\mu_k = \sigma \mu_{k-1}$ , where  $\sigma \in (0, 1)$
  - ▶ Find Newton direction  $(\Delta x^k, \Delta y^k, \Delta s^k)$  by solving

$$\begin{bmatrix} A & 0 & 0 \\ -Q & A^\top & I \\ S^k & 0 & X^k \end{bmatrix} \begin{bmatrix} \Delta x^k \\ \Delta y^k \\ \Delta s^k \end{bmatrix} = \begin{bmatrix} b - Ax^k \\ c - A^\top y^k - s^k + Qx^k \\ \mu_k e - X^k S^k e \end{bmatrix}$$

- ▶ Find step length  $\alpha_k$  so  $(x^k + \alpha_k \Delta x^k, y^k + \alpha_k \Delta y^k, s^k + \alpha_k \Delta s^k)$  is feasible.
- ▶ Make step  $(x^{k+1}, y^{k+1}, s^{k+1}) = (x^k + \alpha_k \Delta x^k, y^k + \alpha_k \Delta y^k, s^k + \alpha_k \Delta s^k)$ .

## Path-following algorithm

- ▶ **Short-step path-following method:**  $\mathcal{O}(\sqrt{n})$  complexity result

## Path-following algorithm

- **Short-step path-following method:**  $\mathcal{O}(\sqrt{n})$  complexity result

Theorem ([Gondzio, 2012, Thm. 3.1])

Given  $\epsilon > 0$ , suppose that a feasible starting point  $(x^0, y^0, s^0) \in \mathcal{N}_2(0.1)$  satisfies

$$(x^0)^\top s^0 = n\mu^0, \text{ where } \mu^0 \leq 1/\epsilon^\kappa,$$

for some positive constant  $\kappa$ . Then for some  $K = \mathcal{O}(\sqrt{n} \ln(1/\epsilon))$ , the optimality gap is bounded by  $\epsilon$  after at most  $K$  iterations:

$$\mu^k \leq \epsilon, \quad \forall k \geq K$$

## Path-following algorithm

- **Short-step path-following method:**  $\mathcal{O}(\sqrt{n})$  complexity result

Theorem ([Gondzio, 2012, Thm. 3.1])

Given  $\epsilon > 0$ , suppose that a feasible starting point  $(x^0, y^0, s^0) \in \mathcal{N}_2(0.1)$  satisfies

$$(x^0)^\top s^0 = n\mu^0, \text{ where } \mu^0 \leq 1/\epsilon^\kappa,$$

for some positive constant  $\kappa$ . Then for some  $K = \mathcal{O}(\sqrt{n} \ln(1/\epsilon))$ , the optimality gap is bounded by  $\epsilon$  after at most  $K$  iterations:

$$\mu^k \leq \epsilon, \quad \forall k \geq K$$

- $\theta$ -neighborhood of the central path:

$$\mathcal{N}_2(\theta) := \{(x, y, s) \in \mathcal{F}^0 \mid \|XSe - \mu e\| \leq \theta\mu\}, \text{ with } \mu = \frac{1}{n}x^\top s.$$

## Path-following algorithm

- ▶ **Short-step path-following method:**  $\mathcal{O}(\sqrt{n})$  complexity result

Theorem ([Gondzio, 2012, Thm. 3.1])

Given  $\epsilon > 0$ , suppose that a feasible starting point  $(x^0, y^0, s^0) \in \mathcal{N}_2(0.1)$  satisfies

$$(x^0)^\top s^0 = n\mu^0, \text{ where } \mu^0 \leq 1/\epsilon^\kappa,$$


for some positive constant  $\kappa$ . Then for some  $K = \mathcal{O}(\sqrt{n} \ln(1/\epsilon))$ , the optimality gap is bounded by  $\epsilon$  after at most  $K$  iterations:

$$\mu^k \leq \epsilon, \quad \forall k \geq K$$

- ▶  $\theta$ -neighborhood of the central path:

$\mathcal{N}_2(\theta) := \{(x, y, s) \in \mathcal{F}^0 \mid \|XSe - \mu e\| \leq \theta\mu\}$ , with  $\mu = \frac{1}{n}x^\top s$ .

- ▶ Slow progress towards optimality



## Augmented system

### Newton direction

$$\begin{bmatrix} A & 0 & 0 \\ -Q & A^\top & I \\ S & 0 & X \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta y \\ \Delta s \end{bmatrix} = \begin{bmatrix} b - Ax \\ c - A^\top y - s + Qx \\ \mu_k e - XSe \end{bmatrix} =: \begin{bmatrix} \xi_p \\ \xi_d \\ \xi_\mu \end{bmatrix}$$

use last (complementarity) block to solve for  $\Delta s$  as a function of  $\Delta x$ .

### Augmented system

Define  $\Theta = XS^{-1}$  (ill-conditioned!). Then  $\Delta x$  and  $\Delta y$  solve the Newton system  
 $\iff$

$$\begin{bmatrix} -Q - \Theta^{-1} & A^\top \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta y \end{bmatrix} = \begin{bmatrix} \xi_d - X^{-1}\xi_\mu \\ \xi_p \end{bmatrix}$$

- ▶ Newton system is nonsymmetric.
- ▶ Augmented system is symmetric but indefinite.

## Normal equations

### Augmented system

$$\begin{bmatrix} -\Theta^{-1} & A^\top \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta y \end{bmatrix} = \begin{bmatrix} \xi_d - X^{-1}\xi_\mu \\ \xi_p \end{bmatrix} \equiv: \begin{bmatrix} g \\ \xi_p \end{bmatrix}$$

### Normal equations

Eliminate  $\Delta x$  to arrive at the *Normal equations*

$$(A\Theta A^\top)\Delta y = A\Theta g + \xi_p$$



## Normal equations

### Augmented system

$$\begin{bmatrix} -\Theta^{-1} & A^\top \\ A & 0 \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta y \end{bmatrix} = \begin{bmatrix} \xi_d - X^{-1}\xi_\mu \\ \xi_p \end{bmatrix} =: \begin{bmatrix} g \\ \xi_p \end{bmatrix}$$

### Normal equations

Eliminate  $\Delta x$  to arrive at the *Normal equations*

$$(A\Theta A^\top)\Delta y = A\Theta g + \xi_p$$

- ▶  $A\Theta A^\top$  is symmetric and positive semidefinite. (Finally!)
- ▶ Normal equations in QP ( $A(Q + \Theta)A^\top$ ) $\Delta y = g$  are generally nearly dense, even when  $A$  and  $Q$  are sparse.
- ▶ LP: Normal equations are often used.
- ▶ QP: usually use the indefinite augmented system.

# Outline

IPM for linear and quadratic programs

IPM for convex nonlinear programming

IPM for conic optimization

## IPM for NLP

### ► Convex NLP

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & g(x) \leq 0 \end{array} \iff \begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & g(x) + z = 0, \quad z \geq 0 \end{array}$$

## IPM for NLP

- Convex NLP

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & g(x) \leq 0 \end{array} \iff \begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & g(x) + z = 0, \quad z \geq 0 \end{array}$$

- Replace inequality  $z \geq 0$  with logarithmic barrier

$$\text{minimize} \quad f(x) - \mu \sum_{i=1}^m \ln(z_i) \quad \text{subject to} \quad g(x) + z = 0$$

## IPM for NLP

- Convex NLP

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & g(x) \leq 0 \end{array} \iff \begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & g(x) + z = 0, \quad z \geq 0 \end{array}$$

- Replace inequality  $z \geq 0$  with logarithmic barrier

$$\text{minimize} \quad f(x) - \mu \sum_{i=1}^m \ln(z_i) \quad \text{subject to} \quad g(x) + z = 0$$

- Write out Lagrangian

$$L(x, y, z, \mu) = f(x) + y^\top (g(x) + z) - \mu \sum_{i=1}^m \ln(z_i)$$

## IPM for NLP

- Write conditions for stationary point

$$\nabla_x L(x, z, y) = \nabla f(x) + J_g(x)^\top y = 0$$

$$\nabla_y L(x, z, y) = g(x) + z = 0$$

$$\nabla_z L(x, z, y) = y - \mu Z^{-1} e = 0$$

## IPM for NLP

- ▶ Write conditions for stationary point

$$\nabla_x L(x, z, y) = \nabla f(x) + J_g(x)^\top y = 0$$

$$\nabla_y L(x, z, y) = g(x) + z = 0$$

$$\nabla_z L(x, z, y) = y - \mu Z^{-1} e = 0$$

- ▶ Write KKT system

$$\nabla f(x) + J_g(x)^\top y = 0,$$

$$g(x) + z = 0$$

$$YZe = \mu e$$

## Newton for KKT of NLP

- ▶ Apply Newton method for KKT system



## Newton for KKT of NLP

- ▶ Apply Newton method for KKT system
- ▶ Jacobian matrix of KKT system

$$J_F(x, z, y) = \begin{bmatrix} Q(x, y) & J_g(x)^\top & 0 \\ J_g(x) & 0 & I \\ 0 & Z & Y \end{bmatrix}$$

where  $Q(x, y) = \nabla^2 f(x) + \sum_{i=1}^m y_i \nabla^2 g_i(x)$  is the Hessian of  $L$

## Newton for KKT of NLP

- ▶ Apply Newton method for KKT system
- ▶ Jacobian matrix of KKT system

$$J_F(x, z, y) = \begin{bmatrix} Q(x, y) & J_g(x)^\top & 0 \\ J_g(x) & 0 & I \\ 0 & Z & Y \end{bmatrix}$$

where  $Q(x, y) = \nabla^2 f(x) + \sum_{i=1}^m y_i \nabla^2 g_i(x)$  is the Hessian of  $L$

- ▶ Newton step for KKT system

$$\begin{bmatrix} Q(x, y) & J_g(x)^\top & 0 \\ J_g(x) & 0 & I \\ 0 & Z & Y \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta y \\ \Delta z \end{bmatrix} = \begin{bmatrix} -\nabla f(x) - J_g(x)^\top y \\ -g(x) - z \\ \mu e - YZe \end{bmatrix}$$

## From QP to NLP

- Newton direction for NLP

$$\begin{bmatrix} Q(x, y) & J_g(x)^\top & 0 \\ J_g(x) & 0 & I \\ 0 & Z & Y \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta y \\ \Delta z \end{bmatrix} = \begin{bmatrix} -\nabla f(x) - J_g(x)^\top y \\ -g(x) - z \\ \mu e - YZe \end{bmatrix}$$

## From QP to NLP

- Newton direction for NLP

$$\begin{bmatrix} Q(x, y) & J_g(x)^\top & 0 \\ J_g(x) & 0 & I \\ 0 & Z & Y \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta y \\ \Delta z \end{bmatrix} = \begin{bmatrix} -\nabla f(x) - J_g(x)^\top y \\ -g(x) - z \\ \mu e - YZe \end{bmatrix}$$

- Augmented system for NLP

$$\begin{bmatrix} Q(x, y) & J_g(x)^\top \\ J_g(x) & -ZY^{-1} \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta y \end{bmatrix} = \begin{bmatrix} -\nabla f(x) - J_g(x)^\top y \\ -g(x) - \mu Y^{-1} e \end{bmatrix}$$

## From QP to NLP

- Newton direction for NLP

$$\begin{bmatrix} Q(x, y) & J_g(x)^\top & 0 \\ J_g(x) & 0 & I \\ 0 & Z & Y \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta y \\ \Delta z \end{bmatrix} = \begin{bmatrix} -\nabla f(x) - J_g(x)^\top y \\ -g(x) - z \\ \mu e - YZe \end{bmatrix}$$

- Augmented system for NLP

$$\begin{bmatrix} Q(x, y) & J_g(x)^\top \\ J_g(x) & -ZY^{-1} \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta y \end{bmatrix} = \begin{bmatrix} -\nabla f(x) - J_g(x)^\top y \\ -g(x) - \mu Y^{-1} e \end{bmatrix}$$

- Need to compute  $Q(x, y)$  and  $J_g(x)$  at each iteration

## From QP to NLP

- ▶ Newton direction for NLP

$$\begin{bmatrix} Q(x, y) & J_g(x)^\top & 0 \\ J_g(x) & 0 & I \\ 0 & Z & Y \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta y \\ \Delta z \end{bmatrix} = \begin{bmatrix} -\nabla f(x) - J_g(x)^\top y \\ -g(x) - z \\ \mu e - YZe \end{bmatrix}$$

- ▶ Augmented system for NLP

$$\begin{bmatrix} Q(x, y) & J_g(x)^\top \\ J_g(x) & -ZY^{-1} \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta y \end{bmatrix} = \begin{bmatrix} -\nabla f(x) - J_g(x)^\top y \\ -g(x) - \mu Y^{-1} e \end{bmatrix}$$

- ▶ Need to compute  $Q(x, y)$  and  $J_g(x)$  at each iteration
- ▶ Caveat: use trust region method to choose stepsize as Hessian may be indefinite.

# Outline

IPM for linear and quadratic programs

IPM for convex nonlinear programming

IPM for conic optimization

## Self-concordant function

### Definition

Function  $f$  is *self-concordant* if for some constant  $M_f \geq 0$ , the inequality

$$\nabla^3 f(x)[u, u, u] \leq M_f \|u\|_{\nabla^2 f(x)}^{3/2}$$

holds for any  $x \in \text{dom } f$  and  $u \in R^n$ .

- ▶ A self-concordant function is always well approximated by a quadratic model because the error of such an approximation can be bounded by the  $\|u\|_{\nabla^2 f(x)}^{3/2}$



## Self-concordant function

### Definition

Function  $f$  is *self-concordant* if for some constant  $M_f \geq 0$ , the inequality

$$\nabla^3 f(x)[u, u, u] \leq M_f \|u\|_{\nabla^2 f(x)}^{3/2}$$

holds for any  $x \in \text{dom } f$  and  $u \in R^n$ .

- ▶ A self-concordant function is always well approximated by a quadratic model because the error of such an approximation can be bounded by the  $\|u\|_{\nabla^2 f(x)}^{3/2}$

## Self-concordant function

### Definition

Function  $f$  is *self-concordant* if for some constant  $M_f \geq 0$ , the inequality

$$\nabla^3 f(x)[u, u, u] \leq M_f \|u\|_{\nabla^2 f(x)}^{3/2}$$

holds for any  $x \in \text{dom } f$  and  $u \in R^n$ .

- ▶ A self-concordant function is always well approximated by a quadratic model because the error of such an approximation can be bounded by the  $\|u\|_{\nabla^2 f(x)}^{3/2}$

### Theorem ([Boyd and Vandenberghe, 2004, Section 11.5])

Newton's method with line search finds an  $\varepsilon$  approximate solution in less than  $T := \text{constant} \times (f(x_0) - f^*) + \log_2 \log_2 \frac{1}{\varepsilon}$  iterations.

## Log-barrier is self-concordant

### Theorem

*The barrier function  $-\ln(x)$  is self-concordant in  $\mathbf{R}_+$ .*

## Log-barrier is self-concordant

### Theorem

*The barrier function  $-\ln(x)$  is self-concordant in  $\mathbf{R}_+$ .*

### Proof.

Consider  $f(x) = -\ln(x)$ . This function is self-concordant in  $\mathbf{R}_+$  because

$$f'(x) = -\frac{1}{x}, \quad f''(x) = \frac{1}{x^2}, \quad f'''(x) = -\frac{2}{x^3}$$

## Log-barrier is self-concordant

### Theorem

*The barrier function  $-\ln(x)$  is self-concordant in  $\mathbf{R}_+$ .*

### Proof.

Consider  $f(x) = -\ln(x)$ . This function is self-concordant in  $\mathbf{R}_+$  because

$$f'(x) = -\frac{1}{x}, \quad f''(x) = \frac{1}{x^2}, \quad f'''(x) = -\frac{2}{x^3}$$

Compute and check that self-concordance holds with  $M_f = 2$ . □

►  $-\ln(1/x^\alpha)$ , with  $\alpha \in (0, \infty)$  is not self-concordant in  $\mathbf{R}_+$ .

## Log-barrier is self-concordant

### Theorem

*The barrier function  $-\ln(x)$  is self-concordant in  $\mathbf{R}_+$ .*

### Proof.

Consider  $f(x) = -\ln(x)$ . This function is self-concordant in  $\mathbf{R}_+$  because

$$f'(x) = -\frac{1}{x}, \quad f''(x) = \frac{1}{x^2}, \quad f'''(x) = -\frac{2}{x^3}$$

Compute and check that self-concordance holds with  $M_f = 2$ . □

- ▶  $-\ln(1/x^\alpha)$ , with  $\alpha \in (0, \infty)$  is not self-concordant in  $\mathbf{R}_+$ .
- ▶  $\exp(1/x)$  is not self-concordant in  $\mathbf{R}_+$ .

## Conic optimization

- Consider the optimization problem

$$\begin{array}{ll}\text{minimize} & c^\top x \\ \text{subject to} & Ax = b \\ & x \in K\end{array}$$

where  $K$  is a convex closed cone.

## Conic optimization

- Consider the optimization problem

$$\begin{array}{ll}\text{minimize} & c^\top x \\ \text{subject to} & Ax = b \\ & x \in K\end{array}$$

where  $K$  is a convex closed cone.

- The associated dual is

$$\begin{array}{ll}\text{maximize} & b^\top y \\ \text{subject to} & A^\top y + s = c \\ & s \in K^* \text{ (Dual cone)}\end{array}$$



## Conic optimization

- Consider the optimization problem

$$\begin{array}{ll}\text{minimize} & c^\top x \\ \text{subject to} & Ax = b \\ & x \in K\end{array}$$

where  $K$  is a convex closed cone.

- The associated dual is

$$\begin{array}{ll}\text{maximize} & b^\top y \\ \text{subject to} & A^\top y + s = c \\ & s \in K^* \text{ (Dual cone)}\end{array}$$

- Weak duality

$$c^\top x - b^\top y = x^\top (c - A^\top y) = x^\top s \geq 0$$

## Conic optimization

- Consider the optimization problem

$$\begin{array}{ll}\text{minimize} & c^\top x \\ \text{subject to} & Ax = b \\ & x \in K\end{array}$$

where  $K$  is a convex closed cone.

- The associated dual is

$$\begin{array}{ll}\text{maximize} & b^\top y \\ \text{subject to} & A^\top y + s = c \\ & s \in K^* \text{ (Dual cone)}\end{array}$$

- Weak duality

$$c^\top x - b^\top y = x^\top (c - A^\top y) = x^\top s \geq 0$$

- Conic optimization can be solved in polynomial time with IPMs

## Second-order conic optimization

- ▶  $K = \mathbb{L} := \{(x, t) \mid x \in \mathbf{R}^{n-1}, t \in \mathbf{R}, \|x\|_2 \leq t, t \geq 0\}$  (Second-order cone)

## Second-order conic optimization

- ▶  $K = \mathbb{L} := \{(x, t) \mid x \in \mathbf{R}^{n-1}, t \in \mathbf{R}, \|x\|_2 \leq t, t \geq 0\}$  (Second-order cone)
- ▶ Logarithmic barrier function for the second-order cone

$$f(x, t) = \begin{cases} -\ln(t^2 - \|x\|_2^2) & \text{if } \|x\| < t \\ +\infty & \text{otherwise} \end{cases}$$

## Second-order conic optimization

- ▶  $K = \mathbb{L} := \{(x, t) \mid x \in \mathbf{R}^{n-1}, t \in \mathbf{R}, \|x\|_2 \leq t, t \geq 0\}$  (Second-order cone)
- ▶ Logarithmic barrier function for the second-order cone

$$f(x, t) = \begin{cases} -\ln(t^2 - \|x\|_2^2) & \text{if } \|x\| < t \\ +\infty & \text{otherwise} \end{cases}$$

## Second-order conic optimization

- ▶  $K = \mathbb{L} := \{(x, t) \mid x \in \mathbf{R}^{n-1}, t \in \mathbf{R}, \|x\|_2 \leq t, t \geq 0\}$  (Second-order cone)
- ▶ Logarithmic barrier function for the second-order cone

$$f(x, t) = \begin{cases} -\ln(t^2 - \|x\|_2^2) & \text{if } \|x\| < t \\ +\infty & \text{otherwise} \end{cases}$$

### Theorem

*The barrier function  $f(x, t)$  is self-concordant on  $\mathbb{L}$ .*

## Second-order conic optimization

- ▶  $K = \mathbb{L} := \{(x, t) \mid x \in \mathbf{R}^{n-1}, t \in \mathbf{R}, \|x\|_2 \leq t, t \geq 0\}$  (Second-order cone)
- ▶ Logarithmic barrier function for the second-order cone

$$f(x, t) = \begin{cases} -\ln(t^2 - \|x\|_2^2) & \text{if } \|x\| < t \\ +\infty & \text{otherwise} \end{cases}$$

### Theorem

*The barrier function  $f(x, t)$  is self-concordant on  $\mathbb{L}$ .*

Exercise: Prove in case  $n = 2$ .

## Semidefinite programming

- ▶ Variable now is a symmetric matrix  $X \in K = \mathbf{S}^n$

### SDP and its dual

$$\begin{array}{ll}\text{minimize} & C \bullet X \\ \text{subject to} & A_i \bullet X = b_i, i = 1, \dots, m \\ & X \succeq 0\end{array}$$

$$\begin{array}{ll}\text{maximize} & b^\top y \\ \text{subject to} & \sum_{i=1}^m y_i A_i + S = C \\ & S \succeq 0\end{array}$$

$A_i, C \in \mathbf{S}^n$  and  $b \in \mathbf{R}^m$  given, and  $X, S \in \mathbf{S}^n$  and  $y \in \mathbf{R}^m$  unknown.



## Semidefinite programming

- ▶ Variable now is a symmetric matrix  $X \in K = \mathbf{S}^n$
- ▶ Define  $X \bullet Y = \text{tr}(X^\top Y)$

### SDP and its dual

$$\begin{array}{ll}\text{minimize} & C \bullet X \\ \text{subject to} & A_i \bullet X = b_i, i = 1, \dots, m \\ & X \succeq 0\end{array}$$

$$\begin{array}{ll}\text{maximize} & b^\top y \\ \text{subject to} & \sum_{i=1}^m y_i A_i + S = C \\ & S \succeq 0\end{array}$$

$A_i, C \in \mathbf{S}^n$  and  $b \in \mathbf{R}^m$  given, and  $X, S \in \mathbf{S}^n$  and  $y \in \mathbf{R}^m$  unknown.

## Semidefinite programming

- ▶ Variable now is a symmetric matrix  $X \in K = \mathbf{S}^n$
- ▶ Define  $X \bullet Y = \text{tr}(X^\top Y)$

### SDP and its dual

$$\begin{array}{ll}\text{minimize} & C \bullet X \\ \text{subject to} & A_i \bullet X = b_i, i = 1, \dots, m \\ & X \succeq 0\end{array}$$

$$\begin{array}{ll}\text{maximize} & b^\top y \\ \text{subject to} & \sum_{i=1}^m y_i A_i + S = C \\ & S \succeq 0\end{array}$$

$A_i, C \in \mathbf{S}^n$  and  $b \in \mathbf{R}^m$  given, and  $X, S \in \mathbf{S}^n$  and  $y \in \mathbf{R}^m$  unknown.

## Semidefinite programming

- ▶ Variable now is a symmetric matrix  $X \in K = \mathbf{S}^n$
- ▶ Define  $X \bullet Y = \text{tr}(X^\top Y)$

### SDP and its dual

$$\begin{array}{ll}\text{minimize} & C \bullet X \\ \text{subject to} & A_i \bullet X = b_i, i = 1, \dots, m \\ & X \succeq 0\end{array}$$

$$\begin{array}{ll}\text{maximize} & b^\top y \\ \text{subject to} & \sum_{i=1}^m y_i A_i + S = C \\ & S \succeq 0\end{array}$$

$A_i, C \in \mathbf{S}^n$  and  $b \in \mathbf{R}^m$  given, and  $X, S \in \mathbf{S}^n$  and  $y \in \mathbf{R}^m$  unknown.

### Theorem (Weak duality for SDP)

*If  $X$  is primal feasible and  $(y, S)$  is dual feasible, then*

$$C \bullet X - b^\top y = X \bullet S \geq 0$$

## Logarithmic barrier for SDP

- Logarithmic barrier function for the semi-definite cone

$$f(X) = \begin{cases} -\ln(\det(X)) & \text{if } X \succ 0 \\ +\infty & \text{otherwise} \end{cases}$$

## Logarithmic barrier for SDP

- ▶ Logarithmic barrier function for the semi-definite cone

$$f(X) = \begin{cases} -\ln(\det(X)) & \text{if } X \succ 0 \\ +\infty & \text{otherwise} \end{cases}$$

- ▶ Facts (for small  $t$ ):

## Logarithmic barrier for SDP

- ▶ Logarithmic barrier function for the semi-definite cone

$$f(X) = \begin{cases} -\ln(\det(X)) & \text{if } X \succ 0 \\ +\infty & \text{otherwise} \end{cases}$$

- ▶ Facts (for small  $t$ ):
  - ▶  $\det(I + tU) = 1 + t \operatorname{tr}(U) + \mathcal{O}(t^2)$

## Logarithmic barrier for SDP

- ▶ Logarithmic barrier function for the semi-definite cone

$$f(X) = \begin{cases} -\ln(\det(X)) & \text{if } X \succ 0 \\ +\infty & \text{otherwise} \end{cases}$$

- ▶ Facts (for small  $t$ ):
  - ▶  $\det(I + tU) = 1 + t \operatorname{tr}(U) + \mathcal{O}(t^2)$
  - ▶  $\ln(1 + t \operatorname{tr}(U)) \approx t \operatorname{tr}(U)$

## Logarithmic barrier for SDP

- ▶ Logarithmic barrier function for the semi-definite cone

$$f(X) = \begin{cases} -\ln(\det(X)) & \text{if } X \succ 0 \\ +\infty & \text{otherwise} \end{cases}$$

- ▶ Facts (for small  $t$ ):
  - ▶  $\det(I + tU) = 1 + t \operatorname{tr}(U) + \mathcal{O}(t^2)$
  - ▶  $\ln(1 + t \operatorname{tr}(U)) \approx t \operatorname{tr}(U)$
- ▶ Let  $X \succ 0$  and  $H \in \mathbf{S}^n$  be given. Then

$$\begin{aligned} f(X + tH) &= -\ln(\det(X + tH)) = -\ln(\det(X(I + tX^{-1}H))) \\ &= -\ln(\det(X)) - \ln(\det(I + tX^{-1}H)) \\ &= -\ln(\det(X)) - \ln(1 + t \operatorname{tr}(X^{-1}H) + \mathcal{O}(t^2)) \\ &= f(X) - tX^{-1} \bullet H + \mathcal{O}(t^2) \end{aligned}$$



## Derivatives of Logarithmic barrier for SDP

- First derivative of  $f(X)$

$$f'(X) = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t} = -X^{-1}$$

## Derivatives of Logarithmic barrier for SDP

- First derivative of  $f(X)$

$$f'(X) = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t} = -X^{-1}$$

## Derivatives of Logarithmic barrier for SDP

- First derivative of  $f(X)$

$$f'(X) = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t} = -X^{-1}$$

So  $Df(X)[H] = -X^{-1} \bullet H$ .

## Derivatives of Logarithmic barrier for SDP

- First derivative of  $f(X)$

$$f'(X) = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t} = -X^{-1}$$

So  $Df(X)[H] = -X^{-1} \bullet H$ .

- Second derivative of  $f(X)$

$$\begin{aligned} f'(X + tH) &= -[X(I + tX^{-1}H)]^{-1} = -[I - tX^{-1}H + \mathcal{O}(t^2)]X^{-1} \\ &= f'(X) + tX^{-1}HX^{-1} + \mathcal{O}(t^2) \end{aligned}$$

## Derivatives of Logarithmic barrier for SDP

- First derivative of  $f(X)$

$$f'(X) = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t} = -X^{-1}$$

So  $Df(X)[H] = -X^{-1} \bullet H$ .

- Second derivative of  $f(X)$

$$\begin{aligned} f'(X + tH) &= -[X(I + tX^{-1}H)]^{-1} = -[I - tX^{-1}H + \mathcal{O}(t^2)]X^{-1} \\ &= f'(X) + tX^{-1}HX^{-1} + \mathcal{O}(t^2) \end{aligned}$$

## Derivatives of Logarithmic barrier for SDP

- First derivative of  $f(X)$

$$f'(X) = \lim_{t \rightarrow 0} \frac{f(X + tH) - f(X)}{t} = -X^{-1}$$

So  $Df(X)[H] = -X^{-1} \bullet H$ .

- Second derivative of  $f(X)$

$$\begin{aligned} f'(X + tH) &= -[X(I + tX^{-1}H)]^{-1} = -[I - tX^{-1}H + \mathcal{O}(t^2)]X^{-1} \\ &= f'(X) + tX^{-1}HX^{-1} + \mathcal{O}(t^2) \end{aligned}$$

so  $f''(X)[H] = X^{-1}HX^{-1}$  and  $D^2f(X)[H, G] = X^{-1}HX^{-1} \bullet G$ .

- $f'''(X)[H, G] = -X^{-1}HX^{-1}GX^{-1} - X^{-1}GX^{-1}HX^{-1}$

## Characterization of self-concordance for SDP

### Theorem

*The function  $f(X) = -\ln \det X$  is a convex barrier for  $\mathbf{S}_+^n$ .*

## Characterization of self-concordance for SDP

### Theorem

*The function  $f(X) = -\ln \det X$  is a convex barrier for  $\mathbf{S}_+^n$ .*

### Proof sketch.

Let  $\varphi(t) = F(X + tH)$ . Then, prove that  $\varphi''(t) \geq 0$  for  $t > 0$  such that  $X + tH \succ 0$ . Therefore, when  $X \succ 0$  approaches a singular matrix, its determinant approaches zero, and the function  $f(X) \rightarrow +\infty$ . □



## Characterization of self-concordance for SDP

### Theorem

*The function  $f(X) = -\ln \det X$  is a convex barrier for  $\mathbf{S}_+^n$ .*

### Proof sketch.

Let  $\varphi(t) = F(X + tH)$ . Then, prove that  $\varphi''(t) \geq 0$  for  $t > 0$  such that  $X + tH \succ 0$ . Therefore, when  $X \succ 0$  approaches a singular matrix, its determinant approaches zero, and the function  $f(X) \rightarrow +\infty$ . □

### Theorem ([Nestervov and Nemirovskii, 1994])

*The barrier function  $f(X) = -\ln \det X$  is self-concordant on  $\mathbf{S}_+^n$ .*

## Solving SDPs with IPMs

- Replace the primal SDP

$$\begin{aligned} &\text{minimize} && C \bullet X \\ &\text{subject to} && \mathcal{A}X = b, \\ &&& X \succeq 0, \end{aligned}$$

with the primal barrier SDP

$$\begin{aligned} &\text{minimize} && C \bullet X + \mu f(X) \\ &\text{subject to} && \mathcal{A}X = b, \end{aligned}$$

(with a barrier parameter  $\mu \geq 0$  ).

## Solving SDPs with IPMs

- Replace the primal SDP

$$\begin{aligned} &\text{minimize} && C \bullet X \\ &\text{subject to} && \mathcal{A}X = b, \\ &&& X \succeq 0, \end{aligned}$$

with the primal barrier SDP

$$\begin{aligned} &\text{minimize} && C \bullet X + \mu f(X) \\ &\text{subject to} && \mathcal{A}X = b, \end{aligned}$$

(with a barrier parameter  $\mu \geq 0$  ).

- Formulate the Lagrangian

$$L(X, y, S) = C \bullet X + \mu f(X) - y^T (\mathcal{A}X - b),$$

with  $y \in \mathcal{R}^m$ , and write the first order conditions (FOC) for a stationary point of  $L$ :

$$C + \mu f'(X) - \mathcal{A}^* y = 0$$

## Solving SDPs with IPMs (cont'd)

- ▶ Use  $f(X) = -\ln \det X$  and  $f'(X) = -X^{-1}$  to obtain

$$C - \mu X^{-1} - \mathcal{A}^* y = 0$$

## Solving SDPs with IPMs (cont'd)

- ▶ Use  $f(X) = -\ln \det X$  and  $f'(X) = -X^{-1}$  to obtain

$$C - \mu X^{-1} - \mathcal{A}^* y = 0$$

- ▶ Denote  $S = \mu X^{-1}$ , i.e.,  $XS = \mu I$ . Then, the FOC can be written as

$$\mathcal{A}X = b$$

$$\mathcal{A}^* y + S = C$$

$$XS = \mu I$$

with  $X, S \in \mathbf{S}_{++}^n$ .

## Newton direction

Differentiating this system is hard! The Newton direction solves:

$$\begin{bmatrix} \mathcal{A} & 0 & 0 \\ 0 & \mathcal{A}^* & \mathcal{I} \\ \mu(X^{-1} \odot X^{-1}) & 0 & \mathcal{I} \end{bmatrix} \cdot \begin{bmatrix} \Delta X \\ \Delta y \\ \Delta S \end{bmatrix} = \begin{bmatrix} \xi_b \\ \xi_C \\ \xi_\mu \end{bmatrix}.$$

We define the Kronecker product  $P \odot Q$  for  $P, Q \in \mathbf{R}^{n \times n}$ , which yields a linear operator from  $\mathbf{S}^n$  to  $\mathbf{S}^n$  given by

$$(P \odot Q)U = \frac{1}{2} (PUQ^T + QUP^T).$$

## Summary

- ▶ IPM for SOCP and SDP with self-concordant barrier:

source: [Gondzio, 2012: Interior Point Methods 25 Years Later]

## Summary

- ▶ IPM for SOCP and SDP with self-concordant barrier:
  - ▶ polynomial complexity (predictable behaviour)

source: [Gondzio, 2012: Interior Point Methods 25 Years Later]



## Summary

- ▶ IPM for SOCP and SDP with self-concordant barrier:
  - ▶ polynomial complexity (predictable behaviour)
- ▶ Unified algorithm with fast convergence

source: [Gondzio, 2012: Interior Point Methods 25 Years Later]

## Summary

- ▶ IPM for SOCP and SDP with self-concordant barrier:
  - ▶ polynomial complexity (predictable behaviour)
- ▶ Unified algorithm with fast convergence
  - ▶ from LP via QP to NLP, SOCP and SDP

source: [Gondzio, 2012: Interior Point Methods 25 Years Later]

## Summary

- ▶ IPM for SOCP and SDP with self-concordant barrier:
  - ▶ polynomial complexity (predictable behaviour)
- ▶ Unified algorithm with fast convergence
  - ▶ from LP via QP to NLP, SOCP and SDP
- ▶ efficient for LP, QP, SOCP

source: [Gondzio, 2012: Interior Point Methods 25 Years Later]

## Summary

- ▶ IPM for SOCP and SDP with self-concordant barrier:
  - ▶ polynomial complexity (predictable behaviour)
- ▶ Unified algorithm with fast convergence
  - ▶ from LP via QP to NLP, SOCP and SDP
- ▶ efficient for LP, QP, SOCP
- ▶ problematic for SDP because solving a problem of size  $n$  involves linear algebra operations in dimension  $n^2$

source: [Gondzio, 2012: Interior Point Methods 25 Years Later]

## Summary

- ▶ IPM for SOCP and SDP with self-concordant barrier:
  - ▶ polynomial complexity (predictable behaviour)
- ▶ Unified algorithm with fast convergence
  - ▶ from LP via QP to NLP, SOCP and SDP
- ▶ efficient for LP, QP, SOCP
- ▶ problematic for SDP because solving a problem of size  $n$  involves linear algebra operations in dimension  $n^2$ 
  - ▶ and this requires  $n^6$  flops!

source: [Gondzio, 2012: Interior Point Methods 25 Years Later]