

CME 307 / MS&E 311: Optimization

LP modeling and solution techniques

Professor Udell

Management Science and Engineering
Stanford

January 17, 2024

Course survey

You're interested in

- ▶ duality
- ▶ modeling real-world problems
- ▶ hyperparameter and blackbox optimization
- ▶ fairness and ethics in optimization
- ▶ ...

Outline

LP standard form

Modeling

LP inequality form

Solving LPs

Duality

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ & x \geq 0 \end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $-\infty$

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ & x \geq 0 \end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $-\infty$

Q: if $p^* = -\infty$, does a solution exist? is it unique?
what about $p^* = \infty$?

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ & x \geq 0 \end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $-\infty$

Q: if $p^* = -\infty$, does a solution exist? is it unique?

what about $p^* = \infty$?

henceforth assume $A \in \mathbf{R}^{m \times n}$ has full row rank m

Q: why? how to check?

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ & x \geq 0 \end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $-\infty$

Q: if $p^* = -\infty$, does a solution exist? is it unique?

what about $p^* = \infty$?

henceforth assume $A \in \mathbf{R}^{m \times n}$ has full row rank m

Q: why? how to check?

A: otherwise infeasible or redundant rows; use gaussian elimination to check and remove

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

► span of A :

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$
- ▶ nullspace of A :

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$
- ▶ nullspace of A : $\text{nullspace}(A) = \{x \in \mathbf{R}^n \mid Ax = 0\} \subseteq \mathbf{R}^n$

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$
- ▶ nullspace of A : $\text{nullspace}(A) = \{x \in \mathbf{R}^n \mid Ax = 0\} \subseteq \mathbf{R}^n$
- ▶ how to compute basis for span and nullspace of A ?

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$
- ▶ nullspace of A : $\text{nullspace}(A) = \{x \in \mathbf{R}^n \mid Ax = 0\} \subseteq \mathbf{R}^n$
- ▶ how to compute basis for span and nullspace of A ?
can use QR factorization or SVD

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$
- ▶ nullspace of A : $\text{nullspace}(A) = \{x \in \mathbf{R}^n \mid Ax = 0\} \subseteq \mathbf{R}^n$
- ▶ how to compute basis for span and nullspace of A ?
can use QR factorization or SVD
- ▶ how to solve $Ax = b$?

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$
- ▶ nullspace of A : $\text{nullspace}(A) = \{x \in \mathbf{R}^n \mid Ax = 0\} \subseteq \mathbf{R}^n$
- ▶ how to compute basis for span and nullspace of A ?
can use QR factorization or SVD
- ▶ how to solve $Ax = b$? factor-solve with QR or SVD; form normal equations $A^T Ax = A^T b$ and use CG; other Krylov methods like LSQR (positive definite), MINRES (indefinite), GMRES (general)

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$
- ▶ nullspace of A : $\text{nullspace}(A) = \{x \in \mathbf{R}^n \mid Ax = 0\} \subseteq \mathbf{R}^n$
- ▶ how to compute basis for span and nullspace of A ?
can use QR factorization or SVD
- ▶ how to solve $Ax = b$? factor-solve with QR or SVD; form normal equations $A^T Ax = A^T b$ and use CG; other Krylov methods like LSQR (positive definite), MINRES (indefinite), GMRES (general)
- ▶ solution to $Ax = b$ is unique if

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$
- ▶ nullspace of A : $\text{nullspace}(A) = \{x \in \mathbf{R}^n \mid Ax = 0\} \subseteq \mathbf{R}^n$
- ▶ how to compute basis for span and nullspace of A ?
can use QR factorization or SVD
- ▶ how to solve $Ax = b$? factor-solve with QR or SVD; form normal equations $A^T Ax = A^T b$ and use CG; other Krylov methods like LSQR (positive definite), MINRES (indefinite), GMRES (general)
- ▶ solution to $Ax = b$ is unique if $m = n$ and A is full rank

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$
- ▶ nullspace of A : $\text{nullspace}(A) = \{x \in \mathbf{R}^n \mid Ax = 0\} \subseteq \mathbf{R}^n$
- ▶ how to compute basis for span and nullspace of A ?
can use QR factorization or SVD
- ▶ how to solve $Ax = b$? factor-solve with QR or SVD; form normal equations $A^T Ax = A^T b$ and use CG; other Krylov methods like LSQR (positive definite), MINRES (indefinite), GMRES (general)
- ▶ solution to $Ax = b$ is unique if $m = n$ and A is full rank
- ▶ if $m < n$ and A is full rank
 - ▶ solution set is a hyperplane of dimension

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$
- ▶ nullspace of A : $\text{nullspace}(A) = \{x \in \mathbf{R}^n \mid Ax = 0\} \subseteq \mathbf{R}^n$
- ▶ how to compute basis for span and nullspace of A ?
can use QR factorization or SVD
- ▶ how to solve $Ax = b$? factor-solve with QR or SVD; form normal equations $A^T Ax = A^T b$ and use CG; other Krylov methods like LSQR (positive definite), MINRES (indefinite), GMRES (general)
- ▶ solution to $Ax = b$ is unique if $m = n$ and A is full rank
- ▶ if $m < n$ and A is full rank
 - ▶ solution set is a hyperplane of dimension $n - m$
 - ▶ null space of A , $\text{nullspace}(A)$, is a hyperplane of dimension

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$
- ▶ nullspace of A : $\text{nullspace}(A) = \{x \in \mathbf{R}^n \mid Ax = 0\} \subseteq \mathbf{R}^n$
- ▶ how to compute basis for span and nullspace of A ?
can use QR factorization or SVD
- ▶ how to solve $Ax = b$? factor-solve with QR or SVD; form normal equations $A^T Ax = A^T b$ and use CG; other Krylov methods like LSQR (positive definite), MINRES (indefinite), GMRES (general)
- ▶ solution to $Ax = b$ is unique if $m = n$ and A is full rank
- ▶ if $m < n$ and A is full rank
 - ▶ solution set is a hyperplane of dimension $n - m$
 - ▶ null space of A , $\text{nullspace}(A)$, is a hyperplane of dimension $n - m$
 - ▶ solution set is $\{x : Ax = b\} = \{x_0 + Vz\}$ where columns of $V \in \mathbf{R}^{n \times n-m}$ span $\text{nullspace}(A)$

Linear algebra review

matrix $A \in \mathbf{R}^{m \times n}$

- ▶ span of A : $\text{span}(A) = \{Ax \mid x \in \mathbf{R}^n\} \subseteq \mathbf{R}^m$
- ▶ nullspace of A : $\text{nullspace}(A) = \{x \in \mathbf{R}^n \mid Ax = 0\} \subseteq \mathbf{R}^n$
- ▶ how to compute basis for span and nullspace of A ?
can use QR factorization or SVD
- ▶ how to solve $Ax = b$? factor-solve with QR or SVD; form normal equations $A^T Ax = A^T b$ and use CG; other Krylov methods like LSQR (positive definite), MINRES (indefinite), GMRES (general)
- ▶ solution to $Ax = b$ is unique if $m = n$ and A is full rank
- ▶ if $m < n$ and A is full rank
 - ▶ solution set is a hyperplane of dimension $n - m$
 - ▶ null space of A , $\text{nullspace}(A)$, is a hyperplane of dimension $n - m$
 - ▶ solution set is $\{x : Ax = b\} = \{x_0 + Vz\}$ where columns of $V \in \mathbf{R}^{n \times n-m}$ span $\text{nullspace}(A)$

if these are confusing: review linear algebra and prove them all!

LP example: diet problem

- ▶ x_i servings of food i
- ▶ c_i cost per serving
- ▶ a_{ij} amount of nutrient j in food i
- ▶ b_j required amount of nutrient j

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

LP example: diet problem

- ▶ x_i servings of food i
- ▶ c_i cost per serving
- ▶ a_{ij} amount of nutrient j in food i
- ▶ b_j required amount of nutrient j

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

extensions:

- ▶ foods come from recipes? $x = By$

LP example: diet problem

- ▶ x_i servings of food i
- ▶ c_i cost per serving
- ▶ a_{ij} amount of nutrient j in food i
- ▶ b_j required amount of nutrient j

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

extensions:

- ▶ foods come from recipes?
- ▶ ensure diversity in diet?

LP example: diet problem

- ▶ x_i servings of food i
- ▶ c_i cost per serving
- ▶ a_{ij} amount of nutrient j in food i
- ▶ b_j required amount of nutrient j

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

extensions:

- ▶ foods come from recipes?
- ▶ ensure diversity in diet? $y \leq u$

LP example: diet problem

- ▶ x_i servings of food i
- ▶ c_i cost per serving
- ▶ a_{ij} amount of nutrient j in food i
- ▶ b_j required amount of nutrient j

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

extensions:

- ▶ foods come from recipes?
- ▶ ensure diversity in diet? $y \leq u$
- ▶ ranges of nutrients?

LP example: diet problem

- ▶ x_i servings of food i
- ▶ c_i cost per serving
- ▶ a_{ij} amount of nutrient j in food i
- ▶ b_j required amount of nutrient j

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

extensions:

- ▶ foods come from recipes?
- ▶ ensure diversity in diet? $y \leq u$
- ▶ ranges of nutrients? $l \leq y \leq u$

Geometry of LP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

the **feasible set** is the set of points x that satisfy all constraints

- ▶ interpretation: add up columns of A so they match b
- ▶ $Ax = b$ defines a **hyperplane**
- ▶ $x_i \geq 0$ is a **halfspace**
- ▶ $x \geq 0$ is the **positive orthant**

Geometry of LP: convexity

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

- ▶ define the **feasible set** $\{x : Ax = b, x \geq 0\}$

Geometry of LP: convexity

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

- ▶ define the **feasible set** $\{x : Ax = b, x \geq 0\}$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

Geometry of LP: convexity

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

- ▶ define the **feasible set** $\{x : Ax = b, x \geq 0\}$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ fact: the feasible set is convex

Geometry of LP: convexity

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

- ▶ define the **feasible set** $\{x : Ax = b, x \geq 0\}$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ fact: the feasible set is convex
- ▶ define **extreme point**: x is extreme in C if it cannot be written as a linear combination of other points in C :

$$x \in C \quad \text{and} \quad x = \theta y + (1 - \theta)z \quad \implies \quad x = y = z$$

Geometry of LP: convexity

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

- ▶ define the **feasible set** $\{x : Ax = b, x \geq 0\}$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ fact: the feasible set is convex
- ▶ define **extreme point**: x is extreme in C if it cannot be written as a linear combination of other points in C :

$$x \in C \quad \text{and} \quad x = \theta y + (1 - \theta)z \quad \implies \quad x = y = z$$

- ▶ fact: if a solution exists, then some extreme point of the feasible set is optimal

Geometry of LP: polytopes

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

- ▶ define **polytope** P : convex hull of its extreme points $v_1, \dots, v_k \in \mathbf{R}^n$:

$$P = \{x \in \mathbf{R}^n \mid x = \sum_{i=1}^k \theta_i v_i, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1\}$$

- ▶ if feasible set is bounded, it is a polytope
- ▶ prove: if a solution exists, then some extreme point of the feasible set is optimal

Outline

LP standard form

Modeling

LP inequality form

Solving LPs

Duality

Let's do some modeling!

- ▶ OptiMUS: <https://optimus-solver.vercel.app/>
- ▶ power systems: https://jump.dev/JuMP.jl/stable/tutorials/applications/power_systems/
- ▶ multicast routing:
<https://colab.research.google.com/drive/1iOn1T1Muh51KaA7mf7UIQOdhSFZhZyry?usp=sharing>

Let's do some modeling!

- ▶ OptiMUS: <https://optimus-solver.vercel.app/>
- ▶ power systems: https://jump.dev/JuMP.jl/stable/tutorials/applications/power_systems/
- ▶ multicast routing:
<https://colab.research.google.com/drive/1iOn1T1Muh51KaA7mf7UIQOdhSFZhZyry?usp=sharing>

practical solvers for MILP:

- ▶ Gurobi and COPT (cardinal optimizer) are the state-of-the-art commercial solvers
- ▶ GLPK is a free solver that is not as fast
- ▶ JuliaOpt/JuMP is a modeling language in Julia that calls solvers like Gurobi and is specialized for MILP applications
- ▶ CVX* (including CVXPY in python) are modeling languages that call solvers like Gurobi with good support for convex problems
- ▶ OptiMUS is a LLM-based modeling tool for MILP

Modeling challenges

model the following as standard form LPs:

- ▶ **inequality constraints.** $Ax \leq b$
- ▶ **free variable.** $x \in \mathbf{R}$
- ▶ **absolute value.** constraint $|x| \leq 10$
- ▶ **piecewise linear.** objective $\max(x_1, x_2)$
- ▶ **assignment.** e.g., every class is assigned exactly one classroom
- ▶ **logic.** e.g., class enrollment \leq capacity of assigned room
- ▶ **flow.** e.g., the least cost way to ship an item from s to t

Outline

LP standard form

Modeling

LP inequality form

Solving LPs

Duality

LP inequality form

another common form for LP is **inequality form**

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

how to transform to standard form?

- ▶ inequality constraints $Ax \leq b$?

LP inequality form

another common form for LP is **inequality form**

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

how to transform to standard form?

- ▶ inequality constraints $Ax \leq b$? slack variables $s \geq 0$
- ▶ free variable $x \in \mathbf{R}^n$?

LP inequality form

another common form for LP is **inequality form**

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

how to transform to standard form?

- ▶ inequality constraints $Ax \leq b$? slack variables $s \geq 0$
- ▶ free variable $x \in \mathbf{R}^n$? split into positive and negative parts

we will see later that these forms are also related by **duality**

LP example: production planning

- ▶ x_i units of product i
- ▶ c_i cost per unit
- ▶ a_{ij} amount of resource j used by product i
- ▶ b_j amount of resource j available
- ▶ d_i demand for product i
- ▶ u_i max production of product i
- ▶ l_i min production of product i

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & l \leq x \leq u\end{array}$$

LP example: production planning

- ▶ x_i units of product i
- ▶ c_i cost per unit
- ▶ a_{ij} amount of resource j used by product i
- ▶ b_j amount of resource j available
- ▶ d_i demand for product i
- ▶ u_i max production of product i
- ▶ l_i min production of product i

extensions:

- ▶ fixed cost for producing product i at all?

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & l \leq x \leq u\end{array}$$

LP example: production planning

- ▶ x_i units of product i
- ▶ c_i cost per unit
- ▶ a_{ij} amount of resource j used by product i
- ▶ b_j amount of resource j available
- ▶ d_i demand for product i
- ▶ u_i max production of product i
- ▶ l_i min production of product i

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & l \leq x \leq u\end{array}$$

extensions:

- ▶ fixed cost for producing product i at all?
 $c^T x + f^T z, z_i \in \{0, 1\}, x_i \leq Mz_i$ for M large

Geometry of LP: inequality form

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

- ▶ $Ax \leq b$ defines a **polyhedron**

Geometry of LP: inequality form

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

- ▶ $Ax \leq b$ defines a **polyhedron**
- ▶ \implies feasible set $P = \{x : Ax \leq b\}$ is a polyhedron

Geometry of LP: inequality form

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

- ▶ $Ax \leq b$ defines a **polyhedron**
- ▶ \implies feasible set $P = \{x : Ax \leq b\}$ is a polyhedron
- ▶ x is a **vertex** of polyhedron P if there is some c so that

$$c^T x < c^T y, \quad \forall y \in P \setminus \{x\}$$

Geometry of LP: inequality form

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

- ▶ $Ax \leq b$ defines a **polyhedron**
- ▶ \implies feasible set $P = \{x : Ax \leq b\}$ is a polyhedron
- ▶ x is a **vertex** of polyhedron P if there is some c so that

$$c^T x < c^T y, \quad \forall y \in P \setminus \{x\}$$

fact: vertex \iff extreme point

Solution of LP is extreme point

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

fact: if a solution exists and the feasible set has an extreme point, then some extreme point of the feasible set is optimal

Solution of LP is extreme point

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

fact: if a solution exists and the feasible set has an extreme point, then some extreme point of the feasible set is optimal

cases: solution x^* is unique / not unique

Solution of LP is extreme point

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

fact: if a solution exists and the feasible set has an extreme point, then some extreme point of the feasible set is optimal

cases: solution x^* is unique / not unique

- ▶ unique: so $c^T x < c^T y$ for all $y \in P \setminus \{x\}$

Solution of LP is extreme point

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

fact: if a solution exists and the feasible set has an extreme point, then some extreme point of the feasible set is optimal

cases: solution x^* is unique / not unique

- ▶ unique: so $c^T x < c^T y$ for all $y \in P \setminus \{x\}$
- ▶ not unique: $\{X^* : c^T x = c^T x^*, x \in P\}$ is a polyhedron. It is not empty (a solution exists) and its complement is not empty (optimal value is bounded). So, it has at least one vertex. That vertex is also a vertex of P .

Basic feasible solution

define: $x \in \mathbf{R}^n$ is a **basic feasible solution** (BFS) if there is a set S of m linearly independent active constraints so that

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0.$$

- ▶ $A_S \in \mathbf{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ BFS \iff extreme point
- ▶ two BFS with S, S' are neighbors if they share $m - 1$ constraints: $|S \cap S'| = m - 1$

Basic feasible solution

define: $x \in \mathbf{R}^n$ is a **basic feasible solution** (BFS) if there is a set S of m linearly independent active constraints so that

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0.$$

- ▶ $A_S \in \mathbf{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ BFS \iff extreme point
- ▶ two BFS with S, S' are neighbors if they share $m - 1$ constraints: $|S \cap S'| = m - 1$

define: the **active set** is the set of constraints that hold with equality

Basic feasible solution

define: $x \in \mathbf{R}^n$ is a **basic feasible solution** (BFS) if there is a set S of m linearly independent active constraints so that

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0.$$

- ▶ $A_S \in \mathbf{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ BFS \iff extreme point
- ▶ two BFS with S, S' are neighbors if they share $m - 1$ constraints: $|S \cap S'| = m - 1$

define: the **active set** is the set of constraints that hold with equality

Q: how to find a BFS?

Basic feasible solution

define: $x \in \mathbf{R}^n$ is a **basic feasible solution** (BFS) if there is a set S of m linearly independent active constraints so that

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0.$$

- ▶ $A_S \in \mathbf{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ BFS \iff extreme point
- ▶ two BFS with S, S' are neighbors if they share $m - 1$ constraints: $|S \cap S'| = m - 1$

define: the **active set** is the set of constraints that hold with equality

Q: how to find a BFS?

A: start at a feasible point; move in a **feasible direction** until you hit another constraint; continue until you reach a BFS

Outline

LP standard form

Modeling

LP inequality form

Solving LPs

Duality

Solving LPs

algorithms:

- ▶ enumerate all vertices and check
- ▶ fourier-motzkin elimination
- ▶ simplex method
- ▶ ellipsoid method
- ▶ interior point methods
- ▶ first-order methods
- ▶ ...

Solving LPs

algorithms:

- ▶ enumerate all vertices and check
- ▶ fourier-motzkin elimination
- ▶ simplex method
- ▶ ellipsoid method
- ▶ interior point methods
- ▶ first-order methods
- ▶ ...

remarks:

- ▶ enumeration and elimination are simple but not practical
- ▶ simplex was the first practical algorithm; still used today
- ▶ ellipsoid method is the first polynomial-time algorithm; not practical
- ▶ interior point methods are polynomial-time and practical
- ▶ first-order methods are practical and scale to large problems

Discuss: how to solve LPs?

write down a method to solve LPs; discuss in groups.

- ▶ idea
- ▶ math
- ▶ pseudocode

complete <https://forms.gle/JbP2fLd6cRVbNUoW9> when you're ready (and before Friday noon)
(link also available from course schedule)

Enumerate vertices of LP

can generate all extreme points of LP: for each $S \subseteq \{1, \dots, n\}$ with $|S| = m$,

- ▶ $A_S \in \mathbf{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ solve $A_S x_S = b$ for x_S and set $x_{\bar{S}} = 0$
- ▶ if $x_S \geq 0$, then x is a BFS
- ▶ evaluate objective $c^T x$

the best BFS is optimal!

Enumerate vertices of LP

can generate all extreme points of LP: for each $S \subseteq \{1, \dots, n\}$ with $|S| = m$,

- ▶ $A_S \in \mathbf{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ solve $A_S x_S = b$ for x_S and set $x_{\bar{S}} = 0$
- ▶ if $x_S \geq 0$, then x is a BFS
- ▶ evaluate objective $c^T x$

the best BFS is optimal!

problem: how many BFSs are there?

Enumerate vertices of LP

can generate all extreme points of LP: for each $S \subseteq \{1, \dots, n\}$ with $|S| = m$,

- ▶ $A_S \in \mathbf{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ solve $A_S x_S = b$ for x_S and set $x_{\bar{S}} = 0$
- ▶ if $x_S \geq 0$, then x is a BFS
- ▶ evaluate objective $c^T x$

the best BFS is optimal!

problem: how many BFSs are there?

n choose m is $\binom{n}{m} = \frac{n!}{m!(n-m)!}$ (“exponentially many”)

Simplex algorithm

basic idea: local search on the vertices of the feasible set

- ▶ start at BFS x and evaluate objective $c^T x$
- ▶ move to a neighboring BFS x' with better objective $c^T x'$
- ▶ repeat until no improvement possible

Simplex algorithm

basic idea: local search on the vertices of the feasible set

- ▶ start at BFS x and evaluate objective $c^T x$
- ▶ move to a neighboring BFS x' with better objective $c^T x'$
- ▶ repeat until no improvement possible

discuss in groups:

- ▶ how to find an initial BFS?
- ▶ how to find a neighboring BFS with better objective?
- ▶ how to prove optimality?

Finding an initial BFS

solve an auxiliary problem for which a BFS is known:

$$\begin{array}{ll}\text{minimize} & \sum_{i=1}^n z_i \\ \text{subject to} & Ax + Dz = b \\ & x, z \geq 0\end{array}$$

where $D \in \mathbf{R}^{m \times m}$ is a diagonal matrix with $D_{ii} = \mathbf{sign}(b_i)$ for $i = 1, \dots, m$.

- ▶ $x = 0, z = b$ is a BFS of this problem
- ▶ $(x, z) = (x, 0)$ is a BFS of this problem $\iff x$ is a BFS of the original problem

Find a better neighboring BFS

start with BFS x with active set S and turn on variable $j \notin S$

$$x^+ \leftarrow x + \theta d, \quad \theta > 0$$

where $d_j = 1$ and $d_i = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S .

Find a better neighboring BFS

start with BFS x with active set S and turn on variable $j \notin S$

$$x^+ \leftarrow x + \theta d, \quad \theta > 0$$

where $d_j = 1$ and $d_i = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S .

- need to stay feasible wrt equality constraints, so

$$Ax = b, \quad A(x + \theta d) = b, \implies Ad = 0$$

Find a better neighboring BFS

start with BFS x with active set S and turn on variable $j \notin S$

$$x^+ \leftarrow x + \theta d, \quad \theta > 0$$

where $d_j = 1$ and $d_i = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S .

- ▶ need to stay feasible wrt equality constraints, so

$$Ax = b, \quad A(x + \theta d) = b, \implies Ad = 0$$

- ▶ construct the **j th basic direction**

$$Ad = A_S d_S + A_j = 0 \implies d_S = -A_S^{-1} A_j$$

Find a better neighboring BFS

start with BFS x with active set S and turn on variable $j \notin S$

$$x^+ \leftarrow x + \theta d, \quad \theta > 0$$

where $d_j = 1$ and $d_i = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S .

- ▶ need to stay feasible wrt equality constraints, so

$$Ax = b, \quad A(x + \theta d) = b, \implies Ad = 0$$

- ▶ construct the **j th basic direction**

$$Ad = A_S d_S + A_j = 0 \implies d_S = -A_S^{-1} A_j$$

- ▶ if $x_S > 0$ is **non-degenerate**, then $\exists \theta > 0$ st $x^+ \geq 0$

Find a better neighboring BFS

start with BFS x with active set S and turn on variable $j \notin S$

$$x^+ \leftarrow x + \theta d, \quad \theta > 0$$

where $d_j = 1$ and $d_i = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S .

- ▶ need to stay feasible wrt equality constraints, so

$$Ax = b, \quad A(x + \theta d) = b, \implies Ad = 0$$

- ▶ construct the **j th basic direction**

$$Ad = A_S d_S + A_j = 0 \implies d_S = -A_S^{-1} A_j$$

- ▶ if $x_S > 0$ is **non-degenerate**, then $\exists \theta > 0$ st $x^+ \geq 0$
- ▶ how does objective change?

$$c^T x^+ = c^T x + \theta c_j^T d = c^T x + c_j - \theta c_S^T A_S^{-1} A_j$$

Reduced cost

define **reduced cost** $\bar{c}_j = c_j - c_S^T A_S^{-1} A_j, j \notin S$

Reduced cost

define **reduced cost** $\bar{c}_j = c_j - c_S^T A_S^{-1} A_j, j \notin S$

fact:

- ▶ if $\bar{c} \geq 0$, x is optimal
- ▶ if x is optimal and nondegenerate ($x_S > 0$), then $\bar{c} \geq 0$

Outline

LP standard form

Modeling

LP inequality form

Solving LPs

Duality

Why duality?

- ▶ certify optimality
 - ▶ turn \forall into \exists
 - ▶ use dual lower bound to derive stopping conditions
- ▶ new algorithms based on the dual
 - ▶ solve dual, then recover primal solution

Warmup: Farkas lemma

Theorem (Farkas lemma)

Given $A \in \mathbf{R}^{m \times n}$ and $b \in \mathbf{R}^m$, exactly one of the following is true:

- ▶ *there exists $x \in \mathbf{R}^n$ so that $Ax = b$ and $x \geq 0$*
- ▶ *there exists $y \in \mathbf{R}^m$ so that $A^T y \geq 0$ and $\langle b, y \rangle < 0$*

\implies can efficiently certify infeasibility of a linear program

Warmup: Farkas lemma

Theorem (Farkas lemma)

Given $A \in \mathbf{R}^{m \times n}$ and $b \in \mathbf{R}^m$, exactly one of the following is true:

- ▶ there exists $x \in \mathbf{R}^n$ so that $Ax = b$ and $x \geq 0$
- ▶ there exists $y \in \mathbf{R}^m$ so that $A^T y \geq 0$ and $\langle b, y \rangle < 0$

\implies can efficiently certify infeasibility of a linear program

proof: suppose we have $x \in \mathbf{R}^n$ so that $Ax = b$ and $x \geq 0$.
then for any $y \in \mathbf{R}^m$,

$$\begin{aligned} 0 &= \langle y, b - Ax \rangle = \langle y, b \rangle - \langle A^T y, x \rangle \\ \langle y, b \rangle &= \langle A^T y, x \rangle \end{aligned}$$

so if $A^T y \geq 0$, then use $x \geq 0$ to conclude $\langle y, b \rangle \geq 0$.

Warmup: Farkas lemma

Theorem (Farkas lemma)

Given $A \in \mathbf{R}^{m \times n}$ and $b \in \mathbf{R}^m$, exactly one of the following is true:

- ▶ there exists $x \in \mathbf{R}^n$ so that $Ax = b$ and $x \geq 0$
- ▶ there exists $y \in \mathbf{R}^m$ so that $A^T y \geq 0$ and $\langle b, y \rangle < 0$

\implies can efficiently certify infeasibility of a linear program

proof: suppose we have $x \in \mathbf{R}^n$ so that $Ax = b$ and $x \geq 0$.
then for any $y \in \mathbf{R}^m$,

$$\begin{aligned} 0 &= \langle y, b - Ax \rangle = \langle y, b \rangle - \langle A^T y, x \rangle \\ \langle y, b \rangle &= \langle A^T y, x \rangle \end{aligned}$$

so if $A^T y \geq 0$, then use $x \geq 0$ to conclude $\langle y, b \rangle \geq 0$.

(opposite direction is similar)

Lagrange duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ & x \geq 0 \end{array} \quad (\mathcal{P})$$

if x is feasible, then $Ax = b$, so $\langle y, Ax - b \rangle = 0$ for $y \in \mathbf{R}^m$.

Lagrange duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ & x \geq 0 \end{array} \quad (\mathcal{P})$$

if x is feasible, then $Ax = b$, so $\langle y, Ax - b \rangle = 0$ for $y \in \mathbf{R}^m$.

define the **Lagrangian**

$$\mathcal{L}(x, y) := c^T x - \langle y, Ax - b \rangle$$

Lagrange duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ & x \geq 0 \end{array} \quad (\mathcal{P})$$

if x is feasible, then $Ax = b$, so $\langle y, Ax - b \rangle = 0$ for $y \in \mathbf{R}^m$.

define the **Lagrangian**

$$\begin{aligned} \mathcal{L}(x, y) &:= c^T x - \langle y, Ax - b \rangle \\ p^* &= \inf_{x: Ax=b, x \geq 0} \mathcal{L}(x, y) \geq \inf_{x \geq 0} \mathcal{L}(x, y) \end{aligned}$$

Lagrange duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ & x \geq 0 \end{array} \quad (\mathcal{P})$$

if x is feasible, then $Ax = b$, so $\langle y, Ax - b \rangle = 0$ for $y \in \mathbf{R}^m$.

define the **Lagrangian**

$$\begin{aligned} \mathcal{L}(x, y) &:= c^T x - \langle y, Ax - b \rangle \\ p^* &= \inf_{x: Ax=b, x \geq 0} \mathcal{L}(x, y) \geq \inf_{x \geq 0} \mathcal{L}(x, y) \\ &= \inf_{x \geq 0} c^T x + \langle y, b - Ax \rangle \\ &= \langle y, b \rangle + \inf_{x \geq 0} \left(c^T x - \langle A^T y, x \rangle \right) \\ &= \langle y, b \rangle + \inf_{x \geq 0} \left(\langle c - A^T y, x \rangle \right) \end{aligned}$$

unbounded below unless $c - A^T y \geq 0$.

Lagrange duality, ctd

we have a lower bound on p^* for any y , and a useful one whenever $c + A^T y = 0$. maximize bound:

$$p^* \geq \begin{array}{ll} \text{maximize} & \langle y, b \rangle \\ \text{subject to} & A^T y \leq c \\ \text{variable} & y \in \mathbf{R}^m \end{array}$$

define the **dual function**

$$g(y) = \begin{cases} \langle y, b \rangle & A^T y \leq c \\ -\infty & \text{otherwise} \end{cases}$$

Lagrange duality

weak duality asserts that $p^* \geq g(y)$ for all $y \in \mathbf{R}^m$.

$$\begin{aligned} p^* &\geq g(y) \quad \forall y \in \mathbf{R}^m \\ &\geq \underbrace{\sup_y g(y)}_{\mathcal{D}} =: d^* \end{aligned}$$

$p^* \geq d^*$ dual optimal value

Strong duality

Definition (Duality gap)

The **duality gap** for a primal-dual pair (x, y) is $c^T x - b^T y \geq 0$

by weak duality, duality gap is always nonnegative

Strong duality

Definition (Duality gap)

The **duality gap** for a primal-dual pair (x, y) is $c^T x - b^T y \geq 0$

by weak duality, duality gap is always nonnegative

Definition (Strong duality)

A primal-dual pair (x^*, y^*) satisfies **strong duality** if

$$p^* = d^* \iff c^T x - b^T y = 0$$

Strong duality

Definition (Duality gap)

The **duality gap** for a primal-dual pair (x, y) is $c^T x - b^T y \geq 0$

by weak duality, duality gap is always nonnegative

Definition (Strong duality)

A primal-dual pair (x^*, y^*) satisfies **strong duality** if

$$p^* = d^* \iff c^T x - b^T y = 0$$

strong duality holds

- ▶ for feasible LPs
- ▶ (later) for convex problems under **constraint qualification** aka **Slater's condition**. feasible region has an **interior point** x so that all inequality constraints hold strictly

strong duality fails if either primal or dual problem is infeasible or unbounded

Strong duality for LPs

primal and dual LP in standard form:

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

$$\begin{array}{ll}\text{maximize} & b^T y \\ \text{subject to} & A^T y \leq c\end{array}$$

claim: if primal LP has a bounded feasible solution x^* , then strong duality holds

i.e., dual LP has a bounded feasible solution y^* and $p^* = d^*$

Proof of strong duality for LPs

consider the following system with variables $x' \in \mathbf{R}^n$, $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

Proof of strong duality for LPs

consider the following system with variables $x' \in \mathbf{R}^n$, $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

claim: this system has no solution. pf by contradiction:

- ▶ if $\tau > 0$, then x'/τ is feasible for LP and $c^T x'/\tau < p^*$
- ▶ if $\tau = 0$, then $x^* + x'$ is feasible for LP and $c^T(x^* + x') < p^*$

Proof of strong duality for LPs

consider the following system with variables $x' \in \mathbf{R}^n$, $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

claim: this system has no solution. pf by contradiction:

- ▶ if $\tau > 0$, then x'/τ is feasible for LP and $c^T x'/\tau < p^*$
- ▶ if $\tau = 0$, then $x^* + x'$ is feasible for LP and $c^T(x^* + x') < p^*$

so use Farkas' lemma:

$$Ax + b = 0, \quad x \geq 0 \quad \text{or} \quad A^T y \geq 0, \quad b^T y < 0$$

Proof of strong duality for LPs

consider the following system with variables $x' \in \mathbf{R}^n$, $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

claim: this system has no solution. pf by contradiction:

- ▶ if $\tau > 0$, then x'/τ is feasible for LP and $c^T x'/\tau < p^*$
- ▶ if $\tau = 0$, then $x^* + x'$ is feasible for LP and $c^T(x^* + x') < p^*$

so use Farkas' lemma:

$$\begin{aligned} Ax + b = 0, \quad x \geq 0 \quad & \text{or} \quad A^T y \geq 0, \quad b^T y < 0 \\ \begin{bmatrix} A & -b \\ c^T & -p^* \end{bmatrix} \begin{bmatrix} x \\ \tau \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \end{bmatrix} \quad & \text{or} \quad \begin{bmatrix} A^T & c \\ -b^T & -p^* \end{bmatrix} \begin{bmatrix} y \\ \sigma \end{bmatrix} \geq 0, \quad \sigma > 0 \end{aligned}$$

Proof of strong duality for LPs

consider the following system with variables $x' \in \mathbf{R}^n$, $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

claim: this system has no solution. pf by contradiction:

- ▶ if $\tau > 0$, then x'/τ is feasible for LP and $c^T x'/\tau < p^*$
- ▶ if $\tau = 0$, then $x^* + x'$ is feasible for LP and $c^T(x^* + x') < p^*$

so use Farkas' lemma:

$$\begin{array}{ll} Ax + b = 0, \quad x \geq 0 & \text{or} \quad A^T y \geq 0, \quad b^T y < 0 \\ \begin{bmatrix} A & -b \\ c^T & -p^* \end{bmatrix} \begin{bmatrix} x \\ \tau \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \end{bmatrix} & \text{or} \quad \begin{bmatrix} A^T & c \\ -b^T & -p^* \end{bmatrix} \begin{bmatrix} y \\ \sigma \end{bmatrix} \geq 0, \quad \sigma > 0 \end{array}$$

use second system to show y/σ is dual feasible and optimal

Strong duality and complementary slackness

Definition (complementary slackness)

The primal-dual pair x and y are **complementary** if

$$\langle y, b - Ax \rangle = 0$$

They satisfy **strict complementary slackness** if $y_i(b_i - a_i^T x) = 0$ for $i = 1, \dots, n$.

for conic problem, strong duality \iff complementary slackness

$$\begin{aligned}\langle y, s \rangle &= \langle y, b - Ax \rangle \\ &= \langle y, b \rangle - \langle A^* y, x \rangle \\ &= \langle y, b \rangle - \langle c, x \rangle\end{aligned}$$