

CME 307 / MS&E 311 / OIT 676: Optimization

Optimality conditions and convexity

Professor Udell

Management Science and Engineering
Stanford

October 13, 2025

Outline

Constrained and unconstrained optimization

Optimality conditions

Convex analysis

Convex optimization

Constrained vs unconstrained optimization

constrained optimization

- ▶ examples: scheduling, routing, packing, logistics, scheduling, control
- ▶ what's hard: finding a feasible point

unconstrained optimization

- ▶ examples: data fitting, statistical/machine learning
- ▶ what's hard: reducing the objective

both are necessary for real-world problems!

Unconstrained smooth optimization

for $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ctly differentiable,

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{variable} & x \in \mathbb{R}^n \end{array}$$

examples:

- ▶ least squares
- ▶ logistic regression
- ▶ neural network training (with smooth activation like tanh, ELU, GeLU, ...)
- ▶ ...

Oracles

an optimization **oracle** is your interface for accessing the problem data:
e.g., an oracle for $f : \mathbb{R}^n \rightarrow \mathbb{R}$ can evaluate for any $x \in \mathbb{R}^n$:

- ▶ **zero-order:** $f_0(x)$
- ▶ **first-order:** $f_0(x)$ and $\nabla f_0(x)$
- ▶ **second-order:** $f_0(x)$, $\nabla f_0(x)$, and $\nabla^2 f_0(x)$

why oracles?

- ▶ can optimize real systems based on observed output (not just models)
- ▶ can use and extend old or complex but trusted code (e.g., NASA, PDE simulations, ...)
- ▶ can prove lower bounds on the oracle complexity of a problem class

source: Nesterov 2004 “Introductory Lectures on Convex Optimization”

Outline

Constrained and unconstrained optimization

Optimality conditions

Convex analysis

Convex optimization

Solution of an optimization problem

$$\text{minimize } f(x)$$

for $f : \mathcal{D} \rightarrow \mathbb{R}$. x^* is a

- ▶ **global minimizer** if $f(x) \geq f(x^*)$ for all $x \in \mathcal{D}$.
- ▶ **local minimizer** if there is a neighborhood \mathcal{N} around x^* so that $f(x) \geq f(x^*)$ for all $x \in \mathcal{N}$.
- ▶ **isolated local minimizer** if the neighborhood \mathcal{N} contains no other local minimizers.
- ▶ **unique minimizer** if it is the only global minimizer.

Solution of an optimization problem

$$\text{minimize } f(x)$$

for $f : \mathcal{D} \rightarrow \mathbb{R}$. x^* is a

- ▶ **global minimizer** if $f(x) \geq f(x^*)$ for all $x \in \mathcal{D}$.
- ▶ **local minimizer** if there is a neighborhood \mathcal{N} around x^* so that $f(x) \geq f(x^*)$ for all $x \in \mathcal{N}$.
- ▶ **isolated local minimizer** if the neighborhood \mathcal{N} contains no other local minimizers.
- ▶ **unique minimizer** if it is the only global minimizer.

pictures!

First order optimality condition

Theorem

If $x^ \in \mathbb{R}^n$ is a local minimizer of a differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, then $\nabla f(x^*) = 0$.*

First order optimality condition

Theorem

If $x^* \in \mathbb{R}^n$ is a local minimizer of a differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, then $\nabla f(x^*) = 0$.

proof: suppose by contradiction that $\nabla f(x^*) \neq 0$. consider points of the form $x_\alpha = x^* - \alpha \nabla f(x^*)$ for $\alpha > 0$. by definition of the gradient,

$$\lim_{\alpha \rightarrow 0} \frac{f(x_\alpha) - f(x^*)}{\alpha} = -\nabla f(x^*)^\top \nabla f(x^*) = -\|\nabla f(x^*)\|^2 < 0$$

so for any sufficiently small $\alpha > 0$, we have $f(x_\alpha) < f(x^*)$, which contradicts the fact that x^* is a local minimizer.

First order optimality condition

Theorem

If $x^* \in \mathbb{R}^n$ is a local minimizer of a differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, then $\nabla f(x^*) = 0$.

proof: suppose by contradiction that $\nabla f(x^*) \neq 0$. consider points of the form $x_\alpha = x^* - \alpha \nabla f(x^*)$ for $\alpha > 0$. by definition of the gradient,

$$\lim_{\alpha \rightarrow 0} \frac{f(x_\alpha) - f(x^*)}{\alpha} = -\nabla f(x^*)^\top \nabla f(x^*) = -\|\nabla f(x^*)\|^2 < 0$$

so for any sufficiently small $\alpha > 0$, we have $f(x_\alpha) < f(x^*)$, which contradicts the fact that x^* is a local minimizer.

Definition

$x^* \in \mathbb{R}^n$ is a **stationary point** of a differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ if $\nabla f(x^*) = 0$.

is a stationary point always a local minimizer?

First order optimality condition

Theorem

If $x^* \in \mathbb{R}^n$ is a local minimizer of a differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, then $\nabla f(x^*) = 0$.

proof: suppose by contradiction that $\nabla f(x^*) \neq 0$. consider points of the form $x_\alpha = x^* - \alpha \nabla f(x^*)$ for $\alpha > 0$. by definition of the gradient,

$$\lim_{\alpha \rightarrow 0} \frac{f(x_\alpha) - f(x^*)}{\alpha} = -\nabla f(x^*)^\top \nabla f(x^*) = -\|\nabla f(x^*)\|^2 < 0$$

so for any sufficiently small $\alpha > 0$, we have $f(x_\alpha) < f(x^*)$, which contradicts the fact that x^* is a local minimizer.

Definition

$x^* \in \mathbb{R}^n$ is a **stationary point** of a differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ if $\nabla f(x^*) = 0$.

is a stationary point always a local minimizer? no! saddle points, local maximizers.

Second order optimality condition

Theorem

If $x^ \in \mathbb{R}^n$ is a local minimizer of a twice differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, then $\nabla^2 f(x^*) \succeq 0$.*

Second order optimality condition

Theorem

If $x^ \in \mathbb{R}^n$ is a local minimizer of a twice differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, then $\nabla^2 f(x^*) \succeq 0$.*

proof: similar to the previous proof. use the fact that the second order approximation

$$f(x_\alpha) \approx f(x^*) + \nabla f(x^*)^\top (x_\alpha - x^*) + \frac{1}{2}(x_\alpha - x^*)^\top \nabla^2 f(x^*)(x_\alpha - x^*)$$

is accurate locally to show a contradiction unless $\nabla^2 f(x^*) \succeq 0$: if not, there is a direction v such that $v^\top \nabla^2 f(x^*) v < 0$. then $f(x + \alpha v) < f(x^*)$ for α arbitrarily small, which contradicts the fact that x^* is a local minimizer.

Symmetric positive semidefinite matrices

Definition

a symmetric matrix $Q \in \mathbb{R}^{n \times n}$ is **positive semidefinite** (psd) if $x^T Q x \geq 0$ for all $x \in \mathbb{R}^n$.

these matrices are so important that there are many ways to write them! for $Q \in \mathbb{R}^{n \times n}$,

$$Q \in \mathbf{S}_+^n \iff Q \succeq 0 \iff Q = Q^T, \lambda_{\min}(Q) \geq 0 \iff v^T Q v \geq 0 \quad \forall v \in \mathbb{R}^n$$

Symmetric positive semidefinite matrices

Definition

a symmetric matrix $Q \in \mathbb{R}^{n \times n}$ is **positive semidefinite** (psd) if $x^T Q x \geq 0$ for all $x \in \mathbb{R}^n$.

these matrices are so important that there are many ways to write them! for $Q \in \mathbb{R}^{n \times n}$,

$$Q \in \mathbf{S}_+^n \iff Q \succeq 0 \iff Q = Q^T, \lambda_{\min}(Q) \geq 0 \iff v^T Q v \geq 0 \quad \forall v \in \mathbb{R}^n$$

$Q \in \mathbf{S}_{++}^n$ is **symmetric positive definite** (spd) ($Q \succ 0$) if $x^T Q x > 0$ for all $x \neq 0$.

Symmetric positive semidefinite matrices

Definition

a symmetric matrix $Q \in \mathbb{R}^{n \times n}$ is **positive semidefinite** (psd) if $x^T Q x \geq 0$ for all $x \in \mathbb{R}^n$.

these matrices are so important that there are many ways to write them! for $Q \in \mathbb{R}^{n \times n}$,

$$Q \in \mathbf{S}_+^n \iff Q \succeq 0 \iff Q = Q^T, \lambda_{\min}(Q) \geq 0 \iff v^T Q v \geq 0 \quad \forall v \in \mathbb{R}^n$$

$Q \in \mathbf{S}_{++}^n$ is **symmetric positive definite** (spd) ($Q \succ 0$) if $x^T Q x > 0$ for all $x \neq 0$.

why care about psd matrices Q ?

- ▶ least-squares objective is quadratic with psd Hessian $A^T A$
- ▶ level sets of $x^T Q x$ are (bounded) ellipsoids if $Q \succ 0$
- ▶ the quadratic form $x^T Q x$ is a metric iff $Q \succ 0$
- ▶ eigenvalue decomp and svd coincide for psd matrices

Outline

Constrained and unconstrained optimization

Optimality conditions

Convex analysis

Convex optimization

Convex sets

Definition

A set $S \subseteq \mathbb{R}^n$ is convex if it contains every chord: for all $\theta \in [0, 1]$, $w, v \in S$,

$$\theta w + (1 - \theta)v \in S$$

Convex sets

Definition

A set $S \subseteq \mathbb{R}^n$ is convex if it contains every chord: for all $\theta \in [0, 1]$, $w, v \in S$,

$$\theta w + (1 - \theta)v \in S$$

Q: Which of these are convex?

ellipsoid, crescent moon, ...

Operations that preserve convexity

if $S \subseteq \mathbb{R}^n$ and $T \subseteq \mathbb{R}^n$ are convex, then so are:

- ▶ intersection: $S \cap T$
- ▶ sum: $S + T = \{s + t \mid s \in S, t \in T\}$
- ▶ projection: $\{x : (x, y) \in S\}$

Convex functions

a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex iff

Convex functions

a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex iff

► **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbb{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

Convex functions

a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex iff

- ▶ **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbb{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.** $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$ is convex

Convex functions

a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex iff

- ▶ **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbb{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.** $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$ is convex
- ▶ **First order condition.** if f is differentiable,

$$f(v) \geq f(w) + \nabla f(w)^\top (v - w), \quad \forall w, v \in \mathbb{R}^n$$

Convex functions

a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex iff

- ▶ **Chords.** it never lies above its chord: $\forall \theta \in [0, 1], w, v \in \mathbb{R}^n$

$$f(\theta w + (1 - \theta)v) \leq \theta f(w) + (1 - \theta)f(v)$$

- ▶ **Epigraph.** $\text{epi}(f) = \{(x, t) : t \geq f(x)\}$ is convex
- ▶ **First order condition.** if f is differentiable,

$$f(v) \geq f(w) + \nabla f(w)^\top (v - w), \quad \forall w, v \in \mathbb{R}^n$$

- ▶ **Second order condition.** If f is twice differentiable, its Hessian is always psd:

$$\lambda_{\min}(\nabla^2 f(x)) \geq 0, \quad \forall x \in \mathbb{R}^n$$

Convexity examples

Q: Which of these functions are convex?

- ▶ quadratic function $f(x) = x^2$ for $x \in \mathbb{R}$
- ▶ absolute value function $f(x) = |x|$ for $x \in \mathbb{R}$
- ▶ quadratic function $f(x) = x^T A x$, $x \in \mathbb{R}^n$, $A \succeq 0$
- ▶ quadratic function $f(x) = x^T A x$, A indefinite
- ▶ rollercoaster function (cubic) $f(x) = (x - 1)(x - 3)(x - 5)$
- ▶ hyperbolic function $f(x) = 1/x$ for $x > 0$
- ▶ jump function $f(x) = 1$ if $x \geq 0$, $f(x) = 0$ otherwise
- ▶ jump to infinity function $f(x) = 1$ if $x \in [-1, 1]$, $f(x) = \infty$ otherwise

Operations that preserve convexity

if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}$ are convex, then so are:

- ▶ cf for $c \geq 0$
- ▶ $f(Ax + b)$ for $A \in \mathbb{R}^{n \times m}$, $b \in \mathbb{R}^n$
- ▶ $f + g$
- ▶ $\max\{f, g\}$

Operations that preserve convexity

if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}$ are convex, then so are:

- ▶ cf for $c \geq 0$
- ▶ $f(Ax + b)$ for $A \in \mathbb{R}^{n \times m}$, $b \in \mathbb{R}^n$
- ▶ $f + g$
- ▶ $\max\{f, g\}$

Q: Pick one and assume f and g are twice-differentiable. What is the easiest way to prove convexity?

Operations that preserve convexity

if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}$ are convex, then so are:

- ▶ cf for $c \geq 0$
- ▶ $f(Ax + b)$ for $A \in \mathbb{R}^{n \times m}$, $b \in \mathbb{R}^n$
- ▶ $f + g$
- ▶ $\max\{f, g\}$

Q: Pick one and assume f and g are twice-differentiable. What is the easiest way to prove convexity?

most general rule:

$f \circ g(x) = f(g(x))$ is convex if g is convex and f is convex and nondecreasing

since

$$(f \circ g)''(x) = f''(g(x))(g'(x))^2 + f'(g(x))g''(x)$$

Jensen's inequality

Jensen's inequality generalizes the chord condition to a distribution of points:

Theorem

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex and X is a random variable, then

$$f(\mathbb{E}[X]) \leq \mathbb{E}[f(X)]$$

Sublevel set

Definition

The **sublevel set** of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ at level t is

$$S_t = \{x \in \mathbb{R}^n \mid f(x) \leq t\}$$

Sublevel set

Definition

The **sublevel set** of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ at level t is

$$S_t = \{x \in \mathbb{R}^n \mid f(x) \leq t\}$$

Theorem

A convex function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ has convex sublevel sets.

Sublevel set

Definition

The **sublevel set** of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ at level t is

$$S_t = \{x \in \mathbb{R}^n \mid f(x) \leq t\}$$

Theorem

A convex function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ has convex sublevel sets.

proof: Jensen's inequality. if $x, y \in S_t$, then for $\theta \in [0, 1]$,

$$f(\theta x + (1 - \theta)y) \leq \theta f(x) + (1 - \theta)f(y) \leq \theta t + (1 - \theta)t = t$$

so $\theta x + (1 - \theta)y \in S_t$.

Quasiconvexity

converse is not true: a function can have all sublevel sets convex, and still be non-convex.

Definition

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is **quasiconvex** if its sublevel sets are convex.

examples of functions that are quasiconvex but not convex?

Supporting hyperplane

Definition

A **supporting hyperplane** to a set $S \subseteq \mathbb{R}^n$ at a point $x \in S$ is a hyperplane that touches S at x and lies entirely on one side of S :

$$H = \{y \in \mathbb{R}^n \mid a^\top y = b\} \text{ supports } S \text{ at } x \text{ if } \begin{array}{ll} a^\top x &= b \\ a^\top y &\geq b \quad \forall y \in S \end{array}$$

Supporting hyperplane

Definition

A **supporting hyperplane** to a set $S \subseteq \mathbb{R}^n$ at a point $x \in S$ is a hyperplane that touches S at x and lies entirely on one side of S :

$$H = \{y \in \mathbb{R}^n \mid a^\top y = b\} \text{ supports } S \text{ at } x \text{ if } \begin{array}{ll} a^\top x &= b \\ a^\top y &\geq b \quad \forall y \in S \end{array}$$

Theorem (Supporting hyperplane)

Any nonempty convex set has a supporting hyperplane at every boundary point.

Supporting hyperplane condition for convexity

Theorem (Partial converse)

If a closed set with nonempty interior has a supporting hyperplane at every boundary point, then it is convex.

Supporting hyperplane condition for convexity

Theorem (Partial converse)

If a closed set with nonempty interior has a supporting hyperplane at every boundary point, then it is convex.

Theorem

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex \iff for all $x \in \mathbf{relint\,dom\,}f$, the epigraph of f has a supporting hyperplane at $(x, f(x))$: for some $g \in \mathbb{R}^n$,

$$f(y) \geq f(x) + g^\top(y - x) \quad \forall y \in \mathbb{R}^n$$

Supporting hyperplane condition for convexity

Theorem (Partial converse)

If a closed set with nonempty interior has a supporting hyperplane at every boundary point, then it is convex.

Theorem

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex \iff for all $x \in \mathbf{relint\,dom\,} f$, the epigraph of f has a supporting hyperplane at $(x, f(x))$: for some $g \in \mathbb{R}^n$,

$$f(y) \geq f(x) + g^\top(y - x) \quad \forall y \in \mathbb{R}^n$$

generalizes first-order condition for convexity to non-differentiable functions!

Supporting hyperplane condition for convexity

Theorem (Partial converse)

If a closed set with nonempty interior has a supporting hyperplane at every boundary point, then it is convex.

Theorem

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is convex \iff for all $x \in \text{relint dom } f$, the epigraph of f has a supporting hyperplane at $(x, f(x))$: for some $g \in \mathbb{R}^n$,

$$f(y) \geq f(x) + g^\top(y - x) \quad \forall y \in \mathbb{R}^n$$

generalizes first-order condition for convexity to non-differentiable functions!

Definition

A vector $g \in \mathbb{R}^n$ is a **subgradient** of $f : \mathbb{R}^n \rightarrow \mathbb{R}$ at $x \in \mathbb{R}^n$ if $f(y) \geq f(x) + g^\top(y - x)$ for all $y \in \mathbb{R}^n$.

Example: subgradients

$f = \max\{f_1, f_2\}$, with f_1, f_2 convex and differentiable

Q: Where is the function f differentiable? Where is the subgradient unique?

Subdifferential

set of all subgradients of f at x is called the **subdifferential** $\partial f(x)$

$$\partial f(x) = \{g : f(y) \geq f(x) + g^T(y - x) \quad \forall y\}$$

Subdifferential

set of all subgradients of f at x is called the **subdifferential** $\partial f(x)$

$$\partial f(x) = \{g : f(y) \geq f(x) + g^T(y - x) \quad \forall y\}$$

for any f ,

- ▶ $\partial f(x)$ is a closed convex set (can be empty)
- ▶ $\partial f(x) = \emptyset$ if $f(x) = \infty$

proof: use the definition

Subdifferential

set of all subgradients of f at x is called the **subdifferential** $\partial f(x)$

$$\partial f(x) = \{g : f(y) \geq f(x) + g^T(y - x) \quad \forall y\}$$

for any f ,

- ▶ $\partial f(x)$ is a closed convex set (can be empty)
- ▶ $\partial f(x) = \emptyset$ if $f(x) = \infty$

proof: use the definition

if f is convex,

- ▶ $\partial f(x)$ is nonempty, for $x \in \mathbf{relint\,dom\,} f$
- ▶ $\partial f(x) = \{\nabla f(x)\}$, if f is differentiable at x
- ▶ if $\partial f(x) = \{g\}$, then f is differentiable at x and $g = \nabla f(x)$

Outline

Constrained and unconstrained optimization

Optimality conditions

Convex analysis

Convex optimization

Convex optimization

an optimization problem is convex if:

► **Geometrically:** the feasible set and the epigraph of the objective are convex

for example, a nonlinear minimization is convex if the objective and inequality constraints are convex functions, and the equality constraints are affine

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq b_i, \quad i = 1, \dots, m_1 \\ & Ax = b_2 \\ \text{variable} & x \in \mathbb{R}^n\end{array}$$

Convex optimization

an optimization problem is convex if:

- ▶ **Geometrically:** the feasible set and the epigraph of the objective are convex

for example, a nonlinear minimization is convex if the objective and inequality constraints are convex functions, and the equality constraints are affine

$$\begin{array}{ll}\text{minimize} & f_0(x) \\ \text{subject to} & f_i(x) \leq b_i, \quad i = 1, \dots, m_1 \\ & Ax = b_2 \\ \text{variable} & x \in \mathbb{R}^n\end{array}$$

concave functions:

- ▶ a function f is concave if $-f$ is convex
- ▶ concave maximization \implies a convex optimization problem

Why care about convex optimization?

- ▶ local optimality \Rightarrow global optimality
- ▶ efficient solvers
- ▶ conceptual tools that generalize linear programming:
duality, stopping conditions, ...

Local minima are global for convex functions

Theorem

If x^ is a local minimizer of a convex function f , then x^* is a global minimizer.*

Local minima are global for convex functions

Theorem

If x^ is a local minimizer of a convex function f , then x^* is a global minimizer.*

proof?

Local minima are global for convex functions

Theorem

If x^ is a local minimizer of a convex function f , then x^* is a global minimizer.*

proof? suppose by contradiction that another point x' is a global minimizer, with $f(x') < f(x^*)$. draw the chord between x' and x^* . since the chord lies above f , every convex combination $x = \theta x^* + (1 - \theta)x'$ of x' and x^* for $\theta \in (0, 1)$ has a value $f(x) < f(x^*)$. this is true even for $x \rightarrow x^*$, contradicting our assumption that x^* is a local minimizer.

Corollary

Corollary

If f is convex and differentiable and $\nabla f(x^) = 0$, then x^* is a global minimizer.*

Corollary

Corollary

If f is convex and differentiable and $\nabla f(x^) = 0$, then x^* is a global minimizer.*

Q: Is a global minimizer of a convex function always unique?

Corollary

Corollary

If f is convex and differentiable and $\nabla f(x^) = 0$, then x^* is a global minimizer.*

Q: Is a global minimizer of a convex function always unique?

A: No. Picture.