

CME 307 / MS&E 311: Optimization

Duality

Professor Udell

Management Science and Engineering
Stanford

May 3, 2023

Announcements

- ▶ meet with course staff to discuss project this week or next (see Ed)
- ▶ project 1 due this Friday 5/5

Outline

Duality

Lagrange duality

Duality

Definition (Dual space)

The **dual** \mathcal{X}^* of a vector space \mathcal{X} is the set of linear functionals on \mathcal{X} .

so if $x \in \mathcal{X}$ and you see someone write

$$w^T x, \quad \langle w, x \rangle, \quad \text{or} \quad w \cdot x$$

you know that $w \in \mathcal{X}^*$ is a dual vector

Duality

Definition (Dual space)

The **dual** \mathcal{X}^* of a vector space \mathcal{X} is the set of linear functionals on \mathcal{X} .

so if $x \in \mathcal{X}$ and you see someone write

$$w^T x, \quad \langle w, x \rangle, \quad \text{or} \quad w \cdot x$$

you know that $w \in \mathcal{X}^*$ is a dual vector

notation: solution to optimization problem x^* vs dual space \mathcal{X}^*

Careful of units!

example 1: suppose $y_i = w^T x_i$ where

$$x_i = \begin{bmatrix} \text{heart rate} \\ \text{blood pressure} \\ \text{age} \end{bmatrix}, \quad \text{with units} \quad \begin{bmatrix} \text{bpm} \\ \text{mmHg} \\ \text{years} \end{bmatrix}$$

and y_i is duration of stay in hospital (units: days)

Careful of units!

example 1: suppose $y_i = w^T x_i$ where

$$x_i = \begin{bmatrix} \text{heart rate} \\ \text{blood pressure} \\ \text{age} \end{bmatrix}, \quad \text{with units} \quad \begin{bmatrix} \text{bpm} \\ \text{mmHg} \\ \text{years} \end{bmatrix}$$

and y_i is duration of stay in hospital (units: days)

then w has units of

$$\begin{bmatrix} \text{days/bpm} \\ \text{days/mmHg} \\ \text{days/year} \end{bmatrix}$$

Careful of units!

example 2: $f(x) = \sum_{i=1}^n \left(1 + \exp \left(\underbrace{y_i w^T x_i}_{\text{input must be a scalar!}} \right) \right)$

Careful of units!

example 2: $f(x) = \sum_{i=1}^n \left(1 + \exp \left(\underbrace{y_i w^T x_i}_{\text{input must be a scalar!}} \right) \right)$

example 3: if $x \in \mathcal{X}$, gradient is a linear function on $\mathcal{X} \implies \nabla f(x_0) \in \mathcal{X}^*$

$$f(x) \approx f(x_0) + \nabla f(x_0)^T (x - x_0),$$

so gradient descent stepsize t has units

$$x^{k+1} = x^k - t \nabla f(x^k)$$

e.g., x (meters m), $\nabla f(x)$ (m^{-1}), and t (m^2)

Careful of units!

example 2: $f(x) = \sum_{i=1}^n \left(1 + \exp \left(\underbrace{y_i w^T x_i}_{\text{input must be a scalar!}} \right) \right)$

example 3: if $x \in \mathcal{X}$, gradient is a linear function on $\mathcal{X} \implies \nabla f(x_0) \in \mathcal{X}^*$

$$f(x) \approx f(x_0) + \nabla f(x_0)^T (x - x_0),$$

so gradient descent stepsize t has units

$$x^{k+1} = x^k - t \nabla f(x^k)$$

e.g., x (meters m), $\nabla f(x)$ (m^{-1}), and t (m^2)

- ▶ no wonder it's hard to choose the stepsize!
- ▶ basic recommendation: standardize your data

Dual of function space

- ▶ $f : [0, 1] \rightarrow \mathbf{R}$ is a function
- ▶ $f(x)$ is a linear function of f , for any x :

$$(f + g)(x) = f(x) + g(x), \quad (cf)(x) = cf(x)$$

- ▶ so is any integral:

$$\int_0^1 f(x) d\mu(x)$$

\implies the dual of the space of functions on $[0, 1]$ is the space of measures on $[0, 1]$

Dual norm

Definition (Dual norm)

The **dual norm** of a norm $\| \cdot \|$ is

$$\|w\|_* = \sup_{\|x\| \leq 1} \langle w, x \rangle$$

equivalently, $\|w\|_* = \sup_x \frac{\langle w, x \rangle}{\|x\|}$

Dual norm

Definition (Dual norm)

The **dual norm** of a norm $\|\cdot\|$ is

$$\|w\|_* = \sup_{\|x\| \leq 1} \langle w, x \rangle$$

equivalently, $\|w\|_* = \sup_x \frac{\langle w, x \rangle}{\|x\|}$

example: ℓ_1 norm dual is ℓ_∞ norm

$$\|w\|_1 = \sum_{i=1}^n |w_i|, \quad \|w\|_\infty = \max_{i=1, \dots, n} |w_i|$$

Dual norm

Definition (Dual norm)

The **dual norm** of a norm $\| \cdot \|$ is

$$\|w\|_* = \sup_{\|x\| \leq 1} \langle w, x \rangle$$

equivalently, $\|w\|_* = \sup_x \frac{\langle w, x \rangle}{\|x\|}$

example: ℓ_1 norm dual is ℓ_∞ norm

$$\|w\|_1 = \sum_{i=1}^n |w_i|, \quad \|w\|_\infty = \max_{i=1, \dots, n} |w_i|$$

example: ℓ_2 norm dual is ℓ_2 norm $\implies \ell_2$ is **self-dual**

Dual norm

Definition (Dual norm)

The **dual norm** of a norm $\|\cdot\|$ is

$$\|w\|_* = \sup_{\|x\| \leq 1} \langle w, x \rangle$$

equivalently, $\|w\|_* = \sup_x \frac{\langle w, x \rangle}{\|x\|}$

example: ℓ_1 norm dual is ℓ_∞ norm

$$\|w\|_1 = \sum_{i=1}^n |w_i|, \quad \|w\|_\infty = \max_{i=1, \dots, n} |w_i|$$

example: ℓ_2 norm dual is ℓ_2 norm $\implies \ell_2$ is **self-dual**

example: for $f : [0, 1] \rightarrow \mathbf{R}$, if $\|f\| = \sup_{x \in [0, 1]} |f(x)|$,

$$\|\mu\|_* = \sup_{\|f\| \leq 1} \int_0^1 f(x) d\mu(x) = \int_0^1 d|\mu|(x)$$

Self-dual norms

given primal space \mathcal{X}

- ▶ dual vector is a linear functional $w(x)$ on $x \in \mathcal{X}$
- ▶ we should define the dual norm on \mathcal{X}^* as

$$\sup_{x \in \mathcal{X}, \|x\| \leq 1} w(x)$$

- ▶ but instead we used the inner product $\langle w, x \rangle$. why?

Self-dual norms

given primal space \mathcal{X}

- ▶ dual vector is a linear functional $w(x)$ on $x \in \mathcal{X}$
- ▶ we should define the dual norm on \mathcal{X}^* as

$$\sup_{x \in \mathcal{X}, \|x\| \leq 1} w(x)$$

- ▶ but instead we used the inner product $\langle w, x \rangle$. why?

Theorem (Riesz representation)

Suppose $\mathcal{X} = H$ is a Hilbert (inner product) space. For any linear functional $\phi \in \mathcal{X}^$, there is a unique vector $w \in H$ so that $w(x) = \langle w, x \rangle$ for all $x \in \mathcal{X} = H$. Moreover, $\|w\|_* = \|w\|$.*

$\|\cdot\|$ is self-dual $\iff \|\cdot\|$ is induced by an inner product

Self-dual norms

given primal space \mathcal{X}

- ▶ dual vector is a linear functional $w(x)$ on $x \in \mathcal{X}$
- ▶ we should define the dual norm on \mathcal{X}^* as

$$\sup_{x \in \mathcal{X}, \|x\| \leq 1} w(x)$$

- ▶ but instead we used the inner product $\langle w, x \rangle$. why?

Theorem (Riesz representation)

Suppose $\mathcal{X} = H$ is a Hilbert (inner product) space. For any linear functional $\phi \in \mathcal{X}^$, there is a unique vector $w \in H$ so that $w(x) = \langle w, x \rangle$ for all $x \in \mathcal{X} = H$. Moreover, $\|w\|_* = \|w\|$.*

$\|\cdot\|$ is self-dual $\iff \|\cdot\|$ is induced by an inner product

example: ℓ_2 norm is self-dual, induced by the inner product

$$\langle w, x \rangle = w^T x$$

Conjugate of linear operator

given $x \in \mathbf{R}^n$, $w \in \mathbf{R}^m$, and $A \in \mathbf{R}^{m \times n}$, conjugate of A is the linear operator A^* defined so that

$$\langle A^* w, x \rangle = \langle w, Ax \rangle$$

Conjugate of linear operator

given $x \in \mathbf{R}^n$, $w \in \mathbf{R}^m$, and $A \in \mathbf{R}^{m \times n}$, conjugate of A is the linear operator A^* defined so that

$$\langle A^* w, x \rangle = \langle w, Ax \rangle$$

example: $x \in \mathbf{R}^n$, $A \in \mathbf{R}^{m \times n}$ defined by

$$Ax = \begin{bmatrix} x_{i_1} \\ \vdots \\ x_{i_m} \end{bmatrix}$$

then $A^* \in \mathbf{R}^{n \times m}$ satisfies

$$\langle A^* w, x \rangle = \langle w, Ax \rangle = \sum_{j=1}^m w_j x_{i_j},$$

so A^* creates a sparse vector from w with

$$(A^* w)_{i_j} = w_j$$

Fenchel dual

Definition (Fenchel dual)

The **Fenchel dual** of a function $f : \mathcal{X} \rightarrow \mathbf{R}$ is

$$f^*(w) = \sup_{x \in \mathcal{X}} \langle w, x \rangle - f(x)$$

also called the **conjugate function**. draw picture!

<https://remilepriol.github.io/dualityviz/>

Fenchel dual

Definition (Fenchel dual)

The **Fenchel dual** of a function $f : \mathcal{X} \rightarrow \mathbf{R}$ is

$$f^*(w) = \sup_{x \in \mathcal{X}} \langle w, x \rangle - f(x)$$

also called the **conjugate function**. draw picture!

<https://remilepriol.github.io/dualityviz/>

example: $f(x) = \|x\|_1, x \in \mathbf{R}^n$

$$f^*(w) = \sup_{x \in \mathbf{R}^n} \langle w, x \rangle - \|x\|_1 = \begin{cases} 0 & \|w\|_\infty \leq 1 \\ \infty & \text{otherwise} \end{cases}$$

\implies fenchel dual of ℓ_1 norm is indicator of ℓ_∞ ball

Biconjugate

Definition (Biconjugate)

The **biconjugate** of a function $f : \mathcal{X} \rightarrow \mathbf{R}$ is

$$f^{**}(x) = \sup_{w \in \mathcal{X}^*} \langle w, x \rangle - f^*(w)$$

- ▶ for convex $f : \mathbf{R} \rightarrow \mathbf{R}$, $f^{**} = f$
- ▶ for nonconvex f , f^{**} is convex hull of f

\implies biconjugate is a convexification operation

Biconjugate

Definition (Biconjugate)

The **biconjugate** of a function $f : \mathcal{X} \rightarrow \mathbf{R}$ is

$$f^{**}(x) = \sup_{w \in \mathcal{X}^*} \langle w, x \rangle - f^*(w)$$

► for convex $f : \mathbf{R} \rightarrow \mathbf{R}$, $f^{**} = f$

► for nonconvex f , f^{**} is convex hull of f

\implies biconjugate is a convexification operation

example: consider $f : \mathbf{R} \rightarrow \mathbf{R}$ defined by

$$f(x) = \begin{cases} 0 & x \in \{-1, 1\} \\ \infty & \text{otherwise} \end{cases}$$

what is f_* ? f^{**} ?

Outline

Duality

Lagrange duality

Why duality?

- ▶ certify optimality
 - ▶ turn \forall into \exists
 - ▶ use dual lower bound to derive stopping conditions
- ▶ new algorithms based on the dual
 - ▶ solve dual, then recover primal solution

Warmup: Farkas lemma

Theorem (Farkas lemma)

Given $A \in \mathbf{R}^{m \times n}$ and $b \in \mathbf{R}^m$, exactly one of the following is true:

- ▶ *there exists $x \in \mathbf{R}^n$ so that $Ax = b$ and $x \geq 0$*
- ▶ *there exists $y \in \mathbf{R}^m$ so that $A^T y \geq 0$ and $\langle b, y \rangle < 0$*

\implies can efficiently certify infeasibility of a linear program

Warmup: Farkas lemma

Theorem (Farkas lemma)

Given $A \in \mathbf{R}^{m \times n}$ and $b \in \mathbf{R}^m$, exactly one of the following is true:

- ▶ there exists $x \in \mathbf{R}^n$ so that $Ax = b$ and $x \geq 0$
- ▶ there exists $y \in \mathbf{R}^m$ so that $A^T y \geq 0$ and $\langle b, y \rangle < 0$

\implies can efficiently certify infeasibility of a linear program

proof: suppose we have $x \in \mathbf{R}^n$ so that $Ax = b$ and $x \geq 0$.
then for any $y \in \mathbf{R}^m$,

$$\begin{aligned} 0 &= \langle y, b - Ax \rangle = \langle y, b \rangle - \langle A^T y, x \rangle \\ \langle y, b \rangle &= \langle A^T y, x \rangle \end{aligned}$$

so if $A^T y \geq 0$, then use $x \geq 0$ to conclude $\langle y, b \rangle \geq 0$.

Warmup: Farkas lemma

Theorem (Farkas lemma)

Given $A \in \mathbf{R}^{m \times n}$ and $b \in \mathbf{R}^m$, exactly one of the following is true:

- ▶ there exists $x \in \mathbf{R}^n$ so that $Ax = b$ and $x \geq 0$
- ▶ there exists $y \in \mathbf{R}^m$ so that $A^T y \geq 0$ and $\langle b, y \rangle < 0$

\implies can efficiently certify infeasibility of a linear program

proof: suppose we have $x \in \mathbf{R}^n$ so that $Ax = b$ and $x \geq 0$.
then for any $y \in \mathbf{R}^m$,

$$\begin{aligned} 0 &= \langle y, b - Ax \rangle = \langle y, b \rangle - \langle A^T y, x \rangle \\ \langle y, b \rangle &= \langle A^T y, x \rangle \end{aligned}$$

so if $A^T y \geq 0$, then use $x \geq 0$ to conclude $\langle y, b \rangle \geq 0$.

(opposite direction is similar)

Lagrange duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

if x is feasible, then $Ax = b$, so $\langle y, Ax - b \rangle = 0$.

Lagrange duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

if x is feasible, then $Ax = b$, so $\langle y, Ax - b \rangle = 0$.

define the **Lagrangian**

$$\mathcal{L}(x, y) := f(x) - \langle y, b - Ax \rangle$$

Lagrange duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

if x is feasible, then $Ax = b$, so $\langle y, Ax - b \rangle = 0$.

define the **Lagrangian**

$$\begin{aligned} \mathcal{L}(x, y) &:= f(x) - \langle y, b - Ax \rangle \\ p^* &= \inf_{x: Ax=b} \mathcal{L}(x, y) \geq \inf_x \mathcal{L}(x, y) \end{aligned}$$

Lagrange duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

if x is feasible, then $Ax = b$, so $\langle y, Ax - b \rangle = 0$.

define the **Lagrangian**

$$\begin{aligned} \mathcal{L}(x, y) &:= f(x) - \langle y, b - Ax \rangle \\ p^* &= \inf_{x: Ax=b} \mathcal{L}(x, y) \geq \inf_x \mathcal{L}(x, y) \\ &= \inf_x f(x) + \langle y, -b + Ax \rangle \\ &= \langle y, -b \rangle + \inf_x \left(f(x) + \langle A^T y, x \rangle \right) \\ &= \langle y, -b \rangle - \sup_x \left(\langle -A^T y, x \rangle - f(x) \right) \\ &= \langle y, -b \rangle - f^*(-A^T y) = g(y) \end{aligned}$$

$g(y)$ is called the **dual function**

Lagrange duality

inequality holds for any $y \in \mathbf{R}^m$, so we have proved **weak duality**

$$\begin{aligned} p^* &\geq g(y) \quad \forall y \in \mathbf{R}^m \\ &\geq \underbrace{\sup_y g(y)}_{\mathcal{D}} =: d^* \end{aligned} \tag{1}$$

dual optimal value $d^* \leq p^*$

Strong duality

Definition (Duality gap)

The **duality gap** for a primal-dual pair (x, y) is $f(x) - g(y)$

by weak duality, duality gap is always nonnegative

Strong duality

Definition (Duality gap)

The **duality gap** for a primal-dual pair (x, y) is $f(x) - g(y)$

by weak duality, duality gap is always nonnegative

Definition (Strong duality)

A primal-dual pair (x^*, y^*) satisfies **strong duality** if

$$p^* = d^* \iff f(x^*) - g(y^*) = 0$$

Strong duality

Definition (Duality gap)

The **duality gap** for a primal-dual pair (x, y) is $f(x) - g(y)$

by weak duality, duality gap is always nonnegative

Definition (Strong duality)

A primal-dual pair (x^*, y^*) satisfies **strong duality** if

$$p^* = d^* \iff f(x^*) - g(y^*) = 0$$

strong duality holds

- ▶ for feasible LPs (pf later)
- ▶ for convex problems under **constraint qualification** aka **Slater's condition**. feasible region has an **interior point** x so that all inequality constraints hold strictly

strong duality fails if either (\mathcal{P}) or (\mathcal{D}) is infeasible or unbounded

Lagrange duality with inequality constraints

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax \leq b : \quad y \geq 0 \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

Lagrange duality with inequality constraints

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax \leq b : \quad y \geq 0 \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

to construct Lagrangian $\mathcal{L}(x, y) = f(x) - \langle y, b - Ax \rangle$, ensure value is **better** (lower) when x and y are feasible

Lagrange duality with inequality constraints

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax \leq b : \quad y \geq 0 \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

to construct Lagrangian $\mathcal{L}(x, y) = f(x) - \langle y, b - Ax \rangle$, ensure value is **better** (lower) when x and y are feasible

$$\mathcal{L}(x, y) := f(x) - \langle y, b - Ax \rangle$$

Lagrange duality with inequality constraints

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax \leq b : \quad y \geq 0 \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

to construct Lagrangian $\mathcal{L}(x, y) = f(x) - \langle y, b - Ax \rangle$, ensure value is **better** (lower) when x and y are feasible

$$\begin{aligned} \mathcal{L}(x, y) &:= f(x) - \langle y, b - Ax \rangle \\ p^* &\geq \inf_{x \text{ feas}} f(x) - \langle y, b - Ax \rangle \\ &\geq \inf_x f(x) - \langle y, b - Ax \rangle \\ &= \langle y, b \rangle - f^*(-A^*y) =: g(y) \end{aligned}$$

Lagrange duality with inequality constraints

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax \leq b : \quad y \geq 0 \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

to construct Lagrangian $\mathcal{L}(x, y) = f(x) - \langle y, b - Ax \rangle$, ensure value is **better** (lower) when x and y are feasible

$$\begin{aligned} \mathcal{L}(x, y) &:= f(x) - \langle y, b - Ax \rangle \\ p^* &\geq \inf_{x \text{ feas}} f(x) - \langle y, b - Ax \rangle \\ &\geq \inf_x f(x) - \langle y, b - Ax \rangle \\ &= \langle y, b \rangle - f^*(-A^*y) =: g(y) \end{aligned}$$

as before, this holds for all y , so we have weak duality

$$p^* \geq \underbrace{\sup_y g(y)}_{\mathcal{D}} =: d^*$$

SVM dual

support vector machine: for $x_i \in \mathbf{R}^n$, $y_i \in \{-1, 1\}$, $i = 1, \dots, m$

$$\text{minimize} \quad \frac{1}{2} \|w\|^2 + 1^T s$$

$$\text{subject to} \quad y_i w^T x_i + s_i \geq 1 \quad i = 1, \dots, m : \quad \alpha \geq 0$$
$$s \geq 0 : \quad \mu \geq 0$$

(SVM)

SVM dual

support vector machine: for $x_i \in \mathbf{R}^n$, $y_i \in \{-1, 1\}$, $i = 1, \dots, m$

$$\begin{aligned} & \text{minimize} && \frac{1}{2} \|w\|^2 + 1^T s \\ & \text{subject to} && y_i w^T x_i + s_i \geq 1 \quad i = 1, \dots, m : \quad \alpha \geq 0 \\ & && s \geq 0 : \quad \mu \geq 0 \end{aligned}$$

(SVM)

Lagrangian: for $\alpha \geq 0$, $\mu \geq 0$,

$$\mathcal{L}(w, s, \alpha, \mu) = \frac{1}{2} \|w\|^2 + 1^T s - \sum_{i=1}^m \alpha_i (y_i w^T x_i + s_i - 1) - \mu^T s$$

► minimize $\mathcal{L}(w, s, \alpha, \mu)$ over w :

$$w = \sum_{i=1}^m \alpha_i y_i x_i$$

► minimize $\mathcal{L}(w, s, \alpha, \mu)$ over $s \implies \alpha + \mu = 1$

SVM dual

so simplify:

$$\begin{aligned}g(\alpha) &= \inf_{w,s} \mathcal{L}(w, s, \alpha, 1 - \alpha) \\&= \frac{1}{2} \|w\|^2 - w^T \sum_{i=1}^m \alpha_i y_i x_i + \mathbf{1}^T \alpha \\&= -\frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 + \mathbf{1}^T \alpha\end{aligned}$$

SVM dual

so simplify:

$$\begin{aligned}g(\alpha) &= \inf_{w,s} \mathcal{L}(w, s, \alpha, 1 - \alpha) \\&= \frac{1}{2} \|w\|^2 - w^T \sum_{i=1}^m \alpha_i y_i x_i + 1^T \alpha \\&= -\frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 + 1^T \alpha\end{aligned}$$

define $K \in \mathbf{R}^m$ so $K_{ij} = y_i y_j x_i^T x_j$. then

$$\left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 = \sum_{i=1}^m \sum_{j=1}^m \alpha_i \alpha_j y_i y_j x_i^T x_j = \alpha^T K \alpha$$

SVM dual

so simplify:

$$\begin{aligned}g(\alpha) &= \inf_{w,s} \mathcal{L}(w, s, \alpha, 1 - \alpha) \\&= \frac{1}{2} \|w\|^2 - w^T \sum_{i=1}^m \alpha_i y_i x_i + 1^T \alpha \\&= -\frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 + 1^T \alpha\end{aligned}$$

define $K \in \mathbf{R}^m$ so $K_{ij} = y_i y_j x_i^T x_j$. then

$$\left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 = \sum_{i=1}^m \sum_{j=1}^m \alpha_i \alpha_j y_i y_j x_i^T x_j = \alpha^T K \alpha$$

dual problem:

$$\begin{array}{ll}\text{maximize} & -\frac{1}{2} \alpha^T K \alpha + 1^T \alpha \\ \text{subject to} & \alpha \geq 0\end{array} \quad (\text{SVM-dual})$$

SVM dual

so simplify:

$$\begin{aligned}g(\alpha) &= \inf_{w,s} \mathcal{L}(w, s, \alpha, 1 - \alpha) \\&= \frac{1}{2} \|w\|^2 - w^T \sum_{i=1}^m \alpha_i y_i x_i + 1^T \alpha \\&= -\frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 + 1^T \alpha\end{aligned}$$

define $K \in \mathbf{R}^m$ so $K_{ij} = y_i y_j x_i^T x_j$. then

$$\left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 = \sum_{i=1}^m \sum_{j=1}^m \alpha_i \alpha_j y_i y_j x_i^T x_j = \alpha^T K \alpha$$

dual problem:

$$\begin{array}{ll}\text{maximize} & -\frac{1}{2} \alpha^T K \alpha + 1^T \alpha \\ \text{subject to} & \alpha \geq 0\end{array} \quad (\text{SVM-dual})$$

new solution ideas! coordinate descent on α (SMO), kernel trick

Generalize Lagrangian duality

Generalize Lagrangian duality

- **nonlinear duality:** replace

$$0 \geq Ax - b \quad \text{with} \quad 0 \geq g(x)$$

(harder to derive explicit form for dual problem)

Generalize Lagrangian duality

- ▶ **nonlinear duality:** replace

$$0 \geq Ax - b \quad \text{with} \quad 0 \geq g(x)$$

(harder to derive explicit form for dual problem)

- ▶ **conic duality:** for cone K , replace

$$b - Ax \geq 0 \quad \text{with} \quad b - Ax \in K$$

define **slack vector** $s = b - Ax \in K$
for weak duality, dual y must satisfy

$$\langle y, s \rangle \geq 0 \quad \forall s \in K$$

Dual cones

this inequality defines the **dual cone** K^* :

Definition (dual cone)

the dual cone K^* of a cone K is the set of vectors y such that

$$\langle y, s \rangle \geq 0 \quad \forall s \in K$$

Dual cones

this inequality defines the **dual cone** K^* :

Definition (dual cone)

the dual cone K^* of a cone K is the set of vectors y such that

$$\langle y, s \rangle \geq 0 \quad \forall s \in K$$

examples of cones and their duals:

- ▶ K acute, K^* obtuse
- ▶ $K = \mathbf{R}_+^m$, $K^* = \mathbf{R}_+^m$
- ▶ $K = \{x \in \mathbf{R}^n \mid \|x\| \leq x_0\}$, $K^* = \{y \in \mathbf{R}^n \mid \|y\| \leq y_0\}$
- ▶ $K = \{X \in \mathbf{S}^n \mid X \succeq 0\}$, $K^* = \{Y \in \mathbf{S}^n \mid Y \succeq 0\}$

Conic duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

Conic duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

for $y \in K^*$, construct Lagrangian $\mathcal{L}(x, y) = \langle c, x \rangle - \langle y, b - Ax \rangle$,
ensure value is **better** (lower) when x and y are feasible

Conic duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll} \text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

for $y \in K^*$, construct Lagrangian $\mathcal{L}(x, y) = \langle c, x \rangle - \langle y, b - Ax \rangle$,
ensure value is **better** (lower) when x and y are feasible

$$\mathcal{L}(x, y) \quad := \quad \langle c, x \rangle - \langle y, b - Ax \rangle$$

Conic duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll}\text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n\end{array} \quad (\mathcal{P})$$

for $y \in K^*$, construct Lagrangian $\mathcal{L}(x, y) = \langle c, x \rangle - \langle y, b - Ax \rangle$,
ensure value is **better** (lower) when x and y are feasible

$$\begin{aligned}\mathcal{L}(x, y) &:= \langle c, x \rangle - \langle y, b - Ax \rangle \\ p^* &\geq \inf_{x \text{ feas}} \langle c, x \rangle - \langle y, b - Ax \rangle \\ &\geq \inf_x \langle c, x \rangle - \langle y, b - Ax \rangle \\ &= \langle y, -b \rangle + \inf_x \langle c + A^*y, x \rangle\end{aligned}$$

Conic duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll}\text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n\end{array} \quad (\mathcal{P})$$

for $y \in K^*$, construct Lagrangian $\mathcal{L}(x, y) = \langle c, x \rangle - \langle y, b - Ax \rangle$,
ensure value is **better** (lower) when x and y are feasible

$$\begin{aligned}\mathcal{L}(x, y) &:= \langle c, x \rangle - \langle y, b - Ax \rangle \\ p^* &\geq \inf_{x \text{ feas}} \langle c, x \rangle - \langle y, b - Ax \rangle \\ &\geq \inf_x \langle c, x \rangle - \langle y, b - Ax \rangle \\ &= \langle y, -b \rangle + \inf_x \langle c + A^*y, x \rangle\end{aligned}$$

which is $-\infty$ unless $c + A^*y = 0$, so

Conic duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll}\text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n\end{array} \quad (\mathcal{P})$$

for $y \in K^*$, construct Lagrangian $\mathcal{L}(x, y) = \langle c, x \rangle - \langle y, b - Ax \rangle$,
ensure value is **better** (lower) when x and y are feasible

$$\begin{aligned}\mathcal{L}(x, y) &:= \langle c, x \rangle - \langle y, b - Ax \rangle \\ p^* &\geq \inf_{x \text{ feas}} \langle c, x \rangle - \langle y, b - Ax \rangle \\ &\geq \inf_x \langle c, x \rangle - \langle y, b - Ax \rangle \\ &= \langle y, -b \rangle + \inf_x \langle c + A^*y, x \rangle\end{aligned}$$

which is $-\infty$ unless $c + A^*y = 0$, so define the **dual problem**

$$\begin{array}{ll}\text{maximize} & \langle y, -b \rangle \\ \text{subject to} & c + A^*y = 0 \\ \text{variable} & y \in K^*\end{array} \quad (\mathcal{D})$$

Conic duality

primal problem with solution $x^* \in \mathbf{R}^n$, optimal value p^* :

$$\begin{array}{ll}\text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n\end{array} \quad (\mathcal{P})$$

for $y \in K^*$, construct Lagrangian $\mathcal{L}(x, y) = \langle c, x \rangle - \langle y, b - Ax \rangle$,
ensure value is **better** (lower) when x and y are feasible

$$\begin{aligned}\mathcal{L}(x, y) &:= \langle c, x \rangle - \langle y, b - Ax \rangle \\ p^* &\geq \inf_{x \text{ feas}} \langle c, x \rangle - \langle y, b - Ax \rangle \\ &\geq \inf_x \langle c, x \rangle - \langle y, b - Ax \rangle \\ &= \langle y, -b \rangle + \inf_x \langle c + A^*y, x \rangle\end{aligned}$$

which is $-\infty$ unless $c + A^*y = 0$, so define the **dual problem**

$$\begin{array}{ll}\text{maximize} & \langle y, -b \rangle \\ \text{subject to} & c + A^*y = 0 \\ \text{variable} & y \in K^*\end{array} \quad (\mathcal{D})$$

Dual of the dual

- ▶ if (\mathcal{P}) is convex, then the dual of (1) is (\mathcal{P})
- ▶ otherwise, the dual of the dual is the **convexification** of the primal

picture

Strong duality for LPs

primal and dual LP in standard form:

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

$$\begin{array}{ll}\text{maximize} & b^T y \\ \text{subject to} & A^T y \leq c\end{array}$$

claim: if primal LP has a bounded feasible solution x^* , then strong duality holds

i.e., dual LP has a bounded feasible solution y^* and $p^* = d^*$

Proof of strong duality for LPs

consider the following system with variables $x' \in \mathbf{R}^n$, $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

Proof of strong duality for LPs

consider the following system with variables $x' \in \mathbf{R}^n$, $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

claim: this system has no solution. pf by contradiction:

- ▶ if $\tau > 0$, then x'/τ is feasible for LP and $c^T x'/\tau < p^*$
- ▶ if $\tau = 0$, then $x^* + x'$ is feasible for LP and $c^T(x^* + x') < p^*$

Proof of strong duality for LPs

consider the following system with variables $x' \in \mathbf{R}^n$, $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

claim: this system has no solution. pf by contradiction:

- ▶ if $\tau > 0$, then x'/τ is feasible for LP and $c^T x'/\tau < p^*$
- ▶ if $\tau = 0$, then $x^* + x'$ is feasible for LP and $c^T(x^* + x') < p^*$

so use Farkas' lemma:

$$Ax + b = 0, \quad x \geq 0 \quad \text{or} \quad A^T y \geq 0, \quad b^T y < 0$$

Proof of strong duality for LPs

consider the following system with variables $x' \in \mathbf{R}^n$, $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

claim: this system has no solution. pf by contradiction:

- ▶ if $\tau > 0$, then x'/τ is feasible for LP and $c^T x'/\tau < p^*$
- ▶ if $\tau = 0$, then $x^* + x'$ is feasible for LP and $c^T(x^* + x') < p^*$

so use Farkas' lemma:

$$\begin{array}{ll} Ax + b = 0, \quad x \geq 0 & \text{or} \quad A^T y \geq 0, \quad b^T y < 0 \\ \begin{bmatrix} A & -b \\ c^T & -p^* \end{bmatrix} \begin{bmatrix} x \\ \tau \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \end{bmatrix} & \text{or} \quad \begin{bmatrix} A^T & c \\ -b^T & -p^* \end{bmatrix} \begin{bmatrix} y \\ \sigma \end{bmatrix} \geq 0, \quad \sigma > 0 \end{array}$$

Proof of strong duality for LPs

consider the following system with variables $x' \in \mathbf{R}^n$, $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

claim: this system has no solution. pf by contradiction:

- ▶ if $\tau > 0$, then x'/τ is feasible for LP and $c^T x'/\tau < p^*$
- ▶ if $\tau = 0$, then $x^* + x'$ is feasible for LP and $c^T(x^* + x') < p^*$

so use Farkas' lemma:

$$\begin{array}{ll} Ax + b = 0, \quad x \geq 0 & \text{or} \quad A^T y \geq 0, \quad b^T y < 0 \\ \begin{bmatrix} A & -b \\ c^T & -p^* \end{bmatrix} \begin{bmatrix} x \\ \tau \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \end{bmatrix} & \text{or} \quad \begin{bmatrix} A^T & c \\ -b^T & -p^* \end{bmatrix} \begin{bmatrix} y \\ \sigma \end{bmatrix} \geq 0, \quad \sigma > 0 \end{array}$$

use second system to show y/σ is dual feasible and optimal

Strong duality and complementary slackness

Definition (complementary slackness)

The primal-dual pair x and y are **complementary** if

$$\langle y, b - Ax \rangle = 0$$

They satisfy **strict complementary slackness** if $y_i(b_i - a_i^T x) = 0$ for $i = 1, \dots, n$.

for conic problem, strong duality \iff complementary slackness

$$\begin{aligned}\langle y, s \rangle &= \langle y, b - Ax \rangle \\ &= \langle y, b \rangle - \langle A^* y, x \rangle \\ &= \langle y, b \rangle - \langle c, x \rangle\end{aligned}$$

KKT conditions

KKT conditions give **necessary** conditions for optimality of convex problem.

Theorem (KKT conditions)

Suppose x^ and y^* are primal and dual optimal, respectively.
Then*

- ▶ **stationarity.** x^* and y^* are a min/max saddle point of the Lagrangian

$$\nabla_x \mathcal{L}(x^*, y^*) = 0, \quad \nabla_y \mathcal{L}(x^*, y^*) = 0$$

- ▶ **feasibility.** x^* is primal feasible; y^* is dual feasible
- ▶ **complementary slackness.** x^* and y^* are complementary:

$$\langle y^*, b - Ax^* \rangle = 0$$

KKT conditions turn optimization problem into a system of equations

KKT Example

Consider the following optimization problem:

$$\begin{array}{ll}\text{minimize} & x^2 + y^2 \\ \text{subject to} & x + y \leq 1 : \quad \lambda \geq 0 \\ & x - y = 0 : \quad \mu\end{array}$$

Lagrangian:

KKT Example

Consider the following optimization problem:

$$\begin{array}{ll}\text{minimize} & x^2 + y^2 \\ \text{subject to} & x + y \leq 1 : \quad \lambda \geq 0 \\ & x - y = 0 : \quad \mu\end{array}$$

Lagrangian:

$$\mathcal{L}(x, y, \lambda, \mu) = x^2 + y^2 + \lambda(x + y - 1) + \mu(x - y)$$

Lagrangian:

$$\mathcal{L}(x, y, \lambda, \mu) = x^2 + y^2 + \lambda(x + y - 1) + \mu(x - y)$$

Lagrangian:

$$\mathcal{L}(x, y, \lambda, \mu) = x^2 + y^2 + \lambda(x + y - 1) + \mu(x - y)$$

KKT conditions:

1. stationarity: $\nabla L(x, y, \lambda, \mu) = 0$
2. feasibility:
 - ▶ primal: $x + y \leq 1$ and $x - y = 0$
 - ▶ dual: $\lambda \geq 0$
3. complementary slackness: $\lambda(x + y - 1) = 0$

Lagrangian:

$$\mathcal{L}(x, y, \lambda, \mu) = x^2 + y^2 + \lambda(x + y - 1) + \mu(x - y)$$

KKT conditions:

1. stationarity: $\nabla L(x, y, \lambda, \mu) = 0$
2. feasibility:
 - ▶ primal: $x + y \leq 1$ and $x - y = 0$
 - ▶ dual: $\lambda \geq 0$
3. complementary slackness: $\lambda(x + y - 1) = 0$

Taking the gradient of L wrt x , y , λ , and μ , we get:

$$\frac{\partial \mathcal{L}}{\partial x} = 2x + \lambda + \mu = 0$$

$$\frac{\partial \mathcal{L}}{\partial y} = 2y + \lambda - \mu = 0$$

$$\frac{\partial \mathcal{L}}{\partial \lambda} = x + y - 1 = 0$$

$$\frac{\partial \mathcal{L}}{\partial \mu} = x - y = 0$$

solve!

Lagrangian:

$$\mathcal{L}(x, y, \lambda, \mu) = x^2 + y^2 + \lambda(x + y - 1) + \mu(x - y)$$

KKT conditions:

1. stationarity: $\nabla L(x, y, \lambda, \mu) = 0$
2. feasibility:
 - ▶ primal: $x + y \leq 1$ and $x - y = 0$
 - ▶ dual: $\lambda \geq 0$
3. complementary slackness: $\lambda(x + y - 1) = 0$

Taking the gradient of L wrt x , y , λ , and μ , we get:

$$\frac{\partial \mathcal{L}}{\partial x} = 2x + \lambda + \mu = 0$$

$$\frac{\partial \mathcal{L}}{\partial y} = 2y + \lambda - \mu = 0$$

$$\frac{\partial \mathcal{L}}{\partial \lambda} = x + y - 1 = 0$$

$$\frac{\partial \mathcal{L}}{\partial \mu} = x - y = 0$$

solve! $\rightarrow x^* = 0.5, y^* = 0.5, \lambda^* = 0, \mu^* = 1$

Semidefinite program (SDP)

consider primal SDP with decision variable $X \in \mathbf{S}^n$:

$$\begin{array}{ll} \text{minimize} & \text{tr}(CX) \\ \text{subject to} & \mathcal{A}X = b \\ & X \succeq 0, \end{array} \quad (\mathcal{P})$$

problem data:

- ▶ cost matrix $C \in \mathbf{S}^n$
- ▶ linear map $\mathcal{A} : \mathbf{R}^{n \times n} \rightarrow \mathbf{R}^m$
- ▶ righthand side $b \in \mathbf{R}^m$

First order methods use too much storage

suppose (\mathcal{P}) has


- ▶ **compact specification:** problem data use $\mathcal{O}(n)$ storage
- ▶ **compact solution:** solution X_\star has constant rank r_\star
 \implies solution uses $\mathcal{O}(n)$ storage

First order methods use too much storage

suppose (\mathcal{P}) has


- ▶ **compact specification:** problem data use $\mathcal{O}(n)$ storage
- ▶ **compact solution:** solution X_\star has constant rank r_\star
 \implies solution uses $\mathcal{O}(n)$ storage

(\mathcal{P}) , using any first order method:



Approximate complementarity requires less storage

(\mathcal{P}), using approximate complementarity:



Motivation: large scale SDP

Max-Cut [goemans1995improved]	Matrix Completion [srebro2005rank]
minimize $\text{tr}(-LX)$ subject to $\text{diag}(X) = \mathbf{1}$ $X \succeq 0$	minimize $\text{tr}(W_1) + \text{tr}(W_2)$ subject to $X_{ij} = M_{ij}, (i, j) \in \Omega$ $\begin{bmatrix} W_1 & X \\ X^* & W_2 \end{bmatrix} \succeq 0$

Motivation: large scale SDP

Max-Cut [goemans1995improved]	Matrix Completion [srebro2005rank]
minimize $\text{tr}(-LX)$ subject to $\text{diag}(X) = \mathbf{1}$ $X \succeq 0$	minimize $\text{tr}(W_1) + \text{tr}(W_2)$ subject to $X_{ij} = M_{ij}, (i, j) \in \Omega$ $\begin{bmatrix} W_1 & X \\ X^* & W_2 \end{bmatrix} \succeq 0$

- ▶ Matrix completion:
 - ▶ 10^9 users, 10^9 products
 - ▶ \implies SDP with 10^{18} variables

Motivation: large scale SDP

Max-Cut [goemans1995improved]	Matrix Completion [srebro2005rank]
minimize $\text{tr}(-LX)$ subject to $\text{diag}(X) = \mathbf{1}$ $X \succeq 0$	minimize $\text{tr}(W_1) + \text{tr}(W_2)$ subject to $X_{ij} = M_{ij}, (i, j) \in \Omega$ $\begin{bmatrix} W_1 & X \\ X^* & W_2 \end{bmatrix} \succeq 0$

- ▶ Matrix completion:
 - ▶ 10^9 users, 10^9 products
 - ▶ \implies SDP with 10^{18} variables
- ▶ MaxCut:
 - ▶ 10^9 people in social network
 - ▶ \implies SDP with 10^{18} variables

Motivation: large scale SDP

Max-Cut [goemans1995improved]	Matrix Completion [srebro2005rank]
minimize $\text{tr}(-LX)$ subject to $\text{diag}(X) = \mathbf{1}$ $X \succeq 0$	minimize $\text{tr}(W_1) + \text{tr}(W_2)$ subject to $X_{ij} = M_{ij}, (i, j) \in \Omega$ $\begin{bmatrix} W_1 & X \\ X^* & W_2 \end{bmatrix} \succeq 0$

- ▶ Matrix completion:
 - ▶ 10^9 users, 10^9 products
 - ▶ \implies SDP with 10^{18} variables
- ▶ MaxCut:
 - ▶ 10^9 people in social network
 - ▶ \implies SDP with 10^{18} variables
- ▶ Phase retrieval:
 - ▶ $10^3 \times 10^3$ discretization of sample
 - ▶ \implies SDP with 10^{12} variables

Optimal Storage

What kind of storage bounds can we hope for?

- Assume black-box implementation of

$$u \mapsto Cu, \quad (u, v) \mapsto \mathcal{A}(uv^*), \text{ and } \quad (u, y) \mapsto (\mathcal{A}^*y)u,$$

where $u, v \in \mathbf{R}^n$, $y \in \mathbf{R}^m$.

Optimal Storage

What kind of storage bounds can we hope for?

- ▶ Assume black-box implementation of

$$u \mapsto Cu, \quad (u, v) \mapsto \mathcal{A}(uv^*), \text{ and } \quad (u, y) \mapsto (\mathcal{A}^*y)u,$$

where $u, v \in \mathbf{R}^n$, $y \in \mathbf{R}^m$.

- ▶ Need $\Omega(m + n)$ storage to apply problem data.

Optimal Storage

What kind of storage bounds can we hope for?

- ▶ Assume black-box implementation of

$$u \mapsto Cu, \quad (u, v) \mapsto \mathcal{A}(uv^*), \text{ and } \quad (u, y) \mapsto (\mathcal{A}^*y)u,$$

where $u, v \in \mathbf{R}^n$, $y \in \mathbf{R}^m$.

- ▶ Need $\Omega(m + n)$ storage to apply problem data.
- ▶ Need $\Theta(rn)$ storage for a rank- r approximate solution.

Optimal Storage

What kind of storage bounds can we hope for?

- ▶ Assume black-box implementation of

$$u \mapsto Cu, \quad (u, v) \mapsto \mathcal{A}(uv^*), \text{ and } (u, y) \mapsto (\mathcal{A}^*y)u,$$

where $u, v \in \mathbf{R}^n$, $y \in \mathbf{R}^m$.

- ▶ Need $\Omega(m + n)$ storage to apply problem data.
- ▶ Need $\Theta(rn)$ storage for a rank- r approximate solution.

Definition. An algorithm to return a rank r (approximate) solution to (\mathcal{P}) has **optimal storage** if it uses working storage

$$\Theta(m + rn).$$

Dual problem

consider dual SDP with decision variable $y \in \mathbf{R}^m$:

$$\begin{array}{ll} \text{maximize} & b^* y \\ \text{subject to} & C - \mathcal{A}^* y \succeq 0 \end{array} \quad (\mathcal{D})$$

Dual problem

consider dual SDP with decision variable $y \in \mathbf{R}^m$:

$$\begin{array}{ll} \text{maximize} & b^*y \\ \text{subject to} & C - \mathcal{A}^*y \succeq 0 \end{array} \quad (\mathcal{D})$$

for sufficiently large $\alpha > 0$, equivalent to **penalized dual**

$$\text{maximize} \quad b^*y + \alpha \min(\lambda_{\min}(C - \mathcal{A}^*y), 0)$$

Dual problem

consider dual SDP with decision variable $y \in \mathbf{R}^m$:

$$\begin{array}{ll} \text{maximize} & b^*y \\ \text{subject to} & C - \mathcal{A}^*y \succeq 0 \end{array} \quad (\mathcal{D})$$

for sufficiently large $\alpha > 0$, equivalent to **penalized dual**

$$\text{maximize} \quad b^*y + \alpha \min(\lambda_{\min}(C - \mathcal{A}^*y), 0)$$

- ▶ any first order method for (\mathcal{D}) uses optimal storage
- ▶ (just compute minimal eigenvalue with iterative method)
- ▶ e.g., subgradient method, AdaGrad [[duchi2011adaptive](#)], AdaNGD [[levy2017online](#)], AccelGrad [[levy2018online](#)],
...

Assumptions

Assumption (Genericity)

- ▶ primal SDP attains its solution(s)
- ▶ dual SDP has a unique solution y_*
- ▶ primal and dual SDP satisfy strong duality

$$0 = \mathbf{tr}(CX_*) - b^*y_* = \mathbf{tr}(X_*(C - \mathcal{A}^*y_*)) = X_*(C - \mathcal{A}^*y_*)$$

- ▶ (for storage optimality) strict complementary slackness

$$\text{Rank}(X_*) + \text{Rank}(C - \mathcal{A}^*y_*) = n$$

\implies solution X_* of primal SDP is unique [lemon2016low]

Assumptions

Assumption (Genericity)

- ▶ primal SDP attains its solution(s)
- ▶ dual SDP has a unique solution y_*
- ▶ primal and dual SDP satisfy strong duality

$$0 = \mathbf{tr}(CX_*) - b^*y_* = \mathbf{tr}(X_*(C - \mathcal{A}^*y_*)) = X_*(C - \mathcal{A}^*y_*)$$

- ▶ (for storage optimality) strict complementary slackness

$$\text{Rank}(X_*) + \text{Rank}(C - \mathcal{A}^*y_*) = n$$

\implies solution X_* of primal SDP is unique [lemon2016low]

Note: these conditions hold generically
[alizadeh1997complementarity]

Primal recovery via exact complementarity

suppose $y_\star \in \mathbf{R}^m$ solves (\mathcal{D})

Primal recovery via exact complementarity

suppose $y_\star \in \mathbf{R}^m$ solves (\mathcal{D})

- ▶ define dual slack matrix $Z_\star = C - \mathcal{A}^* y_\star$

Primal recovery via exact complementarity

suppose $y_\star \in \mathbf{R}^m$ solves (\mathcal{D})

- ▶ define dual slack matrix $Z_\star = C - \mathcal{A}^* y_\star$
- ▶ strict complementarity holds between X_\star and Z_\star

$$\text{Rank}(X_\star) + \text{Rank}(Z_\star) = n \quad \implies \quad \text{range}(X_\star) = \mathbf{nullspace}(Z_\star)$$

Primal recovery via exact complementarity

suppose $y_\star \in \mathbf{R}^m$ solves (\mathcal{D})

- ▶ define dual slack matrix $Z_\star = C - \mathcal{A}^* y_\star$
- ▶ strict complementarity holds between X_\star and Z_\star

$$\text{Rank}(X_\star) + \text{Rank}(Z_\star) = n \implies \text{range}(X_\star) = \mathbf{nullspace}(Z_\star)$$

- ▶ let V_\star be a basis for nullspace of dual slack matrix Z_\star

Primal recovery via exact complementarity

suppose $y_\star \in \mathbf{R}^m$ solves (\mathcal{D})

- ▶ define dual slack matrix $Z_\star = C - \mathcal{A}^*y_\star$
- ▶ strict complementarity holds between X_\star and Z_\star

$$\text{Rank}(X_\star) + \text{Rank}(Z_\star) = n \quad \implies \quad \text{range}(X_\star) = \mathbf{nullspace}(Z_\star)$$

- ▶ let V_\star be a basis for nullspace of dual slack matrix Z_\star
- ▶ constrain $X = V_\star S V_\star^*$ in primal SDP for some $S \in \mathbf{S}_\star^{r_\star}$.
 \implies solution is preserved!

SDP via exact complementarity: algorithm

Algorithm SDP via exact complementarity

Given: problem data C, \mathcal{A}, b

SDP via exact complementarity: algorithm

Algorithm SDP via exact complementarity

Given: problem data C, \mathcal{A}, b

1. compute solution $y_\star \in \mathbf{R}^m$ to (\mathcal{D})

SDP via exact complementarity: algorithm

Algorithm SDP via exact complementarity

Given: problem data C, \mathcal{A}, b

1. compute solution $y_\star \in \mathbf{R}^m$ to (\mathcal{D})
2. compute basis V_\star for nullspace of dual slack matrix
 $Z_\star = C - \mathcal{A}^* y_\star$

SDP via exact complementarity: algorithm

Algorithm SDP via exact complementarity

Given: problem data C, \mathcal{A}, b

1. compute solution $y_\star \in \mathbf{R}^m$ to (\mathcal{D})
2. compute basis V_\star for nullspace of dual slack matrix
 $Z_\star = C - \mathcal{A}^* y_\star$
3. solve **reduced primal** (\mathcal{P}_{V_\star}) with variable $S \in \mathbf{S}_+^{r_\star}$

$$\begin{aligned} & \text{minimize} && \text{tr}(CV_\star S V_\star^*) \\ & \text{subject to} && \mathcal{A}(V_\star S V_\star^*) = b \\ & && S \succeq 0 \end{aligned} \quad (\mathcal{P}_{V_\star})$$

to find primal solution $X_\star = V_\star S V_\star^*$

SDP via exact complementarity: algorithm

Algorithm SDP via exact complementarity

Given: problem data C, \mathcal{A}, b

1. compute solution $y_\star \in \mathbf{R}^m$ to (\mathcal{D})
2. compute basis V_\star for nullspace of dual slack matrix
 $Z_\star = C - \mathcal{A}^* y_\star$
3. solve **reduced primal** (\mathcal{P}_{V_\star}) with variable $S \in \mathbf{S}_+^{r_\star}$

$$\begin{aligned} & \text{minimize} && \text{tr}(CV_\star SV_\star^*) \\ & \text{subject to} && \mathcal{A}(V_\star SV_\star^*) = b \\ & && S \succeq 0 \end{aligned} \quad (\mathcal{P}_{V_\star})$$

to find primal solution $X_\star = V_\star SV_\star^*$

Reduced primal is easy: e.g., when $r_\star=1$, (\mathcal{P}_{V_\star}) solves a 1D problem over $S \in \mathbf{R}_+$