

Duality

Lecture 6

October 8, 2025

Quiz

Recap From Last Time & Today's Plan

Last time...

- **Separating Hyperplane Thm \Rightarrow Farkas Lemma \Rightarrow Strong duality**

Agenda for today:

- Two motivating applications
- Implications of strong duality
- Optimality conditions and primal/dual simplex
- Complementary slackness
- Global sensitivity & Shadow prices as marginal costs
- One more application: network revenue management

Polynomially-Sized CVaR Representation

- Recall homework: ensure CVaR of portfolio payoff exceeds a lower limit
- CVaR was defined as the average over the k -smallest values (for suitable integer k)
- If payoffs in the scenarios are v_1, v_2, \dots, v_n , the key constraint is:

$$\sum_{i=1}^k v_{[i]} \geq b, \tag{1}$$

where $v_{[1]} \leq v_{[2]} \leq \dots \leq v_{[n]}$ is the sorted vector of payoffs.

Polynomially-Sized CVaR Representation

- Recall homework: ensure CVaR of portfolio payoff exceeds a lower limit
- CVaR was defined as the average over the k -smallest values (for suitable integer k)
- If payoffs in the scenarios are v_1, v_2, \dots, v_n , the key constraint is:

$$\sum_{i=1}^k v_{[i]} \geq b, \tag{1}$$

where $v_{[1]} \leq v_{[2]} \leq \dots \leq v_{[n]}$ is the sorted vector of payoffs.

- Can write one constraint for each vector in $\{0, 1\}^n$ with exactly k values of 1.
- *How to formulate with a polynomial number of variables and constraints?*

Application in Robust Optimization

- Consider an LP with an uncertain constraint:

$$a^T x \leq b, \tag{2}$$

where a satisfies $a \in \mathcal{A}$ and \mathcal{A} is polyhedral

- We seek decisions x that are **robustly feasible**, i.e.,

$$a^T x \leq b, \forall a \in \mathcal{A} := \{a \in \mathbb{R}^n : Ca \leq d\} \tag{3}$$

Infinitely many constraints : “semi-infinite” LP. *Any ideas?*

Application in Robust Optimization

- Consider an LP with an uncertain constraint:

$$a^T x \leq b, \tag{2}$$

where a satisfies $a \in \mathcal{A}$ and \mathcal{A} is polyhedral

- We seek decisions x that are **robustly feasible**, i.e.,

$$a^T x \leq b, \forall a \in \mathcal{A} := \{a \in \mathbb{R}^n : Ca \leq d\} \tag{3}$$

Infinitely many constraints : “semi-infinite” LP. *Any ideas?*

Strong Duality

Consider the following primal-dual pair:

$$\begin{aligned} (\mathcal{P}) \quad & \text{minimize } c^T x \\ & \text{subject to } Ax \geq b \end{aligned}$$

Strong Duality

Consider the following primal-dual pair:

$$\begin{aligned} (\mathcal{P}) \quad & \text{minimize } c^T x \\ & \text{subject to } Ax \geq b \end{aligned}$$

Theorem (**Strong Duality**)

If (\mathcal{P}) has an optimal solution, so does (\mathcal{D}) , and their optimal values are equal.

Implications

Strong duality leaves only a few possibilities for a primal-dual pair:

		Dual		
		Finite Optimum	Unbounded	Infeasible
Primal	Finite Optimum	?	?	?
	Unbounded	?	?	?
	Infeasible	?	?	?

Implications

Strong duality leaves only a few possibilities for a primal-dual pair:

		Dual		
		Finite Optimum	Unbounded	Infeasible
Primal	Finite Optimum	?	?	?
	Unbounded	?	?	?
	Infeasible	?	?	?

		Dual		
		Finite Optimum	Unbounded	Infeasible
Primal	Finite Optimum	Possible	Impossible	Impossible
	Unbounded	Impossible	Impossible	Possible
	Infeasible	Impossible	Possible	?

Strong Duality and Theorems of Alternative

- Strong duality allows you to **prove** various “theorems of alternative”

Example (Farkas Lemma)

Prove that exactly one of the following is true:

- (i) $\exists x \geq 0$ such that $Ax = b$,
- (ii) $\exists \lambda$ such that $\lambda^T A \geq 0$ and $\lambda^T b < 0$.

Optimality for Standard-Form LPs

$$(\mathcal{P}) \min c^T x$$

$$Ax = b, \quad x \geq 0$$

$$(\mathcal{D}) \max \lambda^T b$$

$$\lambda^T A \leq c^T$$

- (\mathcal{P}) achieves optimality at a **basic feasible solution** x :

Optimality for Standard-Form LPs

$$(\mathcal{P}) \min c^T x$$

$$Ax = b, \quad x \geq 0$$

$$(\mathcal{D}) \max \lambda^T b$$

$$\lambda^T A \leq c^T$$

- (\mathcal{P}) achieves optimality at a **basic feasible solution** x :
 - If $B \subseteq \{1, \dots, n\}$ is a basis, the b.f.s. is: $x = [x_B, 0]$, $x_B = A_B^{-1}b$.
 - Simplex algorithm: feasibility and optimality for (\mathcal{P}) are given by:

$$\text{Feasibility-}(\mathcal{P}) : \quad x_B := A_B^{-1}b \geq 0 \quad (4a)$$

$$\text{Optimality-}(\mathcal{P}) : \quad c^T - c_B^T A_B^{-1} A \geq 0 \quad (4b)$$

Optimality for Standard-Form LPs

$$(\mathcal{P}) \min c^T x$$

$$Ax = b, \quad x \geq 0$$

$$(\mathcal{D}) \max \lambda^T b$$

$$\lambda^T A \leq c^T$$

- (\mathcal{P}) achieves optimality at a **basic feasible solution** x :
 - If $B \subseteq \{1, \dots, n\}$ is a basis, the b.f.s. is: $x = [x_B, 0]$, $x_B = A_B^{-1}b$.
 - Simplex algorithm: feasibility and optimality for (\mathcal{P}) are given by:

$$\text{Feasibility-}(\mathcal{P}) : \quad x_B := A_B^{-1}b \geq 0 \quad (4a)$$

$$\text{Optimality-}(\mathcal{P}) : \quad c^T - c_B^T A_B^{-1} A \geq 0 \quad (4b)$$

- (\mathcal{D}) : same basis B can also be used to determine a **dual vector** λ :

$$\lambda^T A_i = c_i, \quad \forall i \in B \quad \Rightarrow \quad \lambda^T = c_B^T A_B^{-1}, \quad \forall i \in B.$$

Optimality for Standard-Form LPs

$$(\mathcal{P}) \min c^T x$$

$$Ax = b, \quad x \geq 0$$

$$(\mathcal{D}) \max \lambda^T b$$

$$\lambda^T A \leq c^T$$

- (\mathcal{P}) achieves optimality at a **basic feasible solution** x :
 - If $B \subseteq \{1, \dots, n\}$ is a basis, the b.f.s. is: $x = [x_B, 0]$, $x_B = A_B^{-1}b$.
 - Simplex algorithm: feasibility and optimality for (\mathcal{P}) are given by:

$$\text{Feasibility-}(\mathcal{P}) : \quad x_B := A_B^{-1}b \geq 0 \quad (4a)$$

$$\text{Optimality-}(\mathcal{P}) : \quad c^T - c_B^T A_B^{-1} A \geq 0 \quad (4b)$$

- (\mathcal{D}) : same basis B can also be used to determine a **dual vector** λ :

$$\lambda^T A_i = c_i, \quad \forall i \in B \quad \Rightarrow \quad \lambda^T = c_B^T A_B^{-1}, \quad \forall i \in B.$$

- The dual objective value corresponding to λ is: $\lambda^T b = c_B^T A_B^{-1} b = c^T x$
- λ is feasible in the dual if and only if:

$$\text{Feasibility-}(\mathcal{D}) : \quad c^T - \lambda^T A \geq 0 \quad \Leftrightarrow \quad c^T - c_B^T A_B^{-1} A \geq 0 \quad (5)$$

Optimality for Standard-Form LPs

$$(\mathcal{P}) \min c^T x$$

$$Ax = b, \quad x \geq 0$$

$$(\mathcal{D}) \max \lambda^T b$$

$$\lambda^T A \leq c^T$$

- (\mathcal{P}) achieves optimality at a **basic feasible solution** x :
 - If $B \subseteq \{1, \dots, n\}$ is a basis, the b.f.s. is: $x = [x_B, 0]$, $x_B = A_B^{-1}b$.
 - Simplex algorithm: feasibility and optimality for (\mathcal{P}) are given by:

$$\text{Feasibility-}(\mathcal{P}) : \quad x_B := A_B^{-1}b \geq 0 \quad (4a)$$

$$\text{Optimality-}(\mathcal{P}) : \quad c^T - c_B^T A_B^{-1} A \geq 0 \quad (4b)$$

- (\mathcal{D}) : same basis B can also be used to determine a **dual vector** λ :

$$\lambda^T A_i = c_i, \quad \forall i \in B \quad \Rightarrow \quad \lambda^T = c_B^T A_B^{-1}, \quad \forall i \in B.$$

- The dual objective value corresponding to λ is: $\lambda^T b = c_B^T A_B^{-1} b = c^T x$
- λ is feasible in the dual if and only if:

$$\text{Feasibility-}(\mathcal{D}) : \quad c^T - \lambda^T A \geq 0 \quad \Leftrightarrow \quad c^T - c_B^T A_B^{-1} A \geq 0 \quad (5)$$

Primal optimality \Leftrightarrow Dual feasibility

Simplex terminates when finding a dual-feasible solution!

Solve (\mathcal{P}) or (\mathcal{D}) ?

$$\begin{aligned} (\mathcal{P}) \quad & \min c^T x \\ & Ax = b, \quad x \geq 0 \end{aligned}$$

$$\begin{aligned} (\mathcal{D}) \quad & \max \lambda^T b \\ & \lambda^T A \leq c^T \end{aligned}$$

Solve (\mathcal{P}) or (\mathcal{D}) ?

$$(\mathcal{P}) \quad \min c^T x \\ Ax = b, \quad x \geq 0$$

$$(\mathcal{D}) \quad \max \lambda^T b \\ \lambda^T A \leq c^T$$

Primal simplex

- maintain a **basic feasible solution**
- basis $B \subset \{1, \dots, n\}$
- stopping criterion: dual feasibility

Dual simplex

- maintain a dual feasible solution
- stopping criterion: primal feasibility
- different from primal simplex: works with an LP with inequalities

- How to choose (\mathcal{P}) or (\mathcal{D}) ?
- Suppose we have x^* , λ^* and must now solve a **larger** problem, i.e., with extra decisions or extra constraints.
- *Any preference between primal and dual simplex?*
- Modern solvers include primal and dual simplex and allow concurrent runs

Optimality Conditions and Complementary Slackness

Primal-Dual Pair of Problems

$$\begin{array}{llll} (\mathcal{P}) & \underset{x}{\text{minimize}} & c^T x & \\ & & Ax \leq b & \\ & & x \geq 0 & \\ \text{variables} & & x \in \mathbb{R}^n & \end{array}$$

$$\begin{array}{llll} (\mathcal{D}) & \underset{\lambda}{\text{maximize}} & \lambda^T b & \\ & & \lambda \geq 0 & \\ & & \lambda^T A \leq c^T & \\ \text{variables} & & \lambda \in \mathbb{R}^m. & \end{array}$$

Consider $x \in P, \lambda \in D$ (each feasible). How to check if they are **optimal**?

Optimality Conditions and Complementary Slackness

Primal-Dual Pair of Problems

(\mathcal{P}) minimize $c^T x$ $Ax \leq b$ $x \geq 0$ variables $x \in \mathbb{R}^n$	(\mathcal{D}) maximize $\lambda^T b$ $\lambda \geq 0$ $\lambda^T A \leq c^T$ variables $\lambda \in \mathbb{R}^m$
---	--

Consider $x \in P, \lambda \in D$ (each feasible). How to check if they are **optimal**?

Theorem (Complementary Slackness)

$x \in P$ and $\lambda \in D$ are **optimal** solutions for (\mathcal{P}) and (\mathcal{D}) , respectively, **if and only if**:

$$\lambda_i (a_i^T x - b_i) = 0, i = 1, \dots, m$$

$$(\lambda^T A_j - c_j) x_j = 0, j = 1, \dots, n.$$

- Follows from primal/dual feasibility and $c^T x = b^T \lambda$

Optimality Conditions and Complementary Slackness

Primal-Dual Pair of Problems

$ \begin{array}{llll} (\mathcal{P}) & \underset{x}{\text{minimize}} & c^T x & \\ & & Ax & \leq b \\ & & x & \geq 0 \\ & \text{variables} & x & \in \mathbb{R}^n \end{array} $	$ \begin{array}{llll} (\mathcal{D}) & \underset{\lambda}{\text{maximize}} & \lambda^T b & \\ & & \lambda & \leq 0 \\ & & \lambda^T A & \leq c^T \\ & \text{variables} & \lambda & \in \mathbb{R}^m. \end{array} $
--	--

Consider $x \in P, \lambda \in D$ (each feasible). How to check if they are **optimal**?

Theorem (Complementary Slackness)

$x \in P$ and $\lambda \in D$ are **optimal** solutions for (\mathcal{P}) and (\mathcal{D}) , respectively, **if and only if**:

$$\begin{aligned}
 \lambda_i (a_i^T x - b_i) &= 0, \quad i = 1, \dots, m \\
 (\lambda^T A_j - c_j) x_j &= 0, \quad j = 1, \dots, n.
 \end{aligned}$$


- Follows from primal/dual feasibility and $c^T x = b^T \lambda$
- Interesting insight: **non-binding constraint** \Rightarrow dual variable is **zero**

Representation of Polyhedra

Important consequence of duality: alternative representation of all polyhedra

Definition

Consider a nonempty polyhedron $P = \{x \in \mathbb{R}^n : Ax \geq b\}$. Then:




Representation of Polyhedra

Important consequence of duality: alternative representation of all polyhedra

Definition

Consider a nonempty polyhedron $P = \{x \in \mathbb{R}^n : Ax \geq b\}$. Then:

1. $\mathcal{C} := \{d \in \mathbb{R}^n : Ad \geq 0\}$ is called the **recession cone** of P .
2. Any $d \in \mathcal{C}$ with $d \neq 0$ is called a **ray** of P .





Representation of Polyhedra

Important consequence of duality: alternative representation of all polyhedra

Definition

Consider a nonempty polyhedron $P = \{x \in \mathbb{R}^n : Ax \geq b\}$. Then:

1. $\mathcal{C} := \{d \in \mathbb{R}^n : Ad \geq 0\}$ is called the **recession cone** of P .
2. Any $d \in \mathcal{C}$ with $d \neq 0$ is called a **ray** of P .




Representation of Polyhedra

Important consequence of duality: alternative representation of all polyhedra

Definition

Consider a nonempty polyhedron $P = \{x \in \mathbb{R}^n : Ax \geq b\}$. Then:

1. $\mathcal{C} := \{d \in \mathbb{R}^n : Ad \geq 0\}$ is called the **recession cone** of P .
2. Any $d \in \mathcal{C}$ with $d \neq 0$ is called a **ray** of P .
3. Any ray d that satisfies $a_i^\top d = 0$ for $n - 1$ linearly independent a_i is called an **extreme ray** of P .



Representation of Polyhedra

Theorem (Resolution Theorem)


Let $P = \{x \in \mathbb{R}^n : Ax \geq b\}$ be a non-empty polyhedron, x^1, x^2, \dots, x^k be its **extreme points**, and w^1, w^2, \dots, w^r be its **extreme rays**. Then,

Representation of Polyhedra

Theorem (Resolution Theorem)

Let $P = \{x \in \mathbb{R}^n : Ax \geq b\}$ be a non-empty polyhedron, x^1, x^2, \dots, x^k be its **extreme points**, and w^1, w^2, \dots, w^r be its **extreme rays**. Then,

$$P = \text{conv}(\{x^1, \dots, x^k\}) + \text{cone}(\{w^1, \dots, w^r\})$$
$$= \left\{ \sum_{i=1}^k \mu_i x^i + \sum_{j=1}^r \theta_j w^j : \mu \geq 0, e^T \mu = 1, \theta \geq 0 \right\}.$$




Representation of Polyhedra

Theorem (Resolution Theorem)

Let $P = \{x \in \mathbb{R}^n : Ax \geq b\}$ be a non-empty polyhedron, x^1, x^2, \dots, x^k be its **extreme points**, and w^1, w^2, \dots, w^r be its **extreme rays**. Then,

$$P = \text{conv}(\{x^1, \dots, x^k\}) + \text{cone}(\{w^1, \dots, w^r\})$$
$$= \left\{ \sum_{i=1}^k \mu_i x^i + \sum_{j=1}^r \theta_j w^j : \mu \geq 0, e^T \mu = 1, \theta \geq 0 \right\}.$$



Note: It is **not** “easy” (i.e., poly-time) to switch between these representations

Dual Variables **As Marginal Costs**

$$\begin{array}{ll} (\mathcal{P}) \min c^T x & (\mathcal{D}) \max \lambda^T b \\ Ax = b, \quad x \geq 0 & \lambda^T A \leq c^T \end{array}$$

- Solved the LP and obtained x^* and λ^*
- Want to show that λ^* is the **gradient of the optimal cost with respect to b** “almost everywhere”
- Related to **sensitivity analysis**
How do the optimal value and solution depend on problem data A, b, c ?

Global Dependency On b, c

$$\begin{array}{ll} (\mathcal{P}) \min c^T x & (\mathcal{D}) \max \lambda^T b \\ Ax = b, \ x \geq 0 & \lambda^T A \leq c^T \end{array}$$

- What to show that the **optimal value** (when finite) **as a function of b** is
- What to show that the **optimal value** (when finite) **as a function of c** is

Global Dependency On b, c

$$\begin{array}{ll} (\mathcal{P}) \min c^T x & (\mathcal{D}) \max \lambda^T b \\ Ax = b, \ x \geq 0 & \lambda^T A \leq c^T \end{array}$$

- What to show that the **optimal value** (when finite) **as a function of b** is piecewise linear and **convex**
- What to show that the **optimal value** (when finite) **as a function of c** is piecewise linear and **concave**


Convex and Concave Functions

Definition

$f : X \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ is **convex** if X is a convex set and

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y), \quad \forall x, y \in X \text{ and } \lambda \in [0, 1]. \quad (6)$$

A function is **concave** if $-f$ is convex.




Convex and Concave Functions

Definition

$f : X \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ is **convex** if X is a convex set and

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y), \quad \forall x, y \in X \text{ and } \lambda \in [0, 1]. \quad (6)$$

A function is **concave** if $-f$ is convex.



Equivalent definition in terms of **epigraph**:

$$\text{epi}(f) = \{(x, t) \in X \times \mathbb{R} : t \geq f(x)\} \quad (7)$$

f is **convex** if and only if $\text{epi}(f)$ is a **convex** set.

Global Dependency On b

- Let $P(b) := \{x \in \mathbb{R}^n : Ax = b, x \geq 0\}$ denote the feasible set of the primal
- Let $S := \{b \in \mathbb{R}^m : P(b) \neq \emptyset\}$: right-hand-side values that yield a feasible primal
- Let $p^*(b)$ denote the optimal objective; assume $p^*(b) > -\infty$ (i.e., dual is feasible)

Global Dependency On b

- Let $P(b) := \{x \in \mathbb{R}^n : Ax = b, x \geq 0\}$ denote the feasible set of the primal
- Let $S := \{b \in \mathbb{R}^m : P(b) \neq \emptyset\}$: right-hand-side values that yield a feasible primal
- Let $p^*(b)$ denote the optimal objective; assume $p^*(b) > -\infty$ (i.e., dual is feasible)
- To argue: $p^* : S \rightarrow \mathbb{R}$ is a **piecewise linear and convex** function of b

Global Dependency On b

- Let $P(b) := \{x \in \mathbb{R}^n : Ax = b, x \geq 0\}$ denote the feasible set of the primal
- Let $S := \{b \in \mathbb{R}^m : P(b) \neq \emptyset\}$: right-hand-side values that yield a feasible primal
- Let $p^*(b)$ denote the optimal objective; assume $p^*(b) > -\infty$ (i.e., dual is feasible)
- To argue: $p^* : S \rightarrow \mathbb{R}$ is a **piecewise linear and convex** function of b

Proof. *Is S a convex set?*

Global Dependency On b

- Let $P(b) := \{x \in \mathbb{R}^n : Ax = b, x \geq 0\}$ denote the feasible set of the primal
- Let $S := \{b \in \mathbb{R}^m : P(b) \neq \emptyset\}$: right-hand-side values that yield a feasible primal
- Let $p^*(b)$ denote the optimal objective; assume $p^*(b) > -\infty$ (i.e., dual is feasible)
- To argue: $p^* : S \rightarrow \mathbb{R}$ is a **piecewise linear and convex** function of b

Proof. *Is S a convex set?*


- Strong duality: $p^*(b) = \min\{c^T x : Ax = b, x \geq 0\} = \max\{\lambda^T b : \lambda^T A \leq c^T\}$

Global Dependency On b

- Let $P(b) := \{x \in \mathbb{R}^n : Ax = b, x \geq 0\}$ denote the feasible set of the primal
- Let $S := \{b \in \mathbb{R}^m : P(b) \neq \emptyset\}$: right-hand-side values that yield a feasible primal
- Let $p^*(b)$ denote the optimal objective; assume $p^*(b) > -\infty$ (i.e., dual is feasible)
- To argue: $p^* : S \rightarrow \mathbb{R}$ is a **piecewise linear and convex** function of b

Proof. *Is S a convex set?*

- Strong duality: $p^*(b) = \min\{c^T x : Ax = b, x \geq 0\} = \max\{\lambda^T b : \lambda^T A \leq c^T\}$
- If $\lambda^1, \lambda^2, \dots, \lambda^r$ are the extreme points of D , then: $p^*(b) = \max_{i=1, \dots, r} b^T \lambda^i, \forall b \in S$




Global Dependency On b

- Let $P(b) := \{x \in \mathbb{R}^n : Ax = b, x \geq 0\}$ denote the feasible set of the primal
- Let $S := \{b \in \mathbb{R}^m : P(b) \neq \emptyset\}$: right-hand-side values that yield a feasible primal
- Let $p^*(b)$ denote the optimal objective; assume $p^*(b) > -\infty$ (i.e., dual is feasible)
- To argue: $p^* : S \rightarrow \mathbb{R}$ is a **piecewise linear and convex** function of b

Proof. *Is S a convex set?*


- Strong duality: $p^*(b) = \min\{c^T x : Ax = b, x \geq 0\} = \max\{\lambda^T b : \lambda^T A \leq c^T\}$
- If $\lambda^1, \lambda^2, \dots, \lambda^r$ are the extreme points of D , then: $p^*(b) = \max_{i=1, \dots, r} b^T \lambda^i, \forall b \in S$



How to prove $p^(b)$ convex?*

Global Dependency On b - Implications


$$p^*(b) = \min\{c^T x : Ax = b, x \geq 0\} = \max\{\lambda^T b : \lambda^T A \leq c^T\}$$



- At any \bar{b} where p^* is differentiable, λ^* is the gradient of p^*

Global Dependency On b - Implications


$$p^*(b) = \min\{c^T x : Ax = b, x \geq 0\} = \max\{\lambda^T b : \lambda^T A \leq c^T\}$$



- At any \bar{b} where p^* is differentiable, λ^* **is the gradient** of p^*
- λ_i^* acts as a **marginal cost** or **shadow price** for the i -th constraint r.h.s. b_i
- λ_i allows estimating **exact change in p^* in a range around \bar{b}_i**

Global Dependency On b - Implications

$$p^*(b) = \min\{c^T x : Ax = b, x \geq 0\} = \max\{\lambda^T b : \lambda^T A \leq c^T\}$$




- At any \bar{b} where p^* is differentiable, λ^* is the gradient of p^*
- λ_i^* acts as a **marginal cost** or **shadow price** for the i -th constraint r.h.s. b_i
- λ_i allows estimating **exact change in p^* in a range around \bar{b}_i**
- Modern solvers give **direct access to λ_i^* and the range**

Gurobipy: for constraint c , the attribute $c.Pi$ is λ_i^* and the range is from $c.SARHSLow$ to $c.SARHSUp$

Global Dependency On b - Implications


$$p^*(b) = \min\{c^T x : Ax = b, x \geq 0\} = \max\{\lambda^T b : \lambda^T A \leq c^T\}$$



- At b where p^* is **not** differentiable, several λ^i are optimal
- All such λ^i are valid **subgradients** of p^*

Global Dependency On b - Implications

$$p^*(b) = \min\{c^T x : Ax = b, x \geq 0\} = \max\{\lambda^T b : \lambda^T A \leq c^T\}$$



- At b where p^* is **not** differentiable, several λ^i are optimal
- All such λ^i are valid **subgradients** of p^*

Definition (Subgradient.)

$f : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ convex function. A vector $g \in \mathbb{R}^n$ is a **subgradient** of f at $\bar{x} \in S$ if

$$f(x) \geq f(\bar{x}) + g^T(x - \bar{x}), \quad \forall x \in S.$$

Global Dependency On c

- Let $d^*(c)$ denote optimal value as function of c ; assume $d^*(c) > -\infty$
- To argue: $d^*(c)$ is a **piecewise linear and concave** function of c

Global Dependency On c

- Let $d^*(c)$ denote optimal value as function of c ; assume $d^*(c) > -\infty$
- To argue: $d^*(c)$ is a **piecewise linear and concave** function of c
- $d^*(c) = \min\{c^T x : Ax = b, x \geq 0\} = \max\{\lambda^T b : \lambda^T A \leq c^T\}$
- Can apply same arguments because d^* is the optimal value of the dual

Global Dependency On c

- Let $d^*(c)$ denote optimal value as function of c ; assume $d^*(c) > -\infty$
- To argue: $d^*(c)$ is a **piecewise linear and concave** function of c
- $d^*(c) = \min\{c^T x : Ax = b, x \geq 0\} = \max\{\lambda^T b : \lambda^T A \leq c^T\}$
- Can apply same arguments because d^* is the optimal value of the dual
- $d^*(c)$ is a **concave** function of c on the set $T := \{c : d^*(c) > -\infty\}$
- If for some c the LP has a **unique** optimal solution x^* , then d^* is linear in the vicinity of c and its gradient is x^* .

Global Dependency On c

- Let $d^*(c)$ denote optimal value as function of c ; assume $d^*(c) > -\infty$
- To argue: $d^*(c)$ is a **piecewise linear and concave** function of c
- $d^*(c) = \min\{c^T x : Ax = b, x \geq 0\} = \max\{\lambda^T b : \lambda^T A \leq c^T\}$
- Can apply same arguments because d^* is the optimal value of the dual
- $d^*(c)$ is a **concave** function of c on the set $T := \{c : d^*(c) > -\infty\}$
- If for some c the LP has a **unique** optimal solution x^* , then d^* is linear in the vicinity of c and its gradient is x^* .
- The optimal primal solution x^* **is a shadow price for the dual constraints**
- x^* remains optimal for a range of change in each objective coefficient c_j
- Modern solvers also allow obtaining the range directly
Gurobi: attributes **SAObjLow** and **SAObjUp** for each decision variable

Signs of Dual Variables Revisited

- There is a direct connection between:
 - the **optimization problem** (max/min)
 - the **constraint type** (\leq , \geq)
 - the **signs of the shadow prices**
- Given two of these, can figure out the third one!
- *What is the sign of the shadow price for a ...*
 - \leq constraint in a **minimization** problem ?
 - \geq constraint in a **minimization** problem ?
 - \leq constraint in a **maximization** problem ?
 - \geq constraint in a **maximization** problem ?
- *What is the dependency of the optimal objective on the r.h.s. of a ...*
 - \leq constraint in a **minimization** problem ?
 - \geq constraint in a **minimization** problem ?
 - \leq constraint in a **maximization** problem ?
 - \geq constraint in a **maximization** problem ?

Signs of Dual Variables Revisited

- There is a direct connection between:
 - **the optimization problem** (max/min)
 - the **constraint type** (\leq , \geq)
 - **the signs of the shadow prices**
- Given two of these, can figure out the third one!

Network Revenue Management

- Airline revenue management (“yield management”): setting **booking limits** to control how many tickets of each type are sold

Network Revenue Management

- Airline revenue management (“yield management”): setting **booking limits** to control how many tickets of each type are sold
- Airline is planning operations for a specific day in the future

Network Revenue Management

- Airline revenue management (“yield management”): setting **booking limits** to control how many tickets of each type are sold
- Airline is planning operations for a specific day in the future
- Airline operates a set F of direct flights in its (hub-and-spoke) network

Network Revenue Management

- Airline revenue management (“yield management”): setting **booking limits** to control how many tickets of each type are sold
- Airline is planning operations for a specific day in the future
- Airline operates a set F of direct flights in its (hub-and-spoke) network
- For each flight leg $f \in F$, we know the capacity of the aircraft c_f

Network Revenue Management

- Airline revenue management (“yield management”): setting **booking limits** to control how many tickets of each type are sold
- Airline is planning operations for a specific day in the future
- Airline operates a set F of direct flights in its (hub-and-spoke) network
- For each flight leg $f \in F$, we know the capacity of the aircraft c_f
- The airline can offer a large number of “products” (i.e., itineraries) I :
 - each itinerary refers to an origin-destination-fare class combination
 - each itinerary i has a price r_i that is fixed
 - for each itinerary, the airline estimates the demand d_i
 - each itinerary requires a seat on several flight legs operated by the airline

Network Revenue Management

- Airline revenue management (“yield management”): setting **booking limits** to control how many tickets of each type are sold
- Airline is planning operations for a specific day in the future
- Airline operates a set F of direct flights in its (hub-and-spoke) network
- For each flight leg $f \in F$, we know the capacity of the aircraft c_f
- The airline can offer a large number of “products” (i.e., itineraries) I :
 - each itinerary refers to an origin-destination-fare class combination
 - each itinerary i has a price r_i that is fixed
 - for each itinerary, the airline estimates the demand d_i
 - each itinerary requires a seat on several flight legs operated by the airline
- Requirements: $A \in \{0, 1\}^{F \times I}$ with $A_{f,i} = 1 \Leftrightarrow$ itinerary i needs seat on flight leg f

		Itinerary 1	Itinerary 2	...	Itinerary $ I $
Resource matrix A :	Flight leg 1	1	0	...	1
	Flight leg 2	0	1	...	0
	\vdots	\vdots	\vdots	\vdots	\vdots
	Flight leg $ F $	1	1	...	0

Network Revenue Management

- Airline revenue management (“yield management”): setting **booking limits** to control how many tickets of each type are sold
- Airline is planning operations for a specific day in the future
- Airline operates a set F of direct flights in its (hub-and-spoke) network
- For each flight leg $f \in F$, we know the capacity of the aircraft c_f
- The airline can offer a large number of “products” (i.e., itineraries) I :
 - each itinerary refers to an origin-destination-fare class combination
 - each itinerary i has a price r_i that is fixed
 - for each itinerary, the airline estimates the demand d_i
 - each itinerary requires a seat on several flight legs operated by the airline
- Requirements: $A \in \{0, 1\}^{F \times I}$ with $A_{f,i} = 1 \Leftrightarrow$ itinerary i needs seat on flight leg f

		Itinerary 1	Itinerary 2	...	Itinerary $ I $
Resource matrix A :	Flight leg 1	1	0	...	1
	Flight leg 2	0	1	...	0
	\vdots	\vdots	\vdots	\vdots	\vdots
	Flight leg $ F $	1	1	...	0

- Goal: decide how many itineraries of each type to sell to maximize revenue

Network Revenue Management

- Let x_i denote the number of itineraries of type i that the airline plans to sell, and let x be the vector with components x_i

Network Revenue Management

- Let x_i denote the number of itineraries of type i that the airline plans to sell, and let x be the vector with components x_i
- The problem can be formulated as follows:

$$\max_{x \in \mathbb{R}^I} \left\{ r^T x : Ax \leq c, x \leq d \right\}$$

Network Revenue Management

- Let x_i denote the number of itineraries of type i that the airline plans to sell, and let x be the vector with components x_i
- The problem can be formulated as follows:

$$\max_{x \in \mathbb{R}^I} \left\{ r^T x : Ax \leq c, x \leq d \right\}$$

- $Ax \leq c$ capture the constraints on plane capacity
- $x \leq d$ states that the planned sales cannot exceed the demand
- In practice, an approach that includes **all possible itineraries** encounters challenges

Network Revenue Management

- Let x_i denote the number of itineraries of type i that the airline plans to sell, and let x be the vector with components x_i
- The problem can be formulated as follows:

$$\max_{x \in \mathbb{R}^I} \left\{ r^T x : Ax \leq c, x \leq d \right\}$$

- $Ax \leq c$ capture the constraints on plane capacity
- $x \leq d$ states that the planned sales cannot exceed the demand
- In practice, an approach that includes **all possible itineraries** encounters challenges
 - gargantuan LP
 - poor demand estimates for some itineraries

Network Revenue Management

- Let x_i denote the number of itineraries of type i that the airline plans to sell, and let x be the vector with components x_i
- The problem can be formulated as follows:

$$\max_{x \in \mathbb{R}^I} \left\{ r^T x : Ax \leq c, x \leq d \right\}$$

- $Ax \leq c$ capture the constraints on plane capacity
- $x \leq d$ states that the planned sales cannot exceed the demand
- In practice, an approach that includes **all possible itineraries** encounters challenges
 - gargantuan LP
 - poor demand estimates for some itineraries
- To sell “exotic itineraries”, use the **shadow prices for the capacity constraints**

Network Revenue Management

- Let x_i denote the number of itineraries of type i that the airline plans to sell, and let x be the vector with components x_i
- The problem can be formulated as follows:

$$\max_{x \in \mathbb{R}^I} \left\{ r^T x : Ax \leq c, x \leq d \right\}$$

- $Ax \leq c$ capture the constraints on plane capacity
- $x \leq d$ states that the planned sales cannot exceed the demand
- In practice, an approach that includes **all possible itineraries** encounters challenges
 - gargantuan LP
 - poor demand estimates for some itineraries
- To sell “exotic itineraries”, use the **shadow prices for the capacity constraints**
 - $\lambda \in \mathbb{R}^F$: dual variables for capacity constraints $Ax \leq c$
 - At optimality, p_f is marginal revenue lost if airline loses one seat on flight f

Network Revenue Management

- Let x_i denote the number of itineraries of type i that the airline plans to sell, and let x be the vector with components x_i
- The problem can be formulated as follows:

$$\max_{x \in \mathbb{R}_+^I} \left\{ r^T x : Ax \leq c, x \leq d \right\}$$

- $Ax \leq c$ capture the constraints on plane capacity
- $x \leq d$ states that the planned sales cannot exceed the demand
- In practice, an approach that includes **all possible itineraries** encounters challenges
 - gargantuan LP
 - poor demand estimates for some itineraries
- To sell “exotic itineraries”, use the **shadow prices for the capacity constraints**
 - $\lambda \in \mathbb{R}^F$: dual variables for capacity constraints $Ax \leq c$
 - At optimality, p_f is marginal revenue lost if airline loses one seat on flight f
 - For an “exotic” itinerary that requires seats on several flights $f \in E$, the **minimum price** to charge is given by the sum of the shadow prices, $\sum_{f \in E} p_f$

Network Revenue Management

- Let x_i denote the number of itineraries of type i that the airline plans to sell, and let x be the vector with components x_i
- The problem can be formulated as follows:

$$\max_{x \in \mathbb{R}_+^I} \left\{ r^T x : Ax \leq c, x \leq d \right\}$$

- $Ax \leq c$ capture the constraints on plane capacity
- $x \leq d$ states that the planned sales cannot exceed the demand
- In practice, an approach that includes **all possible itineraries** encounters challenges
 - gargantuan LP
 - poor demand estimates for some itineraries
- To sell “exotic itineraries”, use the **shadow prices for the capacity constraints**
 - $\lambda \in \mathbb{R}^F$: dual variables for capacity constraints $Ax \leq c$
 - At optimality, p_f is marginal revenue lost if airline loses one seat on flight f
 - For an “exotic” itinerary that requires seats on several flights $f \in E$, the **minimum price** to charge is given by the sum of the shadow prices, $\sum_{f \in E} p_f$
- **Bid-price heuristic** in network revenue management
- Broader principle of how to price “products” through resource usage/cost