

# CME 307 / MS&E 311: Optimization

## Duality

Professor Udell

Management Science and Engineering  
Stanford

May 1, 2023

## Announcements

- ▶ meet with course staff to discuss project this week or next (see Ed)
- ▶ project 1 due this Friday 5/5

# Outline

Duality

Lagrange duality

# Duality

## Definition (Dual space)

The **dual**  $\mathcal{X}^*$  of a vector space  $\mathcal{X}$  is the set of linear functionals on  $\mathcal{X}$ .

so if  $x \in \mathcal{X}$  and you see someone write

$$w^T x, \quad \langle w, x \rangle, \quad \text{or} \quad w \cdot x$$

you know that  $w \in \mathcal{X}^*$  is a dual vector

# Duality

## Definition (Dual space)

The **dual**  $\mathcal{X}^*$  of a vector space  $\mathcal{X}$  is the set of linear functionals on  $\mathcal{X}$ .

so if  $x \in \mathcal{X}$  and you see someone write

$$w^T x, \quad \langle w, x \rangle, \quad \text{or} \quad w \cdot x$$

you know that  $w \in \mathcal{X}^*$  is a dual vector

notation: solution to optimization problem  $x^*$  vs dual space  $\mathcal{X}^*$

## Careful of units!

**example 1:** suppose  $y_i = w^T x_i$  where

$$x_i = \begin{bmatrix} \text{heart rate} \\ \text{blood pressure} \\ \text{age} \end{bmatrix}, \quad \text{with units} \quad \begin{bmatrix} \text{bpm} \\ \text{mmHg} \\ \text{years} \end{bmatrix}$$

and  $y_i$  is duration of stay in hospital (units: days)

## Careful of units!

**example 1:** suppose  $y_i = w^T x_i$  where

$$x_i = \begin{bmatrix} \text{heart rate} \\ \text{blood pressure} \\ \text{age} \end{bmatrix}, \quad \text{with units} \quad \begin{bmatrix} \text{bpm} \\ \text{mmHg} \\ \text{years} \end{bmatrix}$$

and  $y_i$  is duration of stay in hospital (units: days)

then  $w$  has units of

$$\begin{bmatrix} \text{days/bpm} \\ \text{days/mmHg} \\ \text{days/year} \end{bmatrix}$$

## Careful of units!

**example 2:**  $f(x) = \sum_{i=1}^n \left( 1 + \exp \left( \underbrace{y_i w^T x_i}_{\text{input must be a scalar!}} \right) \right)$



## Careful of units!

**example 2:**  $f(x) = \sum_{i=1}^n \left( 1 + \exp \left( \underbrace{y_i w^T x_i}_{\text{input must be a scalar!}} \right) \right)$

**example 3:** if  $x \in \mathcal{X}$ , gradient is a linear function on  $\mathcal{X} \implies \nabla f(x_0) \in \mathcal{X}^*$

$$f(x) \approx f(x_0) + \nabla f(x_0)^T (x - x_0),$$

so gradient descent stepsize  $t$  has units

$$x^{k+1} = x^k - t \nabla f(x^k)$$

e.g.,  $x$  (meters  $m$ ),  $\nabla f(x)$  ( $m^{-1}$ ), and  $t$  ( $m^2$ )

## Careful of units!

**example 2:**  $f(x) = \sum_{i=1}^n \left( 1 + \exp \left( \underbrace{y_i w^T x_i}_{\text{input must be a scalar!}} \right) \right)$

**example 3:** if  $x \in \mathcal{X}$ , gradient is a linear function on  $\mathcal{X} \implies \nabla f(x_0) \in \mathcal{X}^*$

$$f(x) \approx f(x_0) + \nabla f(x_0)^T (x - x_0),$$

so gradient descent stepsize  $t$  has units

$$x^{k+1} = x^k - t \nabla f(x^k)$$

e.g.,  $x$  (meters  $m$ ),  $\nabla f(x)$  ( $m^{-1}$ ), and  $t$  ( $m^2$ )

- ▶ no wonder it's hard to choose the stepsize!
- ▶ basic recommendation: standardize your data

## Dual of function space

- ▶  $f : [0, 1] \rightarrow \mathbf{R}$  is a function
- ▶  $f(x)$  is a linear function of  $f$ , for any  $x$ :

$$(f + g)(x) = f(x) + g(x), \quad (cf)(x) = cf(x)$$

- ▶ so is any integral:

$$\int_0^1 f(x) d\mu(x)$$

$\implies$  the dual of the space of functions on  $[0, 1]$  is the space of measures on  $[0, 1]$

## Dual norm

### Definition (Dual norm)

The **dual norm** of a norm  $\| \cdot \|$  is

$$\|w\|_* = \sup_{\|x\| \leq 1} \langle w, x \rangle$$

equivalently,  $\|w\|_* = \sup_x \frac{\langle w, x \rangle}{\|x\|}$

## Dual norm

### Definition (Dual norm)

The **dual norm** of a norm  $\|\cdot\|$  is

$$\|w\|_* = \sup_{\|x\| \leq 1} \langle w, x \rangle$$

equivalently,  $\|w\|_* = \sup_x \frac{\langle w, x \rangle}{\|x\|}$

**example:**  $\ell_1$  norm dual is  $\ell_\infty$  norm

$$\|w\|_1 = \sum_{i=1}^n |w_i|, \quad \|w\|_\infty = \max_{i=1, \dots, n} |w_i|$$

## Dual norm

### Definition (Dual norm)

The **dual norm** of a norm  $\| \cdot \|$  is

$$\|w\|_* = \sup_{\|x\| \leq 1} \langle w, x \rangle$$

equivalently,  $\|w\|_* = \sup_x \frac{\langle w, x \rangle}{\|x\|}$

**example:**  $\ell_1$  norm dual is  $\ell_\infty$  norm

$$\|w\|_1 = \sum_{i=1}^n |w_i|, \quad \|w\|_\infty = \max_{i=1, \dots, n} |w_i|$$

**example:**  $\ell_2$  norm dual is  $\ell_2$  norm  $\implies \ell_2$  is **self-dual**

## Dual norm

### Definition (Dual norm)

The **dual norm** of a norm  $\|\cdot\|$  is

$$\|w\|_* = \sup_{\|x\| \leq 1} \langle w, x \rangle$$

equivalently,  $\|w\|_* = \sup_x \frac{\langle w, x \rangle}{\|x\|}$

**example:**  $\ell_1$  norm dual is  $\ell_\infty$  norm

$$\|w\|_1 = \sum_{i=1}^n |w_i|, \quad \|w\|_\infty = \max_{i=1, \dots, n} |w_i|$$

**example:**  $\ell_2$  norm dual is  $\ell_2$  norm  $\implies \ell_2$  is **self-dual**

**example:** for  $f : [0, 1] \rightarrow \mathbf{R}$ , if  $\|f\| = \sup_{x \in [0, 1]} |f(x)|$ ,

$$\|\mu\|_* = \sup_{\|f\| \leq 1} \int_0^1 f(x) d\mu(x) = \int_0^1 d|\mu|(x)$$

## Self-dual norms

given primal space  $\mathcal{X}$

- ▶ dual vector is a linear functional  $w(x)$  on  $x \in \mathcal{X}$
- ▶ we should define the dual norm on  $\mathcal{X}^*$  as

$$\sup_{x \in \mathcal{X}, \|x\| \leq 1} w(x)$$

- ▶ but instead we used the inner product  $\langle w, x \rangle$ . why?



## Self-dual norms

given primal space  $\mathcal{X}$

- ▶ dual vector is a linear functional  $w(x)$  on  $x \in \mathcal{X}$
- ▶ we should define the dual norm on  $\mathcal{X}^*$  as

$$\sup_{x \in \mathcal{X}, \|x\| \leq 1} w(x)$$

- ▶ but instead we used the inner product  $\langle w, x \rangle$ . why?

### Theorem (Riesz representation)

*Suppose  $\mathcal{X} = H$  is a Hilbert (inner product) space. For any linear functional  $\phi \in \mathcal{X}^*$ , there is a unique vector  $w \in H$  so that  $w(x) = \langle w, x \rangle$  for all  $x \in \mathcal{X} = H$ . Moreover,  $\|w\|_* = \|w\|$ .*

$\|\cdot\|$  is self-dual  $\iff \|\cdot\|$  is induced by an inner product

## Self-dual norms

given primal space  $\mathcal{X}$

- ▶ dual vector is a linear functional  $w(x)$  on  $x \in \mathcal{X}$
- ▶ we should define the dual norm on  $\mathcal{X}^*$  as

$$\sup_{x \in \mathcal{X}, \|x\| \leq 1} w(x)$$

- ▶ but instead we used the inner product  $\langle w, x \rangle$ . why?

### Theorem (Riesz representation)

*Suppose  $\mathcal{X} = H$  is a Hilbert (inner product) space. For any linear functional  $\phi \in \mathcal{X}^*$ , there is a unique vector  $w \in H$  so that  $w(x) = \langle w, x \rangle$  for all  $x \in \mathcal{X} = H$ . Moreover,  $\|w\|_* = \|w\|$ .*

$\|\cdot\|$  is self-dual  $\iff \|\cdot\|$  is induced by an inner product

**example:**  $\ell_2$  norm is self-dual, induced by the inner product

$$\langle w, x \rangle = w^T x$$

## Conjugate of linear operator

given  $x \in \mathbf{R}^n$ ,  $w \in \mathbf{R}^m$ , and  $A \in \mathbf{R}^{m \times n}$ , conjugate of  $A$  is the linear operator  $A^*$  defined so that

$$\langle A^* w, x \rangle = \langle w, Ax \rangle$$

## Conjugate of linear operator

given  $x \in \mathbf{R}^n$ ,  $w \in \mathbf{R}^m$ , and  $A \in \mathbf{R}^{m \times n}$ , conjugate of  $A$  is the linear operator  $A^*$  defined so that

$$\langle A^* w, x \rangle = \langle w, Ax \rangle$$

**example:**  $x \in \mathbf{R}^n$ ,  $A \in \mathbf{R}^{m \times n}$  defined by

$$Ax = \begin{bmatrix} x_{i_1} \\ \vdots \\ x_{i_m} \end{bmatrix}$$

then  $A^* \in \mathbf{R}^{n \times m}$  satisfies

$$\langle A^* w, x \rangle = \langle w, Ax \rangle = \sum_{j=1}^m w_j x_{i_j},$$

so  $A^*$  creates a sparse vector from  $w$  with

$$(A^* w)_{i_j} = w_j$$

## Fenchel dual

### Definition (Fenchel dual)

The **Fenchel dual** of a function  $f : \mathcal{X} \rightarrow \mathbf{R}$  is

$$f^*(w) = \sup_{x \in \mathcal{X}} \langle w, x \rangle - f(x)$$

also called the **conjugate function**. draw picture!

## Fenchel dual

### Definition (Fenchel dual)

The **Fenchel dual** of a function  $f : \mathcal{X} \rightarrow \mathbf{R}$  is

$$f^*(w) = \sup_{x \in \mathcal{X}} \langle w, x \rangle - f(x)$$

also called the **conjugate function**. draw picture!

**example:**  $f(x) = \|x\|_1, x \in \mathbf{R}^n$

$$f^*(w) = \sup_{x \in \mathbf{R}^n} \langle w, x \rangle - \|x\|_1 = \begin{cases} 0 & \|w\|_\infty \leq 1 \\ \infty & \text{otherwise} \end{cases}$$

$\implies$  fenchel dual of  $\ell_1$  norm is indicator of  $\ell_\infty$  ball

# Biconjugate

## Definition (Biconjugate)

The **biconjugate** of a function  $f : \mathcal{X} \rightarrow \mathbf{R}$  is

$$f^{**}(x) = \sup_{w \in \mathcal{X}^*} \langle w, x \rangle - f^*(w)$$

- ▶ for convex  $f : \mathbf{R} \rightarrow \mathbf{R}$ ,  $f^{**} = f$
- ▶ for nonconvex  $f$ ,  $f^{**}$  is convex hull of  $f$

$\implies$  biconjugate is a convexification operation

## Biconjugate

### Definition (Biconjugate)

The **biconjugate** of a function  $f : \mathcal{X} \rightarrow \mathbf{R}$  is

$$f^{**}(x) = \sup_{w \in \mathcal{X}^*} \langle w, x \rangle - f^*(w)$$

► for convex  $f : \mathbf{R} \rightarrow \mathbf{R}$ ,  $f^{**} = f$

► for nonconvex  $f$ ,  $f^{**}$  is convex hull of  $f$

$\implies$  biconjugate is a convexification operation

**example:** consider  $f : \mathbf{R} \rightarrow \mathbf{R}$  defined by

$$f(x) = \begin{cases} 0 & x \in \{-1, 1\} \\ \infty & \text{otherwise} \end{cases}$$

what is  $f_*$ ?  $f^{**}$ ?



# Outline

Duality

Lagrange duality

## Why duality?

- ▶ certify optimality
  - ▶ turn  $\forall$  into  $\exists$
  - ▶ use dual lower bound to derive stopping conditions
- ▶ new algorithms based on the dual
  - ▶ solve dual, then recover primal solution

## Warmup: Farkas lemma

### Theorem (Farkas lemma)

*Given  $A \in \mathbf{R}^{m \times n}$  and  $b \in \mathbf{R}^m$ , exactly one of the following is true:*

- ▶ *there exists  $x \in \mathbf{R}^n$  so that  $Ax = b$  and  $x \geq 0$*
- ▶ *there exists  $y \in \mathbf{R}^m$  so that  $A^T y \geq 0$  and  $\langle b, y \rangle < 0$*

$\implies$  can efficiently certify infeasibility of a linear program

## Warmup: Farkas lemma

### Theorem (Farkas lemma)

Given  $A \in \mathbf{R}^{m \times n}$  and  $b \in \mathbf{R}^m$ , exactly one of the following is true:

- ▶ there exists  $x \in \mathbf{R}^n$  so that  $Ax = b$  and  $x \geq 0$
- ▶ there exists  $y \in \mathbf{R}^m$  so that  $A^T y \geq 0$  and  $\langle b, y \rangle < 0$

$\implies$  can efficiently certify infeasibility of a linear program

**proof:** suppose we have  $x \in \mathbf{R}^n$  so that  $Ax = b$  and  $x \geq 0$ .  
then for any  $y \in \mathbf{R}^m$ ,

$$\begin{aligned} 0 &= \langle y, b - Ax \rangle = \langle y, b \rangle - \langle A^T y, x \rangle \\ \langle y, b \rangle &= \langle A^T y, x \rangle \end{aligned}$$

so if  $A^T y \geq 0$ , then use  $x \geq 0$  to conclude  $\langle y, b \rangle \geq 0$ .

## Warmup: Farkas lemma

### Theorem (Farkas lemma)

Given  $A \in \mathbf{R}^{m \times n}$  and  $b \in \mathbf{R}^m$ , exactly one of the following is true:

- ▶ there exists  $x \in \mathbf{R}^n$  so that  $Ax = b$  and  $x \geq 0$
- ▶ there exists  $y \in \mathbf{R}^m$  so that  $A^T y \geq 0$  and  $\langle b, y \rangle < 0$

$\implies$  can efficiently certify infeasibility of a linear program

**proof:** suppose we have  $x \in \mathbf{R}^n$  so that  $Ax = b$  and  $x \geq 0$ .  
then for any  $y \in \mathbf{R}^m$ ,

$$\begin{aligned} 0 &= \langle y, b - Ax \rangle = \langle y, b \rangle - \langle A^T y, x \rangle \\ \langle y, b \rangle &= \langle A^T y, x \rangle \end{aligned}$$

so if  $A^T y \geq 0$ , then use  $x \geq 0$  to conclude  $\langle y, b \rangle \geq 0$ .

(opposite direction is similar)

## Lagrange duality

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

if  $x$  is feasible, then  $Ax = b$ , so  $\langle y, Ax - b \rangle = 0$ .

## Lagrange duality

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

if  $x$  is feasible, then  $Ax = b$ , so  $\langle y, Ax - b \rangle = 0$ .

define the **Lagrangian**

$$\mathcal{L}(x, y) := f(x) - \langle y, b - Ax \rangle$$

## Lagrange duality

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

if  $x$  is feasible, then  $Ax = b$ , so  $\langle y, Ax - b \rangle = 0$ .

define the **Lagrangian**

$$\begin{aligned} \mathcal{L}(x, y) &:= f(x) - \langle y, b - Ax \rangle \\ p^* &= \inf_{x: Ax=b} \mathcal{L}(x, y) \geq \inf_x \mathcal{L}(x, y) \end{aligned}$$



## Lagrange duality

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax = b : \quad \text{dual } y \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

if  $x$  is feasible, then  $Ax = b$ , so  $\langle y, Ax - b \rangle = 0$ .

define the **Lagrangian**

$$\begin{aligned} \mathcal{L}(x, y) &:= f(x) - \langle y, b - Ax \rangle \\ p^* &= \inf_{x: Ax=b} \mathcal{L}(x, y) \geq \inf_x \mathcal{L}(x, y) \\ &= \inf_x f(x) + \langle y, -b + Ax \rangle \\ &= \langle y, -b \rangle + \inf_x \left( f(x) + \langle A^T y, x \rangle \right) \\ &= \langle y, -b \rangle - \sup_x \left( \langle -A^T y, x \rangle - f(x) \right) \\ &= \langle y, -b \rangle - f^*(-A^T y) = g(y) \end{aligned}$$

$g(y)$  is called the **dual function**

## Lagrange duality

inequality holds for any  $y \in \mathbf{R}^m$ , so we have proved **weak duality**

$$\begin{aligned} p^* &\geq g(y) \quad \forall y \in \mathbf{R}^m \\ &\geq \underbrace{\sup_y g(y)}_{\mathcal{D}} =: d^* \end{aligned} \tag{1}$$

dual optimal value  $d^* \leq p^*$

## Strong duality

### Definition (Duality gap)

The **duality gap** for a primal-dual pair  $(x, y)$  is  $f(x) - g(y)$

by weak duality, duality gap is always nonnegative

## Strong duality

### Definition (Duality gap)

The **duality gap** for a primal-dual pair  $(x, y)$  is  $f(x) - g(y)$

by weak duality, duality gap is always nonnegative

### Definition (Strong duality)

A primal-dual pair  $(x^*, y^*)$  satisfies **strong duality** if

$$p^* = d^* \iff f(x^*) - g(y^*) = 0$$

## Strong duality

### Definition (Duality gap)

The **duality gap** for a primal-dual pair  $(x, y)$  is  $f(x) - g(y)$

by weak duality, duality gap is always nonnegative

### Definition (Strong duality)

A primal-dual pair  $(x^*, y^*)$  satisfies **strong duality** if

$$p^* = d^* \iff f(x^*) - g(y^*) = 0$$

strong duality holds

- ▶ for feasible LPs (pf later)
- ▶ for convex problems under **constraint qualification** aka **Slater's condition**. feasible region has an **interior point**  $x$  so that all inequality constraints hold strictly

strong duality fails if either ( $\mathcal{P}$ ) or ( $\mathcal{D}$ ) is infeasible or unbounded

## Lagrange duality with inequality constraints

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax \leq b : \quad y \geq 0 \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

## Lagrange duality with inequality constraints

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax \leq b : \quad y \geq 0 \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

to construct Lagrangian  $\mathcal{L}(x, y) = f(x) - \langle y, b - Ax \rangle$ , ensure value is **better** (lower) when  $x$  and  $y$  are feasible

## Lagrange duality with inequality constraints

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax \leq b : \quad y \geq 0 \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

to construct Lagrangian  $\mathcal{L}(x, y) = f(x) - \langle y, b - Ax \rangle$ , ensure value is **better** (lower) when  $x$  and  $y$  are feasible

$$\mathcal{L}(x, y) := f(x) - \langle y, b - Ax \rangle$$



## Lagrange duality with inequality constraints

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax \leq b : \quad y \geq 0 \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

to construct Lagrangian  $\mathcal{L}(x, y) = f(x) - \langle y, b - Ax \rangle$ , ensure value is **better** (lower) when  $x$  and  $y$  are feasible

$$\begin{aligned} \mathcal{L}(x, y) &:= f(x) - \langle y, b - Ax \rangle \\ p^* &\geq \inf_{x \text{ feas}} f(x) - \langle y, b - Ax \rangle \\ &\geq \inf_x f(x) - \langle y, b - Ax \rangle \\ &= \langle y, b \rangle - f^*(-A^*y) =: g(y) \end{aligned}$$

## Lagrange duality with inequality constraints

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & f(x) \\ \text{subject to} & Ax \leq b : \quad y \geq 0 \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

to construct Lagrangian  $\mathcal{L}(x, y) = f(x) - \langle y, b - Ax \rangle$ , ensure value is **better** (lower) when  $x$  and  $y$  are feasible

$$\begin{aligned} \mathcal{L}(x, y) &:= f(x) - \langle y, b - Ax \rangle \\ p^* &\geq \inf_{x \text{ feas}} f(x) - \langle y, b - Ax \rangle \\ &\geq \inf_x f(x) - \langle y, b - Ax \rangle \\ &= \langle y, b \rangle - f^*(-A^*y) =: g(y) \end{aligned}$$

as before, this holds for all  $y$ , so we have weak duality

$$p^* \geq \underbrace{\sup_y g(y)}_{\mathcal{D}} =: d^*$$

## SVM dual

support vector machine: for  $x_i \in \mathbf{R}^n$ ,  $y_i \in \{-1, 1\}$ ,  $i = 1, \dots, m$

$$\text{minimize} \quad \frac{1}{2} \|w\|^2 + 1^T s$$

$$\text{subject to} \quad y_i w^T x_i + s_i \geq 1 \quad i = 1, \dots, m : \quad \alpha \geq 0$$

$$s \geq 0 : \quad \mu \geq 0$$

(SVM)

## SVM dual

support vector machine: for  $x_i \in \mathbf{R}^n$ ,  $y_i \in \{-1, 1\}$ ,  $i = 1, \dots, m$

$$\begin{aligned} &\text{minimize} && \frac{1}{2} \|w\|^2 + 1^T s \\ &\text{subject to} && y_i w^T x_i + s_i \geq 1 \quad i = 1, \dots, m : \quad \alpha \geq 0 \\ &&& s \geq 0 : \quad \mu \geq 0 \end{aligned}$$

(SVM)

Lagrangian: for  $\alpha \geq 0$ ,  $\mu \geq 0$ ,

$$\mathcal{L}(w, s, \alpha, \mu) = \frac{1}{2} \|w\|^2 + 1^T s - \sum_{i=1}^m \alpha_i (y_i w^T x_i + s_i - 1) - \mu^T s$$

► minimize  $\mathcal{L}(w, s, \alpha, \mu)$  over  $w$ :

$$w = \sum_{i=1}^m \alpha_i y_i x_i$$

► minimize  $\mathcal{L}(w, s, \alpha, \mu)$  over  $s \implies \alpha + \mu = 1$

## SVM dual

so simplify:

$$\begin{aligned}g(\alpha) &= \inf_{w,s} \mathcal{L}(w, s, \alpha, 1 - \alpha) \\&= \frac{1}{2} \|w\|^2 - w^T \sum_{i=1}^m \alpha_i y_i x_i + \mathbf{1}^T \alpha \\&= -\frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 + \mathbf{1}^T \alpha\end{aligned}$$

## SVM dual

so simplify:

$$\begin{aligned}g(\alpha) &= \inf_{w,s} \mathcal{L}(w, s, \alpha, 1 - \alpha) \\&= \frac{1}{2} \|w\|^2 - w^T \sum_{i=1}^m \alpha_i y_i x_i + 1^T \alpha \\&= -\frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 + 1^T \alpha\end{aligned}$$

define  $K \in \mathbf{R}^m$  so  $K_{ij} = y_i y_j x_i^T x_j$ . then

$$\left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 = \sum_{i=1}^m \sum_{j=1}^m \alpha_i \alpha_j y_i y_j x_i^T x_j = \alpha^T K \alpha$$

## SVM dual

so simplify:

$$\begin{aligned}g(\alpha) &= \inf_{w,s} \mathcal{L}(w, s, \alpha, 1 - \alpha) \\&= \frac{1}{2} \|w\|^2 - w^T \sum_{i=1}^m \alpha_i y_i x_i + 1^T \alpha \\&= -\frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 + 1^T \alpha\end{aligned}$$

define  $K \in \mathbf{R}^m$  so  $K_{ij} = y_i y_j x_i^T x_j$ . then

$$\left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 = \sum_{i=1}^m \sum_{j=1}^m \alpha_i \alpha_j y_i y_j x_i^T x_j = \alpha^T K \alpha$$

dual problem:

$$\begin{array}{ll}\text{maximize} & -\frac{1}{2} \alpha^T K \alpha + 1^T \alpha \\ \text{subject to} & \alpha \geq 0\end{array} \quad (\text{SVM-dual})$$

## SVM dual

so simplify:

$$\begin{aligned}g(\alpha) &= \inf_{w,s} \mathcal{L}(w, s, \alpha, 1 - \alpha) \\&= \frac{1}{2} \|w\|^2 - w^T \sum_{i=1}^m \alpha_i y_i x_i + 1^T \alpha \\&= -\frac{1}{2} \left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 + 1^T \alpha\end{aligned}$$

define  $K \in \mathbf{R}^m$  so  $K_{ij} = y_i y_j x_i^T x_j$ . then

$$\left\| \sum_{i=1}^m \alpha_i y_i x_i \right\|^2 = \sum_{i=1}^m \sum_{j=1}^m \alpha_i \alpha_j y_i y_j x_i^T x_j = \alpha^T K \alpha$$

dual problem:

$$\begin{array}{ll}\text{maximize} & -\frac{1}{2} \alpha^T K \alpha + 1^T \alpha \\ \text{subject to} & \alpha \geq 0\end{array} \quad (\text{SVM-dual})$$

new solution ideas! coordinate descent on  $\alpha$  (SMO), kernel trick



## Generalize Lagrangian duality

## Generalize Lagrangian duality

- **nonlinear duality:** replace

$$0 \geq Ax - b \quad \text{with} \quad 0 \geq g(x)$$

(harder to derive explicit form for dual problem)

## Generalize Lagrangian duality

- **nonlinear duality:** replace

$$0 \geq Ax - b \quad \text{with} \quad 0 \geq g(x)$$

(harder to derive explicit form for dual problem)

- **conic duality:** for cone  $K$ , replace

$$b - Ax \geq 0 \quad \text{with} \quad b - Ax \in K$$

define **slack vector**  $s = b - Ax \in K$   
for weak duality, dual  $y$  must satisfy

$$\langle y, s \rangle \geq 0 \quad \forall s \in K$$

## Dual cones

this inequality defines the **dual cone**  $K^*$ :

### Definition (dual cone)

the dual cone  $K^*$  of a cone  $K$  is the set of vectors  $y$  such that

$$\langle y, s \rangle \geq 0 \quad \forall s \in K$$

## Dual cones

this inequality defines the **dual cone**  $K^*$ :

### Definition (dual cone)

the dual cone  $K^*$  of a cone  $K$  is the set of vectors  $y$  such that

$$\langle y, s \rangle \geq 0 \quad \forall s \in K$$

examples of cones and their duals:

- ▶  $K$  acute,  $K^*$  obtuse
- ▶  $K = \mathbf{R}_+^m$ ,  $K^* = \mathbf{R}_+^m$
- ▶  $K = \{x \in \mathbf{R}^n \mid \|x\| \leq x_0\}$ ,  $K^* = \{y \in \mathbf{R}^n \mid \|y\| \leq y_0\}$
- ▶  $K = \{X \in \mathbf{S}^n \mid X \succeq 0\}$ ,  $K^* = \{Y \in \mathbf{S}^n \mid Y \succeq 0\}$

## Conic duality

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

## Conic duality

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

for  $y \in K^*$ , construct Lagrangian  $\mathcal{L}(x, y) = \langle c, x \rangle - \langle y, b - Ax \rangle$ ,  
ensure value is **better** (lower) when  $x$  and  $y$  are feasible

## Conic duality

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

for  $y \in K^*$ , construct Lagrangian  $\mathcal{L}(x, y) = \langle c, x \rangle - \langle y, b - Ax \rangle$ ,  
ensure value is **better** (lower) when  $x$  and  $y$  are feasible

$$\mathcal{L}(x, y) \quad := \quad \langle c, x \rangle - \langle y, b - Ax \rangle$$



## Conic duality

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll} \text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n \end{array} \quad (\mathcal{P})$$

for  $y \in K^*$ , construct Lagrangian  $\mathcal{L}(x, y) = \langle c, x \rangle - \langle y, b - Ax \rangle$ ,  
ensure value is **better** (lower) when  $x$  and  $y$  are feasible

$$\begin{aligned} \mathcal{L}(x, y) &:= \langle c, x \rangle - \langle y, b - Ax \rangle \\ p^* &\geq \inf_{x \text{ feas}} \langle c, x \rangle - \langle y, b - Ax \rangle \\ &\geq \inf_x \langle c, x \rangle - \langle y, b - Ax \rangle \\ &= \langle y, -b \rangle + \inf_x \langle c + A^*y, x \rangle \end{aligned}$$

## Conic duality

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll}\text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n\end{array} \quad (\mathcal{P})$$

for  $y \in K^*$ , construct Lagrangian  $\mathcal{L}(x, y) = \langle c, x \rangle - \langle y, b - Ax \rangle$ ,  
ensure value is **better** (lower) when  $x$  and  $y$  are feasible

$$\begin{aligned}\mathcal{L}(x, y) &:= \langle c, x \rangle - \langle y, b - Ax \rangle \\ p^* &\geq \inf_{x \text{ feas}} \langle c, x \rangle - \langle y, b - Ax \rangle \\ &\geq \inf_x \langle c, x \rangle - \langle y, b - Ax \rangle \\ &= \langle y, -b \rangle + \inf_x \langle c + A^*y, x \rangle\end{aligned}$$

which is  $-\infty$  unless  $c + A^*y = 0$ , so

## Conic duality

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll}\text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n\end{array} \quad (\mathcal{P})$$

for  $y \in K^*$ , construct Lagrangian  $\mathcal{L}(x, y) = \langle c, x \rangle - \langle y, b - Ax \rangle$ ,  
ensure value is **better** (lower) when  $x$  and  $y$  are feasible

$$\begin{aligned}\mathcal{L}(x, y) &:= \langle c, x \rangle - \langle y, b - Ax \rangle \\ p^* &\geq \inf_{x \text{ feas}} \langle c, x \rangle - \langle y, b - Ax \rangle \\ &\geq \inf_x \langle c, x \rangle - \langle y, b - Ax \rangle \\ &= \langle y, -b \rangle + \inf_x \langle c + A^*y, x \rangle\end{aligned}$$

which is  $-\infty$  unless  $c + A^*y = 0$ , so define the **dual problem**

$$\begin{array}{ll}\text{maximize} & \langle y, -b \rangle \\ \text{subject to} & c + A^*y = 0 \\ \text{variable} & y \in K^*\end{array} \quad (\mathcal{D})$$

## Conic duality

primal problem with solution  $x^* \in \mathbf{R}^n$ , optimal value  $p^*$ :

$$\begin{array}{ll}\text{minimize} & \langle c, x \rangle \\ \text{subject to} & b - Ax \in K \\ \text{variable} & x \in \mathbf{R}^n\end{array} \quad (\mathcal{P})$$

for  $y \in K^*$ , construct Lagrangian  $\mathcal{L}(x, y) = \langle c, x \rangle - \langle y, b - Ax \rangle$ ,  
ensure value is **better** (lower) when  $x$  and  $y$  are feasible

$$\begin{aligned}\mathcal{L}(x, y) &:= \langle c, x \rangle - \langle y, b - Ax \rangle \\ p^* &\geq \inf_{x \text{ feas}} \langle c, x \rangle - \langle y, b - Ax \rangle \\ &\geq \inf_x \langle c, x \rangle - \langle y, b - Ax \rangle \\ &= \langle y, -b \rangle + \inf_x \langle c + A^*y, x \rangle\end{aligned}$$

which is  $-\infty$  unless  $c + A^*y = 0$ , so define the **dual problem**

$$\begin{array}{ll}\text{maximize} & \langle y, -b \rangle \\ \text{subject to} & c + A^*y = 0 \\ \text{variable} & y \in K^*\end{array} \quad (\mathcal{D})$$

## Dual of the dual

- ▶ if  $(\mathcal{P})$  is convex, then the dual of  $(1)$  is  $(\mathcal{P})$
- ▶ otherwise, the dual of the dual is the **convexification** of the primal

picture

## Strong duality for LPs

primal and dual LP in standard form:

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

$$\begin{array}{ll}\text{maximize} & b^T y \\ \text{subject to} & A^T y \leq c\end{array}$$

**claim:** if primal LP has a bounded feasible solution  $x^*$ , then strong duality holds

*i.e.*, dual LP has a bounded feasible solution  $y^*$  and  $p^* = d^*$

## Proof of strong duality for LPs

consider the following system with variables  $x' \in \mathbf{R}^n$ ,  $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

## Proof of strong duality for LPs

consider the following system with variables  $x' \in \mathbf{R}^n$ ,  $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

**claim:** this system has no solution. pf by contradiction:

- ▶ if  $\tau > 0$ , then  $x'/\tau$  is feasible for LP and  $c^T x'/\tau < p^*$
- ▶ if  $\tau = 0$ , then  $x^* + x'$  is feasible for LP and  $c^T(x^* + x') < p^*$



## Proof of strong duality for LPs

consider the following system with variables  $x' \in \mathbf{R}^n$ ,  $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

**claim:** this system has no solution. pf by contradiction:

- ▶ if  $\tau > 0$ , then  $x'/\tau$  is feasible for LP and  $c^T x'/\tau < p^*$
- ▶ if  $\tau = 0$ , then  $x^* + x'$  is feasible for LP and  $c^T(x^* + x') < p^*$

**so use Farkas' lemma:**

$$Ax + b = 0, \quad x \geq 0 \quad \text{or} \quad A^T y \geq 0, \quad b^T y < 0$$

## Proof of strong duality for LPs

consider the following system with variables  $x' \in \mathbf{R}^n$ ,  $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

**claim:** this system has no solution. pf by contradiction:

- ▶ if  $\tau > 0$ , then  $x'/\tau$  is feasible for LP and  $c^T x'/\tau < p^*$
- ▶ if  $\tau = 0$ , then  $x^* + x'$  is feasible for LP and  $c^T(x^* + x') < p^*$

**so use Farkas' lemma:**

$$\begin{array}{ll} Ax + b = 0, \quad x \geq 0 & \text{or} \quad A^T y \geq 0, \quad b^T y < 0 \\ \begin{bmatrix} A & -b \\ c^T & -p^* \end{bmatrix} \begin{bmatrix} x \\ \tau \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \end{bmatrix} & \text{or} \quad \begin{bmatrix} A^T & c \\ -b^T & -p^* \end{bmatrix} \begin{bmatrix} y \\ \sigma \end{bmatrix} \geq 0, \quad \sigma > 0 \end{array}$$

## Proof of strong duality for LPs

consider the following system with variables  $x' \in \mathbf{R}^n$ ,  $\tau \in \mathbf{R}$

$$Ax' - b\tau = 0, \quad c^T x' = p^* \tau - 1, \quad (x', \tau) \geq 0$$

**claim:** this system has no solution. pf by contradiction:

- ▶ if  $\tau > 0$ , then  $x'/\tau$  is feasible for LP and  $c^T x'/\tau < p^*$
- ▶ if  $\tau = 0$ , then  $x^* + x'$  is feasible for LP and  $c^T(x^* + x') < p^*$

**so use Farkas' lemma:**

$$\begin{array}{ll} Ax + b = 0, \quad x \geq 0 & \text{or} \quad A^T y \geq 0, \quad b^T y < 0 \\ \begin{bmatrix} A & -b \\ c^T & -p^* \end{bmatrix} \begin{bmatrix} x \\ \tau \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \end{bmatrix} & \text{or} \quad \begin{bmatrix} A^T & c \\ -b^T & -p^* \end{bmatrix} \begin{bmatrix} y \\ \sigma \end{bmatrix} \geq 0, \quad \sigma > 0 \end{array}$$

use second system to show  $y/\sigma$  is dual feasible and optimal

## Strong duality and complementary slackness

### Definition (complementary slackness)

The primal-dual pair  $x$  and  $y$  are **complementary** if

$$\langle y, b - Ax \rangle = 0$$

They satisfy **strict complementary slackness** if  $y_i(b_i - a_i^T x) = 0$  for  $i = 1, \dots, n$ .

for conic problem, strong duality  $\iff$  complementary slackness

$$\begin{aligned}\langle y, s \rangle &= \langle y, b - Ax \rangle \\ &= \langle y, b \rangle - \langle A^* y, x \rangle \\ &= \langle y, b \rangle - \langle c, x \rangle\end{aligned}$$

## KKT conditions

KKT conditions give **necessary** conditions for optimality

### Theorem (KKT conditions)

*Suppose  $x^*$  and  $y^*$  are primal and dual optimal, respectively.  
Then*

- ▶  *$x^*$  and  $y^*$  are a saddle point of the Lagrangian*
- ▶  *$x^*$  is primal feasible*
- ▶  *$y^*$  is dual feasible*
- ▶  *$x^*$  and  $y^*$  are complementary*

KKT conditions turn optimization problem into a system of equations