

Optimization Under Uncertainty
(but really, just Robust Optimization)

Lecture 18

December 2, 2024

Quick Announcements

- Homework 5 due on Tuesday (Dec 3)
- Office Hours this week - extended schedule (Ed Announcement coming up)
- Final exam topics
- Any questions?

Outline for Today

1 Introduction

- Some Motivating Examples
- A History Detour
- Pros and Cons of Probabilistic Models

2 Robust Optimization


- Basic Premises
- Modeling with Basic Uncertainty Sets
- Reformulating and Solving Robust Models
- Extensions
- Some Applications
- Calibrating Uncertainty Sets
- Distributionally Robust Optimization
- Connections with Other Areas

3 Dynamic Robust Optimization

- Properly Writing a Robust DP
- An Inventory Example
- Tractable Approximations with Decision Rules
- Some Practical Issues
- Bellman Optimality
- An Application in Monitoring

The Flaw of Averages

Optimization based on *nominal* values can lead to *severe* issues...



Taken from "Flaw of averages" Sam Savage (2009, 2012)

How Robust Are Optimal Solutions? (Ben-Tal & Nemirovski)

How Robust Are Optimal Solutions? (Ben-Tal & Nemirovski)

- Consider a **real-world scheduling problem** problem (PILOT4) in NETLIB Library

- One of the constraints is the following linear constraint $\bar{\mathbf{a}}^T \mathbf{x} \geq b$:

$$\begin{aligned} & -15.79081 \cdot x_{826} - 8.598819 \cdot x_{827} - 1.88789 \cdot x_{828} - 1.362417 \cdot x_{829} \\ & -1.526049 \cdot x_{830} - 0.031883 \cdot x_{849} - 28.725555 \cdot x_{850} - 10.792065 \cdot x_{851} \\ & -0.19004 \cdot x_{852} - 2.757176 \cdot x_{853} - 12.290832 \cdot x_{854} + 717.562256 \cdot x_{855} \\ & -0.057865x \cdot x_{856} - 3.785417 \cdot x_{857} - 78.30661 \cdot x_{858} - 122.163055 \cdot x_{859} \\ & -6.46609 \cdot x_{860} - 0.48371 \cdot x_{861} - 0.615264 \cdot x_{862} - 1.353783 \cdot x_{863} \\ & -84.644257 \cdot x_{864} - 122.459045 \cdot x_{865} - 43.15593 \cdot x_{866} - 1.712592 \cdot x_{870} \\ & -0.401597 \cdot x_{871} + x_{880} - 0.946049 \cdot x_{898} - 0.946049 \cdot x_{916} \geq 23.387405 \end{aligned}$$

- Coefficients like 8.598819 are estimated and potentially inaccurate

How Robust Are Optimal Solutions? (Ben-Tal & Nemirovski)

- Consider a **real-world scheduling problem** problem (PILOT4) in NETLIB Library

- One of the constraints is the following linear constraint $\bar{\mathbf{a}}^T \mathbf{x} \geq b$:

$$\begin{aligned} & -15.79081 \cdot x_{826} - 8.598819 \cdot x_{827} - 1.88789 \cdot x_{828} - 1.362417 \cdot x_{829} \\ & -1.526049 \cdot x_{830} - 0.031883 \cdot x_{849} - 28.725555 \cdot x_{850} - 10.792065 \cdot x_{851} \\ & -0.19004 \cdot x_{852} - 2.757176 \cdot x_{853} - 12.290832 \cdot x_{854} + 717.562256 \cdot x_{855} \\ & -0.057865x \cdot x_{856} - 3.785417 \cdot x_{857} - 78.30661 \cdot x_{858} - 122.163055 \cdot x_{859} \\ & -6.46609 \cdot x_{860} - 0.48371 \cdot x_{861} - 0.615264 \cdot x_{862} - 1.353783 \cdot x_{863} \\ & -84.644257 \cdot x_{864} - 122.459045 \cdot x_{865} - 43.15593 \cdot x_{866} - 1.712592 \cdot x_{870} \\ & -0.401597 \cdot x_{871} + x_{880} - 0.946049 \cdot x_{898} - 0.946049 \cdot x_{916} \geq 23.387405 \end{aligned}$$

- Coefficients like 8.598819 are estimated and potentially inaccurate
- What if these coefficients are just 0.1% inaccurate?
 - i.e., suppose the true \mathbf{a} is not $\bar{\mathbf{a}}$, but $|\alpha_i - \bar{\alpha}_i| \leq 0.001|\bar{\alpha}_i|$?
- Will the optimal solution to the problem still be feasible?
- How can we test?

How Robust Are Optimal Solutions? (Ben-Tal & Nemirovski)

- Original constraint: $\bar{\mathbf{a}}^T \mathbf{x} \geq b$, optimal solution \mathbf{x}^*
- Suppose true \mathbf{a} satisfies $|a_i - \bar{a}_i| \leq 0.001|\bar{a}_i|, \forall i$
- How to determine if the constraint is violated?

How Robust Are Optimal Solutions? (Ben-Tal & Nemirovski)

- Original constraint: $\bar{\mathbf{a}}^T \mathbf{x} \geq b$, optimal solution \mathbf{x}^*
- Suppose true \mathbf{a} satisfies $|a_i - \bar{a}_i| \leq 0.001|\bar{a}_i|, \forall i$

- How to determine if the constraint is violated?

$$\begin{aligned} \min_{\mathbf{a}} \quad & \mathbf{a}^T \mathbf{x}^* - b \\ \text{s.t.} \quad & |a_i - \bar{a}_i| \leq 0.001|\bar{a}_i|, \forall i \end{aligned}$$

- ▶ For PILOT4, this comes to $-128.8 \approx -4.5b$, so 450% violation!

How Robust Are Optimal Solutions? (Ben-Tal & Nemirovski)

- Original constraint: $\bar{\mathbf{a}}^T \mathbf{x} \geq b$, optimal solution \mathbf{x}^*
- Suppose true \mathbf{a} satisfies $|a_i - \bar{a}_i| \leq 0.001|\bar{a}_i|, \forall i$

- How to determine if the constraint is violated?

$$\begin{aligned} \min_{\mathbf{a}} \quad & \mathbf{a}^T \mathbf{x}^* - b \\ \text{s.t.} \quad & |a_i - \bar{a}_i| \leq 0.001|\bar{a}_i|, \forall i \end{aligned}$$

- For PILOT4, this comes to $-128.8 \approx -4.5b$, so 450% violation!

- OK, but perhaps we're too conservative?

- Suppose $a_i = \bar{a}_i + \varepsilon_i|\bar{a}_i|$, where $\varepsilon_i \sim \text{Uniform}[-0.001, 0.001]$
 - Using Monte-Carlo simulation with 1,000 samples:

$$\star \quad \mathbb{P}(\text{infeasible}) = 50\%, \mathbb{P}(\text{violation} > 150\%) = 18\%, \mathbb{E}[\text{violation}] = 125\%$$

How Robust Are Optimal Solutions? (Ben-Tal & Nemirovski)

- Original constraint: $\bar{\mathbf{a}}^T \mathbf{x} \geq b$, optimal solution \mathbf{x}^*
- Suppose true \mathbf{a} satisfies $|a_i - \bar{a}_i| \leq 0.001|\bar{a}_i|, \forall i$

- How to determine if the constraint is violated?

$$\begin{aligned} \min_{\mathbf{a}} \quad & \mathbf{a}^T \mathbf{x}^* - b \\ \text{s.t.} \quad & |a_i - \bar{a}_i| \leq 0.001|\bar{a}_i|, \forall i \end{aligned}$$

- For PILOT4, this comes to $-128.8 \approx -4.5b$, so 450% violation!

- OK, but perhaps we're too conservative?

- Suppose $a_i = \bar{a}_i + \varepsilon_i|\bar{a}_i|$, where $\varepsilon_i \sim \text{Uniform}[-0.001, 0.001]$
 - Using Monte-Carlo simulation with 1,000 samples:

$$\star \quad \mathbb{P}(\text{infeasible}) = 50\%, \mathbb{P}(\text{violation} > 150\%) = 18\%, \mathbb{E}[\text{violation}] = 125\%$$

- Disturbing that nominal solutions are likely highly infeasible
- Turns out to be the case for many **NETLIB** problems
- We should **capture uncertainty more explicitly** apriori!

Decisions Under Uncertainty

- Decision Maker (DM) must choose x , without knowing z
- DM incurs a **cost** $C(x, z)$

Decisions Under Uncertainty

- Decision Maker (DM) must choose x , without knowing z
- DM incurs a **cost** $C(x, z)$
- How to model z ? How to properly formalize the decision problem?

Decisions Under Uncertainty

- Decision Maker (DM) must choose x , without knowing z
- DM incurs a **cost** $C(x, z)$
- How to model z ? How to properly formalize the decision problem?
- “Standard” probabilistic model:
 - There is a unique probability distribution \mathbb{P} for z
 - DM considers an objective: $\min_x \mathbb{E}_{z \sim \mathbb{P}}[C(x, z)]$

Classical Probabilistic Model: DM knows \mathbb{P} , solves $\min_{\mathbf{x}} \mathbb{E}_{\mathbf{z} \sim \mathbb{P}} [C(\mathbf{x}, \mathbf{z})]$

Classical Probabilistic Model: DM knows \mathbb{P} , solves $\min_{\mathbf{x}} \mathbb{E}_{\mathbf{z} \sim \mathbb{P}} [C(\mathbf{x}, \mathbf{z})]$

- What if there are constraints?

$$f_i(\mathbf{x}, \mathbf{z}) \geq 0, \forall i \in I$$

- Where is \mathbb{P} coming from?
- When is this reasonable?
- What if \mathbb{P} is **not** the actual distribution?
- What if \mathbb{P} is not exogenous?

Classical Probabilistic Model: DM knows \mathbb{P} , solves $\min_{\mathbf{x}} \mathbb{E}_{\mathbf{z} \sim \mathbb{P}}[C(\mathbf{x}, \mathbf{z})]$

- What if there are constraints?

$$f_i(\mathbf{x}, \mathbf{z}) \geq 0, \forall i \in I$$

- Where is \mathbb{P} coming from?
- When is this reasonable?
- What if \mathbb{P} is **not** the actual distribution?
- What if \mathbb{P} is not exogenous?

- expectation constraint:

$$\mathbb{E}_{\mathbb{P}}[f_i(\mathbf{x}, \mathbf{z})] \geq 0, \forall i$$

- chance constraint:

$$\text{individual: } \mathbb{P}[f_i(\mathbf{x}, \mathbf{z}) \geq 0] \geq 1 - \epsilon, \forall i$$

$$\text{joint: } \mathbb{P}[f_i(\mathbf{x}, \mathbf{z}) \geq 0, \forall i] \geq 1 - \epsilon$$

- robust (a.s.) constraint: $F(\mathbf{x}, \mathbf{z}) \geq 0, \forall \mathbf{z}$
- easy to check / enforce?

Classical Probabilistic Model: DM knows \mathbb{P} , solves $\min_{\mathbf{x}} \mathbb{E}_{\mathbf{z} \sim \mathbb{P}} [C(\mathbf{x}, \mathbf{z})]$

- What if there are constraints?

$$f_i(\mathbf{x}, \mathbf{z}) \geq 0, \forall i \in I$$

- Where is \mathbb{P} coming from?
- When is this reasonable?
- What if \mathbb{P} is **not** the actual distribution?
- What if \mathbb{P} is not exogenous?

- expectation constraint:

$$\mathbb{E}_{\mathbb{P}} [f_i(\mathbf{x}, \mathbf{z})] \geq 0, \forall i$$

- chance constraint:

$$\text{individual: } \mathbb{P}[f_i(\mathbf{x}, \mathbf{z}) \geq 0] \geq 1 - \epsilon, \forall i$$

$$\text{joint: } \mathbb{P}[f_i(\mathbf{x}, \mathbf{z}) \geq 0, \forall i] \geq 1 - \epsilon$$

- robust (a.s.) constraint: $F(\mathbf{x}, \mathbf{z}) \geq 0, \forall \mathbf{z}$
- easy to check / enforce?

- Theory unable to analyze **complex, real-world** dynamics
 - poor data, changing environments (future \neq past), many agents, ...
- Framework not geared towards **computing decisions**
 - Limited computational tractability, particularly in higher dimensions
- With $C = -u(\cdot)$ (u utility function), unclear if this is a good behavioral model

An Alternative Model of Uncertainty

An Alternative Model of Uncertainty

- Let's admit **explicitly** that our model of reality is **incorrect**
- From **classical view**: “we know distribution \mathbb{P} for \mathbf{z} , and solve: $\min_x \mathbb{E}_{\mathbb{P}}[C(\mathbf{x}, \mathbf{z})]$ ”
to **robust view**: “we only know that $\mathbb{P} \in \mathcal{P}$, and solve: $\min_x \max_{\mathbb{P} \in \mathcal{P}} \mathbb{E}_{\mathbb{P}}[C(\mathbf{x}, \mathbf{z})]$ ”

An Alternative Model of Uncertainty

- Let's admit **explicitly** that our model of reality is **incorrect**
- From **classical view**: “we know distribution \mathbb{P} for \mathbf{z} , and solve: $\min_x \mathbb{E}_{\mathbb{P}}[C(\mathbf{x}, \mathbf{z})]$ ”
to **robust view**: “we only know that $\mathbb{P} \in \mathcal{P}$, and solve: $\min_x \max_{\mathbb{P} \in \mathcal{P}} \mathbb{E}_{\mathbb{P}}[C(\mathbf{x}, \mathbf{z})]$ ”

Long history of **robust decision-making** and **model misspecification**:

- **Economics**:
 - ▶ Frank Knight (1921) - risk vs. Knightian uncertainty, Abraham Wald (1939), John von Neumann (1944) zero-sum games
 - ▶ Savage (1951): minimax regret, Scarf (1958): robust Newsvendor model
 - ▶ Schmeidler, Gilboa (1980s): axiomatic frameworks, Ben-Haim (1980s): info-gap theory
 - ▶ Hansen & Sargent (2008): “*Robustness*” - robust control in macroeconomics
 - ▶ Bergemann & Morris (2012): “*Robust mechanism design*” book, Carroll (2015), ...
- **Engineering and robust control**: Bertsekas (1970s), Doyle (1980s), etc.
- **Computer science**: complexity analysis; adversarial training (modern!)
- **Statistics**: M-estimators Huber (1981)
- **Operations Research**:
 - ▶ Early work by Soyster (1973), Libura (1980), Bard (1984), Kouvelis (1997)
 - ▶ **Robust Optimization**: Ben-Tal, Nemirovski, El-Ghaoui ('90s), Bertsimas, Sim ('00s)
 - ▶ Two books: Ben-Tal, El-Ghaoui, Nemirovski (2009), Bertsimas, den Hertog (2020)
 - ▶ Many tutorials!

An Alternative Model of Uncertainty

- Let's admit **explicitly** that our model of reality is **incorrect**
- From **classical view**: “we know distribution \mathbb{P} for \mathbf{z} , and solve: $\min_{\mathbf{x}} \mathbb{E}_{\mathbb{P}}[C(\mathbf{x}, \mathbf{z})]$ ”
to **robust view**: “we only know that $\mathbb{P} \in \mathcal{P}$, and solve: $\min_{\mathbf{x}} \max_{\mathbb{P} \in \mathcal{P}} \mathbb{E}_{\mathbb{P}}[C(\mathbf{x}, \mathbf{z})]$ ”

Why robust optimization? (in my view)

1. Very sensible
2. Modest modeling requirements
3. Modest in its premise: “*always under-promises, and over-delivers*”
4. Tractable: quickly becoming “technology”
5. Very sensible results: can rationalize simple rules in complex problems

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- ① Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
 - Other options possible, based on notions of **regret**
- Conservative?

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- ① Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
 - Other options possible, based on notions of **regret**
- Conservative?
 - **Not necessarily!**
 - \mathcal{U} directly trades off robustness and conservatism, and is ultimately a **modeling choice**

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- ① Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
 - Other options possible, based on notions of **regret**
- Conservative?
 - **Not necessarily!**
 - \mathcal{U} directly trades off robustness and conservatism, and is ultimately a **modeling choice**
- Is there a probabilistic interpretation?

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- ① Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
 - Other options possible, based on notions of **regret**
- Conservative?
 - **Not necessarily!**
 - \mathcal{U} directly trades off robustness and conservatism, and is ultimately a **modeling choice**
- Is there a probabilistic interpretation?
 - Objective = $\sup_{\mathbb{P} \in \mathcal{P}} \mathbb{E}_{\mathbf{z} \sim \mathbb{P}}[C(\mathbf{x}, \mathbf{z})]$ where \mathcal{P} is the set of all measures with support \mathcal{U}
 - So we are assuming that the only information about \mathbb{P} is the support \mathcal{U}

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
- Each constraint is “hard”: must be satisfied *robustly*, for any realization of \mathbf{z}

What is the optimal value of the following robust LP?

$$\begin{array}{ll} \min_{\mathbf{x}} \max_{\mathbf{a} \in \mathcal{U}} & -(x_1 + x_2) \\ \text{such that} & x_1 \leq a_1 \\ & x_2 \leq a_2 \\ & x_1 + x_2 \leq 1. \end{array} \quad \text{where } \mathcal{U} = \{(a_1, a_2) \in [0, 1]^2 : a_1 + a_2 \leq 1\}$$

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- 1 Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
- 2 Each constraint is “hard”: must be satisfied *robustly*, for any realization of \mathbf{z}

What is the optimal value of the following robust LP?

$$\begin{array}{ll} \min_{\mathbf{x}} \max_{\mathbf{a} \in \mathcal{U}} & -(x_1 + x_2) \\ \text{such that} & x_1 \leq a_1 \\ & x_2 \leq a_2 \\ & x_1 + x_2 \leq 1. \end{array} \quad \text{where } \mathcal{U} = \{(a_1, a_2) \in [0, 1]^2 : a_1 + a_2 \leq 1\}$$

Optimal value 0. In RO, **each constraint must be satisfied separately, robustly.**

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- 1 Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
- 2 Each constraint is “hard”: must be satisfied *robustly*, for any realization of \mathbf{z}
- 3 Each constraint can be re-written as an optimization problem

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- 1 Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
- 2 Each constraint is “hard”: must be satisfied *robustly*, for any realization of \mathbf{z}
- 3 Each constraint can be re-written as an optimization problem

$$\boxed{f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}} \quad \Leftrightarrow \quad \boxed{\sup_{\mathbf{z} \in \mathcal{U}} f_i(\mathbf{x}, \mathbf{z}) \leq 0}$$

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- 1 Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
- 2 Each constraint is “hard”: must be satisfied *robustly*, for any realization of \mathbf{z}
- 3 Each constraint can be re-written as an optimization problem
- 4 Without loss, we can consider a problem where \mathbf{z} only appears in constraints

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- 1 Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
- 2 Each constraint is “hard”: must be satisfied *robustly*, for any realization of \mathbf{z}
- 3 Each constraint can be re-written as an optimization problem
- 4 Without loss, we can consider a problem where \mathbf{z} only appears in constraints
(P) is equivalent to the following problem:

$$\begin{array}{ll} \inf_{\mathbf{x}, t} & t \\ \text{s.t. } & t \geq C(\mathbf{x}, \mathbf{z}), \forall \mathbf{z} \in \mathcal{U} \\ & f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- 1 Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
- 2 Each constraint is “hard”: must be satisfied *robustly*, for any realization of \mathbf{z}
- 3 Each constraint can be re-written as an optimization problem
- 4 Without loss, we can consider a problem where \mathbf{z} only appears in constraints
(P) is equivalent to the following problem:

$$\begin{array}{ll} \inf_{\mathbf{x}, t} & t \\ \text{s.t. } & t \geq C(\mathbf{x}, \mathbf{z}), \forall \mathbf{z} \in \mathcal{U} \\ & f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Many RO models are in this *epigraph reformulation*, and focus on constraints

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- 1 Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
- 2 Each constraint is “hard”: must be satisfied *robustly*, for any realization of \mathbf{z}
- 3 Each constraint can be re-written as an optimization problem
- 4 Without loss, we can consider a problem where \mathbf{z} only appears in constraints
- 5 DM only responsible for objective and constraints when $\mathbf{z} \in \mathcal{U}$
 - If $\mathbf{z} \notin \mathcal{U}$ actually occurs, all bets are off
 - Can extend framework to ensure **gradual** degradation of performance:
Globalized robust counterparts (Ben-Tal & Nemirovski)

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

Remarks.

- 1 Objective: worst-case performance $\sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z})$
- 2 Each constraint is “hard”: must be satisfied *robustly*, for any realization of \mathbf{z}
- 3 Each constraint can be re-written as an optimization problem
- 4 Without loss, we can consider a problem where \mathbf{z} only appears in constraints
- 5 DM only responsible for objective and constraints when $\mathbf{z} \in \mathcal{U}$
- 6 Robust model seems to lead to a **difficult** optimization problem
 - For any given \mathbf{x} , checking constraints/solving the “adversary” problem may be tough
 - We must also solve our original problem of finding \mathbf{x} !

“Classical” Robust Optimization (RO)

- Robust Optimization: the values of \mathbf{z} belong to an **uncertainty set** \mathcal{U}
- DM reformulates the original optimization problem as:

$$\begin{array}{ll} \text{(P)} & \inf_{\mathbf{x}} \sup_{\mathbf{z} \in \mathcal{U}} C(\mathbf{x}, \mathbf{z}) \\ & \text{s.t. } f_i(\mathbf{x}, \mathbf{z}) \leq 0, \forall \mathbf{z} \in \mathcal{U}, \forall i \in I \end{array}$$

1. How to model \mathcal{U}
2. How to formulate and solve the **robust counterpart**
3. Why is this useful, in theory and in practice

Intuition for Some Basic Uncertainty Sets

- Recall PILOT4; how to build some “safety buffers” for constraint like #372:

$$\begin{aligned} & -15.79081 \cdot x_{826} - 8.598819 \cdot x_{827} - 1.88789 \cdot x_{828} - 1.362417 \cdot x_{829} \\ & -1.526049 \cdot x_{830} - 0.031883 \cdot x_{849} - 28.725555 \cdot x_{850} - 10.792065 \cdot x_{851} \\ & -0.19004 \cdot x_{852} - 2.757176 \cdot x_{853} - 12.290832 \cdot x_{854} + 717.562256 \cdot x_{855} \\ & -0.057865x \cdot x_{856} - 3.785417 \cdot x_{857} - 78.30661 \cdot x_{858} - 122.163055 \cdot x_{859} \\ & -6.46609 \cdot x_{860} - 0.48371 \cdot x_{861} - 0.615264 \cdot x_{862} - 1.353783 \cdot x_{863} \\ & -84.644257 \cdot x_{864} - 122.459045 \cdot x_{865} - 43.15593 \cdot x_{866} - 1.712592 \cdot x_{870} \\ & -0.401597 \cdot x_{871} + x_{880} - 0.946049 \cdot x_{898} - 0.946049 \cdot x_{916} \geq 23.387405 \end{aligned}$$

- Consider a **linear constraint** in x with coefficients that depend **linearly** on z

$$(\bar{a} + Pz)^T x \leq b, \forall z \in \mathcal{U}$$

Intuition for Some Basic Uncertainty Sets

- Recall PILOT4; how to build some “safety buffers” for constraint like #372:

$$\begin{aligned} & -15.79081 \cdot x_{826} - 8.598819 \cdot x_{827} - 1.88789 \cdot x_{828} - 1.362417 \cdot x_{829} \\ & -1.526049 \cdot x_{830} - 0.031883 \cdot x_{849} - 28.725555 \cdot x_{850} - 10.792065 \cdot x_{851} \\ & -0.19004 \cdot x_{852} - 2.757176 \cdot x_{853} - 12.290832 \cdot x_{854} + 717.562256 \cdot x_{855} \\ & -0.057865x \cdot x_{856} - 3.785417 \cdot x_{857} - 78.30661 \cdot x_{858} - 122.163055 \cdot x_{859} \\ & -6.46609 \cdot x_{860} - 0.48371 \cdot x_{861} - 0.615264 \cdot x_{862} - 1.353783 \cdot x_{863} \\ & -84.644257 \cdot x_{864} - 122.459045 \cdot x_{865} - 43.15593 \cdot x_{866} - 1.712592 \cdot x_{870} \\ & -0.401597 \cdot x_{871} + x_{880} - 0.946049 \cdot x_{898} - 0.946049 \cdot x_{916} \geq 23.387405 \end{aligned}$$

- Consider a **linear constraint** in x with coefficients that depend **linearly** on z

$$(\bar{a} + Pz)^T x \leq b, \forall z \in \mathcal{U}$$

- P is a known matrix; z is primitive uncertainty

- Q:** Why this more general form?

A: For modeling flexibility:

- Suppose the same physical quantity (i.e., coefficient) appears in multiple constraints
- Can capture “correlations”, e.g., with a factor model

Intuition for Some Basic Uncertainty Sets

- Recall PILOT4; how to build some “safety buffers” for constraint like #372:

$$\begin{aligned} & -15.79081 \cdot x_{826} - 8.598819 \cdot x_{827} - 1.88789 \cdot x_{828} - 1.362417 \cdot x_{829} \\ & -1.526049 \cdot x_{830} - 0.031883 \cdot x_{849} - 28.725555 \cdot x_{850} - 10.792065 \cdot x_{851} \\ & -0.19004 \cdot x_{852} - 2.757176 \cdot x_{853} - 12.290832 \cdot x_{854} + 717.562256 \cdot x_{855} \\ & -0.057865x \cdot x_{856} - 3.785417 \cdot x_{857} - 78.30661 \cdot x_{858} - 122.163055 \cdot x_{859} \\ & -6.46609 \cdot x_{860} - 0.48371 \cdot x_{861} - 0.615264 \cdot x_{862} - 1.353783 \cdot x_{863} \\ & -84.644257 \cdot x_{864} - 122.459045 \cdot x_{865} - 43.15593 \cdot x_{866} - 1.712592 \cdot x_{870} \\ & -0.401597 \cdot x_{871} + x_{880} - 0.946049 \cdot x_{898} - 0.946049 \cdot x_{916} \geq 23.387405 \end{aligned}$$

- Consider a **linear constraint** in x with coefficients that depend **linearly** on z

$$(\bar{a} + Pz)^T x \leq b, \forall z \in \mathcal{U}$$

- How about a **box** uncertainty set? For some confidence level ρ :

$$\mathcal{U}_{\text{box}} := \{z : -\rho \leq z_i \leq \rho\} = \{z : \|z\|_{\infty} \leq \rho\}$$

“Too conservative?”

- In PILOT4, **robust** solution is within 1% of x^* for objective
- Recall that x^* would violate this constraint by 450%
- Sometimes not much is sacrificed for robustness!

Intuition for Some Basic Uncertainty Sets

- Recall PILOT4; how to build some “safety buffers” for constraint like #372:

$$\begin{aligned} & -15.79081 \cdot x_{826} - 8.598819 \cdot x_{827} - 1.88789 \cdot x_{828} - 1.362417 \cdot x_{829} \\ & -1.526049 \cdot x_{830} - 0.031883 \cdot x_{849} - 28.725555 \cdot x_{850} - 10.792065 \cdot x_{851} \\ & -0.19004 \cdot x_{852} - 2.757176 \cdot x_{853} - 12.290832 \cdot x_{854} + 717.562256 \cdot x_{855} \\ & -0.057865x \cdot x_{856} - 3.785417 \cdot x_{857} - 78.30661 \cdot x_{858} - 122.163055 \cdot x_{859} \\ & -6.46609 \cdot x_{860} - 0.48371 \cdot x_{861} - 0.615264 \cdot x_{862} - 1.353783 \cdot x_{863} \\ & -84.644257 \cdot x_{864} - 122.459045 \cdot x_{865} - 43.15593 \cdot x_{866} - 1.712592 \cdot x_{870} \\ & -0.401597 \cdot x_{871} + x_{880} - 0.946049 \cdot x_{898} - 0.946049 \cdot x_{916} \geq 23.387405 \end{aligned}$$

- Consider a **linear constraint** in x with coefficients that depend **linearly** on z

$$(\bar{a} + Pz)^T x \leq b, \forall z \in \mathcal{U}$$

- How about a **box** uncertainty set? For some confidence level ρ :

$$\mathcal{U}_{\text{box}} := \{z : -\rho \leq z_i \leq \rho\} = \{z : \|z\|_{\infty} \leq \rho\}$$

- Or maybe an **ellipsoid** would be less conservative:

$$\mathcal{U}_{\text{ellipsoid}} := \{z : \|z\|_2 \leq \rho\}$$

Intuition for Some Basic Uncertainty Sets

- Recall PILOT4; how to build some “safety buffers” for constraint like #372:

$$\begin{aligned} & -15.79081 \cdot x_{826} - 8.598819 \cdot x_{827} - 1.88789 \cdot x_{828} - 1.362417 \cdot x_{829} \\ & -1.526049 \cdot x_{830} - 0.031883 \cdot x_{849} - 28.725555 \cdot x_{850} - 10.792065 \cdot x_{851} \\ & -0.19004 \cdot x_{852} - 2.757176 \cdot x_{853} - 12.290832 \cdot x_{854} + 717.562256 \cdot x_{855} \\ & -0.057865x \cdot x_{856} - 3.785417 \cdot x_{857} - 78.30661 \cdot x_{858} - 122.163055 \cdot x_{859} \\ & -6.46609 \cdot x_{860} - 0.48371 \cdot x_{861} - 0.615264 \cdot x_{862} - 1.353783 \cdot x_{863} \\ & -84.644257 \cdot x_{864} - 122.459045 \cdot x_{865} - 43.15593 \cdot x_{866} - 1.712592 \cdot x_{870} \\ & -0.401597 \cdot x_{871} + x_{880} - 0.946049 \cdot x_{898} - 0.946049 \cdot x_{916} \geq 23.387405 \end{aligned}$$

- Consider a **linear constraint** in x with coefficients that depend **linearly** on z

$$(\bar{a} + Pz)^T x \leq b, \forall z \in \mathcal{U}$$

- How about a **box** uncertainty set? For some confidence level ρ :

$$\mathcal{U}_{\text{box}} := \{z : -\rho \leq z_i \leq \rho\} = \{z : \|z\|_{\infty} \leq \rho\}$$

- Or maybe an **ellipsoid** would be less conservative:

$$\mathcal{U}_{\text{ellipsoid}} := \{z : \|z\|_2 \leq \rho\}$$

- Or what if we gave “nature” a **budget** on how many coefficients it could change:

$$\mathcal{U}_{\text{budget}} := \{z : \|z\|_{\infty} \leq \rho, \|z\|_1 \leq \Gamma\rho\}$$

Intuition for Some Basic Uncertainty Sets

- Recall PILOT4; how to build some “safety buffers” for constraint like #372:

$$\begin{aligned} & -15.79081 \cdot x_{826} - 8.598819 \cdot x_{827} - 1.88789 \cdot x_{828} - 1.362417 \cdot x_{829} \\ & -1.526049 \cdot x_{830} - 0.031883 \cdot x_{849} - 28.725555 \cdot x_{850} - 10.792065 \cdot x_{851} \\ & -0.19004 \cdot x_{852} - 2.757176 \cdot x_{853} - 12.290832 \cdot x_{854} + 717.562256 \cdot x_{855} \\ & -0.057865x \cdot x_{856} - 3.785417 \cdot x_{857} - 78.30661 \cdot x_{858} - 122.163055 \cdot x_{859} \\ & -6.46609 \cdot x_{860} - 0.48371 \cdot x_{861} - 0.615264 \cdot x_{862} - 1.353783 \cdot x_{863} \\ & -84.644257 \cdot x_{864} - 122.459045 \cdot x_{865} - 43.15593 \cdot x_{866} - 1.712592 \cdot x_{870} \\ & -0.401597 \cdot x_{871} + x_{880} - 0.946049 \cdot x_{898} - 0.946049 \cdot x_{916} \geq 23.387405 \end{aligned}$$

- Consider a **linear constraint** in x with coefficients that depend **linearly** on z

$$(\bar{a} + Pz)^T x \leq b, \forall z \in \mathcal{U}$$

- How about a **box** uncertainty set? For some confidence level ρ :

$$\mathcal{U}_{\text{box}} := \{z : -\rho \leq z_i \leq \rho\} = \{z : \|z\|_{\infty} \leq \rho\}$$

- Or maybe an **ellipsoid** would be less conservative:

$$\mathcal{U}_{\text{ellipsoid}} := \{z : \|z\|_2 \leq \rho\}$$

- Or what if we gave “nature” a **budget** on how many coefficients it could change:

$$\mathcal{U}_{\text{budget}} := \{z : \|z\|_{\infty} \leq \rho, \|z\|_1 \leq \Gamma\rho\}$$

- How to formulate the robust counterpart? How to set ρ, Γ ? How to use in practice?

Formulating the Robust Counterpart (RC) for Box Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \quad \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{box}} = \{\mathbf{z} : \|\mathbf{z}\|_{\infty} \leq \rho\}$, satisfying the constraint robustly is equivalent to:

$$\max_{\mathbf{z}: \|\mathbf{z}\|_{\infty} \leq \rho} (\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b,$$

Formulating the Robust Counterpart (RC) for Box Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \quad \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{box}} = \{\mathbf{z} : \|\mathbf{z}\|_{\infty} \leq \rho\}$, satisfying the constraint robustly is equivalent to:

$$\max_{\mathbf{z} : \|\mathbf{z}\|_{\infty} \leq \rho} (\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b,$$

or

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z} : \|\mathbf{z}\|_{\infty} \leq \rho} (\mathbf{P}^T \mathbf{x})^T \mathbf{z} \leq b,$$

Formulating the Robust Counterpart (RC) for Box Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \quad \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{box}} = \{\mathbf{z} : \|\mathbf{z}\|_{\infty} \leq \rho\}$, satisfying the constraint robustly is equivalent to:

$$\max_{\mathbf{z}: \|\mathbf{z}\|_{\infty} \leq \rho} (\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b,$$

or

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z}: \|\mathbf{z}\|_{\infty} \leq \rho} (\mathbf{P}^T \mathbf{x})^T \mathbf{z} \leq b,$$

or

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z}: |\mathbf{z}_i| \leq \rho} \sum_i (\mathbf{P}^T \mathbf{x})_i \mathbf{z}_i \leq b,$$

Formulating the Robust Counterpart (RC) for Box Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \quad \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{box}} = \{\mathbf{z} : \|\mathbf{z}\|_{\infty} \leq \rho\}$, satisfying the constraint robustly is equivalent to:

$$\max_{\mathbf{z}: \|\mathbf{z}\|_{\infty} \leq \rho} (\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b,$$

or

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z}: \|\mathbf{z}\|_{\infty} \leq \rho} (\mathbf{P}^T \mathbf{x})^T \mathbf{z} \leq b,$$

or

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z}: |\mathbf{z}_i| \leq \rho} \sum_i (\mathbf{P}^T \mathbf{x})_i \mathbf{z}_i \leq b,$$

or

$$\bar{\mathbf{a}}^T \mathbf{x} + \rho \sum_i |(\mathbf{P}^T \mathbf{x})_i| \leq b,$$

Formulating the Robust Counterpart (RC) for Box Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \quad \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{box}} = \{\mathbf{z} : \|\mathbf{z}\|_{\infty} \leq \rho\}$, satisfying the constraint robustly is equivalent to:

$$\max_{\mathbf{z}: \|\mathbf{z}\|_{\infty} \leq \rho} (\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b,$$

or

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z}: \|\mathbf{z}\|_{\infty} \leq \rho} (\mathbf{P}^T \mathbf{x})^T \mathbf{z} \leq b,$$

or

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z}: |\mathbf{z}_i| \leq \rho} \sum_i (\mathbf{P}^T \mathbf{x})_i \mathbf{z}_i \leq b,$$

or

$$\bar{\mathbf{a}}^T \mathbf{x} + \rho \sum_i |(\mathbf{P}^T \mathbf{x})_i| \leq b,$$

or

$$\bar{\mathbf{a}}^T \mathbf{x} + \rho \|\mathbf{P}^T \mathbf{x}\|_1 \leq b.$$

Formulating the Robust Counterpart (RC) for Polyhedral Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^\top \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{polyhedral}} = \{\mathbf{z} : \mathbf{D}\mathbf{z} \leq \mathbf{d}\}$, satisfying the constraint robustly is equivalent to:

Formulating the Robust Counterpart (RC) for Polyhedral Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{polyhedral}} = \{\mathbf{z} : \mathbf{D}\mathbf{z} \leq \mathbf{d}\}$, satisfying the constraint robustly is equivalent to:

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z} : \mathbf{D}\mathbf{z} \leq \mathbf{d}} (\mathbf{P}^T \mathbf{x})^T \mathbf{z} \leq b. \quad (1)$$

Formulating the Robust Counterpart (RC) for Polyhedral Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{polyhedral}} = \{\mathbf{z} : \mathbf{D}\mathbf{z} \leq \mathbf{d}\}$, satisfying the constraint robustly is equivalent to:

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z} : \mathbf{D}\mathbf{z} \leq \mathbf{d}} (\mathbf{P}^T \mathbf{x})^T \mathbf{z} \leq b. \quad (1)$$

By strong LP duality, when the left-hand-side in (1) is finite, we must have:

$$\max\{(\mathbf{P}^T \mathbf{x})^T \mathbf{z} : \mathbf{D}\mathbf{z} \leq \mathbf{d}\} = \min\{\mathbf{d}^T \mathbf{y} : \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x}, \mathbf{y} \geq 0\}.$$

Formulating the Robust Counterpart (RC) for Polyhedral Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{polyhedral}} = \{\mathbf{z} : \mathbf{D}\mathbf{z} \leq \mathbf{d}\}$, satisfying the constraint robustly is equivalent to:

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z} : \mathbf{D}\mathbf{z} \leq \mathbf{d}} (\mathbf{P}^T \mathbf{x})^T \mathbf{z} \leq b. \quad (1)$$

By strong LP duality, when the left-hand-side in (1) is finite, we must have:

$$\max\{(\mathbf{P}^T \mathbf{x})^T \mathbf{z} : \mathbf{D}\mathbf{z} \leq \mathbf{d}\} = \min\{\mathbf{d}^T \mathbf{y} : \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x}, \mathbf{y} \geq 0\}.$$

Hence (1) is equivalent to

$$\bar{\mathbf{a}}^T \mathbf{x} + \min_{\mathbf{y}} \{\mathbf{d}^T \mathbf{y} : \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x}, \mathbf{y} \geq 0\} \leq b,$$

Formulating the Robust Counterpart (RC) for Polyhedral Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \quad \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{polyhedral}} = \{\mathbf{z} : \mathbf{D}\mathbf{z} \leq \mathbf{d}\}$, satisfying the constraint robustly is equivalent to:

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z} : \mathbf{D}\mathbf{z} \leq \mathbf{d}} (\mathbf{P}^T \mathbf{x})^T \mathbf{z} \leq b. \quad (1)$$

By strong LP duality, when the left-hand-side in (1) is finite, we must have:

$$\max\{(\mathbf{P}^T \mathbf{x})^T \mathbf{z} : \mathbf{D}\mathbf{z} \leq \mathbf{d}\} = \min\{\mathbf{d}^T \mathbf{y} : \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x}, \mathbf{y} \geq 0\}.$$

Hence (1) is equivalent to

$$\bar{\mathbf{a}}^T \mathbf{x} + \min_{\mathbf{y}} \{\mathbf{d}^T \mathbf{y} : \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x}, \mathbf{y} \geq 0\} \leq b,$$

or

$$\exists \mathbf{y} : \bar{\mathbf{a}}^T \mathbf{x} + \mathbf{d}^T \mathbf{y} \leq b, \quad \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x}, \quad \mathbf{y} \geq 0.$$

Formulating the Robust Counterpart for Polyhedral Uncertainty Set

- Consider a **linear constraint** in x with coefficients that depend **linearly** on z

$$(\bar{a} + Pz)^T x \leq b, \quad \forall z \in \mathcal{U} \quad (2)$$

- For $\mathcal{U}_{\text{polyhedral}} = \{z : Dz \leq d\}$, satisfying the constraint robustly is equivalent to:

$$\exists y : \bar{a}^T x + d^T y \leq b, \quad D^T y = P^T x, \quad y \geq 0.$$

Remarks.

- To formulate the RC for (2), we must introduce a set of **auxiliary decision variables** y
 - these are **decision variables**, chosen together with x
- How many **auxiliary variables** are needed to derive the RC for (2)?*
- How many **constraints** are needed to derive the RC for (2)?*
- Suppose we were solving $\min_x \{c^T x : Ax \leq b\}$, with $A \in \mathbb{R}^{m \times n}$ being uncertain. Under $\mathcal{U}_{\text{polyhedral}}$ and $D \in \mathbb{R}^{p \times q}$, what kind of problem is the RC of this LO, and how large is it?*

Formulating the Robust Counterpart for Polyhedral Uncertainty Set

- Consider a **linear constraint** in x with coefficients that depend **linearly** on z

$$(\bar{a} + Pz)^T x \leq b, \quad \forall z \in \mathcal{U} \quad (2)$$

- For $\mathcal{U}_{\text{polyhedral}} = \{z : Dz \leq d\}$, satisfying the constraint robustly is equivalent to:

$$\exists y : \bar{a}^T x + d^T y \leq b, \quad D^T y = P^T x, \quad y \geq 0.$$

Remarks.

- To formulate the RC for (2), we must introduce a set of **auxiliary decision variables** y
 - these are **decision variables**, chosen together with x
- How many **auxiliary variables** are needed to derive the RC for (2)?*
 - $\#$ **rows of D**, i.e., as many as the constraints defining $\mathcal{U}_{\text{polyhedral}}$
- How many **constraints** are needed to derive the RC for (2)?*
 - $1 + (\# \text{columns of } D) + (\# \text{rows of } D)$
- Suppose we were solving $\min_x \{c^T x : Ax \leq b\}$, with $A \in \mathbb{R}^{m \times n}$ being uncertain. Under $\mathcal{U}_{\text{polyhedral}}$ and $D \in \mathbb{R}^{p \times q}$, what kind of problem is the RC of this LO, and how large is it?*

Formulating the Robust Counterpart for Polyhedral Uncertainty Set

- Consider a **linear constraint** in x with coefficients that depend **linearly** on z

$$(\bar{a} + Pz)^T x \leq b, \quad \forall z \in \mathcal{U} \quad (2)$$

- For $\mathcal{U}_{\text{polyhedral}} = \{z : Dz \leq d\}$, satisfying the constraint robustly is equivalent to:

$$\exists y : \bar{a}^T x + d^T y \leq b, \quad D^T y = P^T x, \quad y \geq 0.$$

Remarks.

- To formulate the RC for (2), we must introduce a set of **auxiliary decision variables** y
 - these are **decision variables**, chosen together with x
- How many **auxiliary variables** are needed to derive the RC for (2)?*
 - $\#$ **rows of D**, i.e., as many as the constraints defining $\mathcal{U}_{\text{polyhedral}}$
- How many **constraints** are needed to derive the RC for (2)?*
 - $1 + (\# \text{columns of } D) + (\# \text{rows of } D)$
- Suppose we were solving $\min_x \{c^T x : Ax \leq b\}$, with $A \in \mathbb{R}^{m \times n}$ being uncertain. Under $\mathcal{U}_{\text{polyhedral}}$ and $D \in \mathbb{R}^{p \times q}$, what kind of problem is the RC of this LO, and how large is it?*
 - the RC of a **linear optimization** with $\mathcal{U}_{\text{polyhedral}}$ **is still a linear optimization**
 - $n + m \cdot p$ variables, $m \cdot (1 + p + q)$ constraints

Formulating the Robust Counterpart (RC) for Ellipsoidal Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^\top \mathbf{x} \leq b, \quad \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{ellipsoid}} = \{\mathbf{z} : \|\mathbf{z}\|_2 \leq \rho\}$, satisfying the constraint robustly is equivalent to:

Formulating the Robust Counterpart (RC) for Ellipsoidal Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{ellipsoid}} = \{\mathbf{z} : \|\mathbf{z}\|_2 \leq \rho\}$, satisfying the constraint robustly is equivalent to:

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z}: \|\mathbf{z}\|_2 \leq \rho} (\mathbf{P}^T \mathbf{x})^T \mathbf{z} \leq b.$$

Formulating the Robust Counterpart (RC) for Ellipsoidal Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{ellipsoid}} = \{\mathbf{z} : \|\mathbf{z}\|_2 \leq \rho\}$, satisfying the constraint robustly is equivalent to:

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z}: \|\mathbf{z}\|_2 \leq \rho} (\mathbf{P}^T \mathbf{x})^T \mathbf{z} \leq b.$$

Intermezzo: $\max \{ \mathbf{q}^T \mathbf{z} : \|\mathbf{z}\|_2 \leq \rho \}$ or $\max \{ \mathbf{q}^T \mathbf{z} : \mathbf{z}^T \mathbf{z} \leq \rho^2 \}$

Lagrange: $\mathbf{z} = \mathbf{q}/\lambda$, and $\lambda = \|\mathbf{q}\|_2/\rho$.

Optimal objective value: $\frac{\mathbf{q}^T \mathbf{q}}{\lambda} = \rho \|\mathbf{q}\|_2$.

Formulating the Robust Counterpart (RC) for Ellipsoidal Uncertainty Set

- Consider a **linear constraint** in \mathbf{x} with coefficients that depend **linearly** on \mathbf{z}

$$(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \quad \forall \mathbf{z} \in \mathcal{U}$$

- For $\mathcal{U}_{\text{ellipsoid}} = \{\mathbf{z} : \|\mathbf{z}\|_2 \leq \rho\}$, satisfying the constraint robustly is equivalent to:

$$\bar{\mathbf{a}}^T \mathbf{x} + \max_{\mathbf{z} : \|\mathbf{z}\|_2 \leq \rho} (\mathbf{P}^T \mathbf{x})^T \mathbf{z} \leq b.$$

Intermezzo: $\max \{ \mathbf{q}^T \mathbf{z} : \|\mathbf{z}\|_2 \leq \rho \}$ or $\max \{ \mathbf{q}^T \mathbf{z} : \mathbf{z}^T \mathbf{z} \leq \rho^2 \}$

Lagrange: $\mathbf{z} = \mathbf{q}/\lambda$, and $\lambda = \|\mathbf{q}\|_2/\rho$.

Optimal objective value: $\frac{\mathbf{q}^T \mathbf{q}}{\lambda} = \rho \|\mathbf{q}\|_2$.

Hence robust counterpart (RC) is:

$$\bar{\mathbf{a}}^T \mathbf{x} + \rho \|\mathbf{P}^T \mathbf{x}\|_2 \leq b.$$

RC for Linear Optimization Problems with Classical Sets

The robust counterpart for $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$ is:

U-set	\mathcal{U}	Robust Counterpart	Tractability
Box	$\ \mathbf{z}\ _{\infty} \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _1 \leq b$	LO
Ellipsoidal	$\ \mathbf{z}\ _2 \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _2 \leq b$	CQO
Polyhedral	$\mathbf{D}\mathbf{z} \leq \mathbf{d}$	$\exists \mathbf{y} : \begin{cases} \bar{\mathbf{a}}^T \mathbf{x} + \mathbf{d}^T \mathbf{y} \leq b \\ \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x} \\ \mathbf{y} \geq 0 \end{cases}$	LO
Budget	$\begin{cases} \ \mathbf{z}\ _{\infty} \leq \rho \\ \ \mathbf{z}\ _1 \leq \Gamma \end{cases}$	$\exists \mathbf{y} : \bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{y}\ _1 + \Gamma \ \mathbf{P}^T \mathbf{x} - \mathbf{y}\ _{\infty} \leq b$	LO

RC for Linear Optimization Problems with Classical Sets

The robust counterpart for $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$ is:

U-set	\mathcal{U}	Robust Counterpart	Tractability
Box	$\ \mathbf{z}\ _{\infty} \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _1 \leq b$	LO
Ellipsoidal	$\ \mathbf{z}\ _2 \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _2 \leq b$	CQO
Polyhedral	$\mathbf{D}\mathbf{z} \leq \mathbf{d}$	$\exists \mathbf{y} : \begin{cases} \bar{\mathbf{a}}^T \mathbf{x} + \mathbf{d}^T \mathbf{y} \leq b \\ \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x} \\ \mathbf{y} \geq 0 \end{cases}$	LO
Budget	$\begin{cases} \ \mathbf{z}\ _{\infty} \leq \rho \\ \ \mathbf{z}\ _1 \leq \Gamma \end{cases}$	$\exists \mathbf{y} : \bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{y}\ _1 + \Gamma \ \mathbf{P}^T \mathbf{x} - \mathbf{y}\ _{\infty} \leq b$	LO

- Problems above can be handled by large-scale modern solvers: CPLEX, Gurobi, etc.
- Some software now also handling automatic problem re-formulation
- If some of the decisions \mathbf{x} are integer, problems above become MI-LO/CQO
- Already a lot of mileage in many practical problems:
logistics and supply chain management, radiation therapy, scheduling, ...

RC for Linear Optimization Problems with Classical Sets

The robust counterpart for $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$ is:

U-set	\mathcal{U}	Robust Counterpart	Tractability
Box	$\ \mathbf{z}\ _{\infty} \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _1 \leq b$	LO
Ellipsoidal	$\ \mathbf{z}\ _2 \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _2 \leq b$	CQO
Polyhedral	$\mathbf{D}\mathbf{z} \leq \mathbf{d}$	$\exists \mathbf{y} : \begin{cases} \bar{\mathbf{a}}^T \mathbf{x} + \mathbf{d}^T \mathbf{y} \leq b \\ \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x} \\ \mathbf{y} \geq 0 \end{cases}$	LO
Budget	$\begin{cases} \ \mathbf{z}\ _{\infty} \leq \rho \\ \ \mathbf{z}\ _1 \leq \Gamma \end{cases}$	$\exists \mathbf{y} : \bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{y}\ _1 + \Gamma \ \mathbf{P}^T \mathbf{x} - \mathbf{y}\ _{\infty} \leq b$	LO

- Uncertainty in the right-hand side:** $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b + \mathbf{p}^T \mathbf{z}, \forall \mathbf{z} \in \mathcal{U}$

$\Leftrightarrow \bar{\mathbf{a}}^T \mathbf{x} + (\mathbf{P}^T \mathbf{x} - \mathbf{p})^T \mathbf{z} \leq b, \forall \mathbf{z} \in \mathcal{U}$, so can use base model

RC for Linear Optimization Problems with Classical Sets

The robust counterpart for $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$ is:

U-set	\mathcal{U}	Robust Counterpart	Tractability
Box	$\ \mathbf{z}\ _{\infty} \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _1 \leq b$	LO
Ellipsoidal	$\ \mathbf{z}\ _2 \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _2 \leq b$	CQO
Polyhedral	$\mathbf{D}\mathbf{z} \leq \mathbf{d}$	$\exists \mathbf{y} : \begin{cases} \bar{\mathbf{a}}^T \mathbf{x} + \mathbf{d}^T \mathbf{y} \leq b \\ \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x} \\ \mathbf{y} \geq 0 \end{cases}$	LO
Budget	$\begin{cases} \ \mathbf{z}\ _{\infty} \leq \rho \\ \ \mathbf{z}\ _1 \leq \Gamma \end{cases}$	$\exists \mathbf{y} : \bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{y}\ _1 + \Gamma \ \mathbf{P}^T \mathbf{x} - \mathbf{y}\ _{\infty} \leq b$	LO

- Uncertainty in the right-hand side:** $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b + \mathbf{p}^T \mathbf{z}, \forall \mathbf{z} \in \mathcal{U}$
 $\Leftrightarrow \bar{\mathbf{a}}^T \mathbf{x} + (\mathbf{P}^T \mathbf{x} - \mathbf{p})^T \mathbf{z} \leq b, \forall \mathbf{z} \in \mathcal{U}$, so can use base model
- General convex uncertainty set:** $\mathcal{U} = \{\mathbf{z} : h_k(\mathbf{z}) \leq 0, k \in K\}$, $h_k(\cdot)$ convex?
 RC is $\exists \{\mathbf{w}_k, \mathbf{u}_k\}_{k \in K} : \bar{\mathbf{a}}^T \mathbf{x} + \sum_k \mathbf{u}_k h_k^*(\mathbf{w}_k/\mathbf{u}_k) \leq b, \sum_k \mathbf{w}_k = \mathbf{P}^T \mathbf{x}, \mathbf{u} \geq 0$. h_k^* is **convex conjugate** of h_k

RC for Linear Optimization Problems with Classical Sets

The robust counterpart for $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$ is:

U-set	\mathcal{U}	Robust Counterpart	Tractability
Box	$\ \mathbf{z}\ _\infty \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _1 \leq b$	LO
Ellipsoidal	$\ \mathbf{z}\ _2 \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _2 \leq b$	CQO
Polyhedral	$\mathbf{D}\mathbf{z} \leq \mathbf{d}$	$\exists \mathbf{y} : \begin{cases} \bar{\mathbf{a}}^T \mathbf{x} + \mathbf{d}^T \mathbf{y} \leq b \\ \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x} \\ \mathbf{y} \geq 0 \end{cases}$	LO
Budget	$\begin{cases} \ \mathbf{z}\ _\infty \leq \rho \\ \ \mathbf{z}\ _1 \leq \Gamma \end{cases}$	$\exists \mathbf{y} : \bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{y}\ _1 + \Gamma \ \mathbf{P}^T \mathbf{x} - \mathbf{y}\ _\infty \leq b$	LO

- Uncertainty in the right-hand side:** $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b + \mathbf{p}^T \mathbf{z}, \forall \mathbf{z} \in \mathcal{U}$
 $\Leftrightarrow \bar{\mathbf{a}}^T \mathbf{x} + (\mathbf{P}^T \mathbf{x} - \mathbf{p})^T \mathbf{z} \leq b, \forall \mathbf{z} \in \mathcal{U}$, so can use base model
- General convex uncertainty set:** $\mathcal{U} = \{\mathbf{z} : h_k(\mathbf{z}) \leq 0, k \in K\}$, $h_k(\cdot)$ convex?
 RC is $\exists \{\mathbf{w}_k, \mathbf{u}_k\}_{k \in K} : \bar{\mathbf{a}}^T \mathbf{x} + \sum_k \mathbf{u}_k h_k^*(\mathbf{w}_k/\mathbf{u}_k) \leq b, \sum_k \mathbf{w}_k = \mathbf{P}^T \mathbf{x}, \mathbf{u} \geq 0$. h_k^* is **convex conjugate** of h_k
- Constraint LHS general in \mathbf{x} , linear in \mathbf{z} :** $(\mathbf{P}\mathbf{z})^T \mathbf{g}(\mathbf{x}) \leq b, \forall \mathbf{z} \in \mathcal{U}$
 To calculate RC, take $\bar{\mathbf{a}} = 0$ and replace \mathbf{x} with $\mathbf{g}(\mathbf{x})$ in our base-case model

RC for Linear Optimization Problems with Classical Sets

The robust counterpart for $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$ is:

U-set	\mathcal{U}	Robust Counterpart	Tractability
Box	$\ \mathbf{z}\ _\infty \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _1 \leq b$	LO
Ellipsoidal	$\ \mathbf{z}\ _2 \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _2 \leq b$	CQO
Polyhedral	$\mathbf{D}\mathbf{z} \leq \mathbf{d}$	$\exists \mathbf{y} : \begin{cases} \bar{\mathbf{a}}^T \mathbf{x} + \mathbf{d}^T \mathbf{y} \leq b \\ \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x} \\ \mathbf{y} \geq 0 \end{cases}$	LO
Budget	$\begin{cases} \ \mathbf{z}\ _\infty \leq \rho \\ \ \mathbf{z}\ _1 \leq \Gamma \end{cases}$	$\exists \mathbf{y} : \bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{y}\ _1 + \Gamma \ \mathbf{P}^T \mathbf{x} - \mathbf{y}\ _\infty \leq b$	LO

- Uncertainty in the right-hand side:** $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b + \mathbf{p}^T \mathbf{z}, \forall \mathbf{z} \in \mathcal{U}$
 $\Leftrightarrow \bar{\mathbf{a}}^T \mathbf{x} + (\mathbf{P}^T \mathbf{x} - \mathbf{p})^T \mathbf{z} \leq b, \forall \mathbf{z} \in \mathcal{U}$, so can use base model
- General convex uncertainty set:** $\mathcal{U} = \{\mathbf{z} : \mathbf{h}_k(\mathbf{z}) \leq 0, k \in K\}$, $\mathbf{h}_k(\cdot)$ convex?
 RC is $\exists \{\mathbf{w}_k, \mathbf{u}_k\}_{k \in K} : \bar{\mathbf{a}}^T \mathbf{x} + \sum_k \mathbf{u}_k \mathbf{h}_k^*(\mathbf{w}_k/\mathbf{u}_k) \leq b, \sum_k \mathbf{w}_k = \mathbf{P}^T \mathbf{x}, \mathbf{u}_k \geq 0$. \mathbf{h}_k^* is **convex conjugate** of \mathbf{h}_k
- Constraint LHS general in \mathbf{x} , linear in \mathbf{z} :** $(\mathbf{P}\mathbf{z})^T \mathbf{g}(\mathbf{x}) \leq b, \forall \mathbf{z} \in \mathcal{U}$
 To calculate RC, take $\bar{\mathbf{a}} = 0$ and replace \mathbf{x} with $\mathbf{g}(\mathbf{x})$ in our base-case model
- Constraint LHS linear in $\mathbf{x} \geq 0$, concave in \mathbf{z} :** $\mathbf{x}^T \mathbf{g}(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z}) \leq b, \forall \mathbf{z} \in \mathcal{U}$, $\mathbf{g}_i(\mathbf{y})$ concave
 $\Leftrightarrow \mathbf{d}^T \mathbf{x} \leq b, \forall (\mathbf{z}, \mathbf{d}) \in \mathcal{U}^+ := \{(\mathbf{z}, \mathbf{d}) \mid \exists \mathbf{a} : \mathbf{a} = \bar{\mathbf{a}} + \mathbf{P}\mathbf{z}, \mathbf{d} \leq \mathbf{f}(\mathbf{a}), \mathbf{z} \in \mathcal{U}\}$; now **linear** in (\mathbf{z}, \mathbf{d}) , and \mathcal{U}^+ convex

RC for Linear Optimization Problems with Classical Sets

The robust counterpart for $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b, \forall \mathbf{z} \in \mathcal{U}$ is:

U-set	\mathcal{U}	Robust Counterpart	Tractability
Box	$\ \mathbf{z}\ _\infty \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _1 \leq b$	LO
Ellipsoidal	$\ \mathbf{z}\ _2 \leq \rho$	$\bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{P}^T \mathbf{x}\ _2 \leq b$	CQO
Polyhedral	$\mathbf{D}\mathbf{z} \leq \mathbf{d}$	$\exists \mathbf{y} : \begin{cases} \bar{\mathbf{a}}^T \mathbf{x} + \mathbf{d}^T \mathbf{y} \leq b \\ \mathbf{D}^T \mathbf{y} = \mathbf{P}^T \mathbf{x} \\ \mathbf{y} \geq 0 \end{cases}$	LO
Budget	$\begin{cases} \ \mathbf{z}\ _\infty \leq \rho \\ \ \mathbf{z}\ _1 \leq \Gamma \end{cases}$	$\exists \mathbf{y} : \bar{\mathbf{a}}^T \mathbf{x} + \rho \ \mathbf{y}\ _1 + \Gamma \ \mathbf{P}^T \mathbf{x} - \mathbf{y}\ _\infty \leq b$	LO

- Uncertainty in the right-hand side:** $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b + \mathbf{p}^T \mathbf{z}, \forall \mathbf{z} \in \mathcal{U}$
 $\Leftrightarrow \bar{\mathbf{a}}^T \mathbf{x} + (\mathbf{P}^T \mathbf{x} - \mathbf{p})^T \mathbf{z} \leq b, \forall \mathbf{z} \in \mathcal{U}$, so can use base model
- General convex uncertainty set:** $\mathcal{U} = \{\mathbf{z} : \mathbf{h}_k(\mathbf{z}) \leq 0, k \in K\}$, $\mathbf{h}_k(\cdot)$ convex?
 RC is $\exists \{\mathbf{w}_k, \mathbf{u}_k\}_{k \in K} : \bar{\mathbf{a}}^T \mathbf{x} + \sum_k \mathbf{u}_k \mathbf{h}_k^*(\mathbf{w}_k/\mathbf{u}_k) \leq b, \sum_k \mathbf{w}_k = \mathbf{P}^T \mathbf{x}, \mathbf{u}_k \geq 0$. \mathbf{h}_k^* is **convex conjugate** of \mathbf{h}_k
- Constraint LHS general in \mathbf{x} , linear in \mathbf{z} :** $(\mathbf{P}\mathbf{z})^T \mathbf{g}(\mathbf{x}) \leq b, \forall \mathbf{z} \in \mathcal{U}$
 To calculate RC, take $\bar{\mathbf{a}} = 0$ and replace \mathbf{x} with $\mathbf{g}(\mathbf{x})$ in our base-case model
- Constraint LHS linear in $\mathbf{x} \geq 0$, concave in \mathbf{z} :** $\mathbf{x}^T \mathbf{g}(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z}) \leq b, \forall \mathbf{z} \in \mathcal{U}$, $\mathbf{g}_i(\mathbf{y})$ concave
 $\Leftrightarrow \mathbf{d}^T \mathbf{x} \leq b, \forall (\mathbf{z}, \mathbf{d}) \in \mathcal{U}^+ := \{(\mathbf{z}, \mathbf{d}) \mid \exists \mathbf{a} : \mathbf{a} = \bar{\mathbf{a}} + \mathbf{P}\mathbf{z}, \mathbf{d} \leq \mathbf{f}(\mathbf{a}), \mathbf{z} \in \mathcal{U}\}$; now **linear** in (\mathbf{z}, \mathbf{d}) , and \mathcal{U}^+ convex
- Constraint LHS convex in \mathbf{x} and convex in \mathbf{z} :** $\mathbf{f}(\mathbf{x}, \mathbf{z}) \leq b$, \mathbf{f} jointly convex
 Tractable if \mathbf{f} has “easy” piece-wise description: $\mathbf{f}(\mathbf{x}, \mathbf{z}) = \max_{k \in K} \mathbf{f}_k(\mathbf{x}, \mathbf{z})$, where \mathbf{f}_k are cases that “worked”

- **Uncertainty in the right-hand side:** $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq \mathbf{b} + \mathbf{p}^T \mathbf{z}, \forall \mathbf{z} \in \mathcal{U}$
 $\Leftrightarrow \bar{\mathbf{a}}^T \mathbf{x} + (\mathbf{P}^T \mathbf{x} - \mathbf{p})^T \mathbf{z} \leq \mathbf{b}, \forall \mathbf{z} \in \mathcal{U}$, so can use base model

- **Uncertainty in the right-hand side:** $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq \mathbf{b} + \mathbf{p}^T \mathbf{z}, \forall \mathbf{z} \in \mathcal{U}$
 $\Leftrightarrow \bar{\mathbf{a}}^T \mathbf{x} + (\mathbf{P}^T \mathbf{x} - \mathbf{p})^T \mathbf{z} \leq \mathbf{b}, \forall \mathbf{z} \in \mathcal{U}$, so can use base model
- **General convex uncertainty set:** $\mathcal{U} = \{\mathbf{z} : h_k(\mathbf{z}) \leq 0, k \in K\}$, $h_k(\cdot)$ convex?
 $\Leftrightarrow \exists \{\mathbf{w}_k, \mathbf{u}_k\}_{k \in K} : \bar{\mathbf{a}}^T \mathbf{x} + \sum_k \mathbf{u}_k h_k^*(\mathbf{w}_k / \mathbf{u}_k) \leq \mathbf{b}, \sum_k \mathbf{w}_k = \mathbf{P}^T \mathbf{x}, \mathbf{u} \geq 0$.
 h_k^* is **convex conjugate** of h_k

- **Uncertainty in the right-hand side:** $(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq \mathbf{b} + \mathbf{p}^T \mathbf{z}, \forall \mathbf{z} \in \mathcal{U}$
 $\Leftrightarrow \bar{\mathbf{a}}^T \mathbf{x} + (\mathbf{P}^T \mathbf{x} - \mathbf{p})^T \mathbf{z} \leq \mathbf{b}, \forall \mathbf{z} \in \mathcal{U}$, so can use base model
- **General convex uncertainty set:** $\mathcal{U} = \{\mathbf{z} : h_k(\mathbf{z}) \leq 0, k \in K\}$, $h_k(\cdot)$ convex?
 $\Leftrightarrow \exists \{\mathbf{w}_k, \mathbf{u}_k\}_{k \in K} : \bar{\mathbf{a}}^T \mathbf{x} + \sum_k \mathbf{u}_k h_k^*(\mathbf{w}_k / \mathbf{u}_k) \leq \mathbf{b}, \sum_k \mathbf{w}_k = \mathbf{P}^T \mathbf{x}, \mathbf{u} \geq 0$.
 h_k^* is **convex conjugate** of h_k
- **LHS general in \mathbf{x} , linear in \mathbf{z} :** $(\mathbf{P}\mathbf{z})^T \mathbf{g}(\mathbf{x}) \leq \mathbf{b}, \forall \mathbf{z} \in \mathcal{U}$
To calculate RC, take $\bar{\mathbf{a}} = 0$ and replace \mathbf{x} with $\mathbf{g}(\mathbf{x})$ in our base-case model

- **Uncertainty in the right-hand side:** $(\bar{a} + Pz)^T x \leq b + p^T z, \forall z \in \mathcal{U}$
 $\Leftrightarrow \bar{a}^T x + (P^T x - p)^T z \leq b, \forall z \in \mathcal{U}$, so can use base model
- **General convex uncertainty set:** $\mathcal{U} = \{z : h_k(z) \leq 0, k \in K\}$, $h_k(\cdot)$ convex?
 $\Leftrightarrow \exists \{w_k, u_k\}_{k \in K} : \bar{a}^T x + \sum_k u_k h_k^*(w_k/u_k) \leq b, \sum_k w_k = P^T x, u \geq 0$.
 h_k^* is **convex conjugate** of h_k
- **LHS general in x , linear in z :** $(Pz)^T g(x) \leq b, \forall z \in \mathcal{U}$
To calculate RC, take $\bar{a} = 0$ and replace x with $g(x)$ in our base-case model
- **LHS linear in $x \geq 0$, concave in z :** $x^T g(\bar{a} + Pz) \leq b, \forall z \in \mathcal{U}$, g concave
 $\Leftrightarrow d^T x \leq b, \forall (z, d) \in \mathcal{U}^+ := \{(z, d) \mid \exists a : a = \bar{a} + Pz, d \leq f(a), z \in \mathcal{U}\}$
now **linear** in (z, d) , and \mathcal{U}^+ convex

- Uncertainty in the right-hand side:** $(\bar{a} + Pz)^T x \leq b + p^T z, \forall z \in \mathcal{U}$
 $\Leftrightarrow \bar{a}^T x + (P^T x - p)^T z \leq b, \forall z \in \mathcal{U}$, so can use base model
- General convex uncertainty set:** $\mathcal{U} = \{z : h_k(z) \leq 0, k \in K\}$, $h_k(\cdot)$ convex?
 $\Leftrightarrow \exists \{w_k, u_k\}_{k \in K} : \bar{a}^T x + \sum_k u_k h_k^*(w_k/u_k) \leq b, \sum_k w_k = P^T x, u \geq 0$.
 h_k^* is **convex conjugate** of h_k
- LHS general in x , linear in z :** $(Pz)^T g(x) \leq b, \forall z \in \mathcal{U}$
 To calculate RC, take $\bar{a} = 0$ and replace x with $g(x)$ in our base-case model
- LHS linear in $x \geq 0$, concave in z :** $x^T g(\bar{a} + Pz) \leq b, \forall z \in \mathcal{U}$, g concave
 $\Leftrightarrow d^T x \leq b, \forall (z, d) \in \mathcal{U}^+ := \{(z, d) \mid \exists a : a = \bar{a} + Pz, d \leq f(a), z \in \mathcal{U}\}$
 now **linear** in (z, d) , and \mathcal{U}^+ convex
- LHS convex in x and convex in z :** $f(x, z) \leq b$, f jointly convex

Tractable if f has "easy" piece-wise description: $f(x, z) = \max_{k \in K} f_k(x, z)$, where f_k are cases that "worked"

Used in many applications

- inventory management e.g., [Ben-Tal et al., 2005, Bertsimas and Thiele, 2006, Bienstock and Özbay, 2008, ...]
- facility location and transportation [Baron et al., 2011, ...]
- scheduling [Lin et al., 2004, Yamashita et al., 2007, Mittal et al., 2014, ...]
- revenue management [Perakis and Roels, 2010, Adida and Perakis, 2006, ...]
- project management [Wiesemann et al., 2012, Ben-Tal et al., 2009, ...]
- energy generation and distribution [Zhao et al., 2013, Lorca and Sun, 2015, ...]
- portfolio optimization [Goldfarb and Iyengar, 2003, Tütüncü and Koenig, 2004, Ceria and Stubbs, 2006, Pinar and Tütüncü, 2005, Bertsimas and Pachamanova, 2008, ...]
- healthcare [Borfeld et al., 2008, Hanne et al., 2009, Chen et al., 2011, I., Trichakis, Yoon (2018), ...]
- humanitarian [Uichano 2017, den Hertog et al., 2019, ...]

A Quick Example: A Facility Location Problem (Baron et al. 2011)

Need to decide where to open facilities, how much capacity to install, and how to assign customer demands over a future planning horizon, in order to maximize profits.

A Quick Example: A Facility Location Problem (Baron et al. 2011)

Need to decide where to open facilities, how much capacity to install, and how to assign customer demands over a future planning horizon, in order to maximize profits.

Step 1. Start with a **deterministic** model formulation:

A Quick Example: A Facility Location Problem (Baron et al. 2011)

Need to decide where to open facilities, how much capacity to install, and how to assign customer demands over a future planning horizon, in order to maximize profits.

Step 1. Start with a **deterministic** model formulation:

$$\begin{aligned} \max_{X, I, Z, P} \quad & \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} \sum_{j \in \mathcal{N}} (\eta - d_{ij}) X_{ij\tau} D_{j\tau} - \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} c_i P_{i\tau} - \sum_{i \in \mathcal{F}} (C_i Z_i + K_i I_i) \\ \text{s.t.} \quad & \sum_{i \in \mathcal{F}} X_{ij\tau} \leq 1, \quad j \in \mathcal{N}, \tau \in \mathcal{T}, \\ & \sum_{j \in \mathcal{N}} X_{ij\tau} D_{j\tau} \leq P_{i\tau}, \quad i \in \mathcal{F}, \tau \in \mathcal{T}, \\ & X_{ij\tau} \geq 0, \quad i \in \mathcal{F}, j \in \mathcal{N}, \tau \in \mathcal{T} \\ & P_{i\tau} \leq Z_i, \quad Z_i \leq M \cdot I_i, \quad i \in \mathcal{F}, \tau \in \mathcal{T} \\ & I \in \{0, 1\}^{|\mathcal{F}|}, \quad (\text{where } M \text{ is a large enough constant.}) \end{aligned}$$

Parameters:

\mathcal{T} : discrete planning horizon, indexed by τ
 \mathcal{F} : potential facility locations, indexed by i
 \mathcal{N} : demand node locations, indexed by j
 η : unit price of goods
 c_i : cost per unit of production at facility i
 C_i : cost per unit of capacity for facility i
 K_i : cost of opening a facility at location i
 d_{ij} : cost of shipping units from location i to j
 $D_{j\tau}$: demand in period τ at location j .

Decision variables:

$X_{ij\tau}$: fraction of demand j in period τ satisfied by i
 $P_{i\tau}$: quantity produced at facility i in period τ
 I_i : whether facility i is open (0/1)
 Z_i : capacity of facility i if open.

A Quick Example: A Facility Location Problem (Baron et al. 2011)

Need to decide where to open facilities, how much capacity to install, and how to assign customer demands over a future planning horizon, in order to maximize profits.

Step 1. Start with a **deterministic** model formulation:

$$\begin{aligned} \max_{X, I, Z, P} \quad & \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} \sum_{j \in \mathcal{N}} (\eta - d_{ij}) X_{ij\tau} D_{j\tau} - \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} c_i P_{i\tau} - \sum_{i \in \mathcal{F}} (C_i Z_i + K_i I_i) \\ \text{s.t.} \quad & \sum_{i \in \mathcal{F}} X_{ij\tau} \leq 1, \quad j \in \mathcal{N}, \tau \in \mathcal{T}, \\ & \sum_{j \in \mathcal{N}} X_{ij\tau} D_{j\tau} \leq P_{i\tau}, \quad i \in \mathcal{F}, \tau \in \mathcal{T}, \\ & X_{ij\tau} \geq 0, \quad i \in \mathcal{F}, j \in \mathcal{N}, \tau \in \mathcal{T} \\ & P_{i\tau} \leq Z_i, \quad Z_i \leq M \cdot I_i, \quad i \in \mathcal{F}, \tau \in \mathcal{T} \\ & I \in \{0, 1\}^{|\mathcal{F}|}, \quad (\text{where } M \text{ is a large enough constant.}) \end{aligned}$$

Step 2. Identify all uncertain parameters and **model** the uncertainty set \mathcal{U} .

Baron et al. 2011 captured uncertain demands:

$$\mathcal{U} = \left\{ \mathbf{D} \in \mathbb{R}^{|\mathcal{N}| \cdot |\mathcal{T}|} \left| \sum_{j \in \mathcal{N}} \sum_{t \in \mathcal{T}} \left(\frac{\mathbf{D}_{jt} - \bar{\mathbf{D}}_{jt}}{\epsilon_t \bar{\mathbf{D}}_{jt}} \right)^2 \leq \rho^2 \right. \right\},$$

$\{\bar{\mathbf{D}}_{jt}\}_{j \in \mathcal{N}; t \in \mathcal{T}}$ are “nominal” demands, ϵ_t is allowed deviation (%), ρ is the size of the ellipsoid.

A Quick Example: A Facility Location Problem (Baron et al. 2011)

Need to decide where to open facilities, how much capacity to install, and how to assign customer demands over a future planning horizon, in order to maximize profits.

Step 1. Start with a **deterministic** model formulation:

$$\begin{aligned} \max_{X, I, Z, P} \quad & \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} \sum_{j \in \mathcal{N}} (\eta - d_{ij}) X_{ij\tau} D_{j\tau} - \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} c_i P_{i\tau} - \sum_{i \in \mathcal{F}} (C_i Z_i + K_i I_i) \\ \text{s.t.} \quad & \sum_{i \in \mathcal{F}} X_{ij\tau} \leq 1, \quad j \in \mathcal{N}, \tau \in \mathcal{T}, \\ & \sum_{j \in \mathcal{N}} X_{ij\tau} D_{j\tau} \leq P_{i\tau}, \quad i \in \mathcal{F}, \tau \in \mathcal{T}, \\ & X_{ij\tau} \geq 0, \quad i \in \mathcal{F}, j \in \mathcal{N}, \tau \in \mathcal{T} \\ & P_{i\tau} \leq Z_i, \quad Z_i \leq M \cdot I_i, \quad i \in \mathcal{F}, \tau \in \mathcal{T} \\ & I \in \{0, 1\}^{|\mathcal{F}|}, \quad (\text{where } M \text{ is a large enough constant.}) \end{aligned}$$

Step 2. Identify all uncertain parameters and **model** the uncertainty set \mathcal{U} .

Baron et al. 2011 captured uncertain demands:

$$\mathcal{U} = \left\{ \mathbf{D} \in \mathbb{R}^{|\mathcal{N}| \cdot |\mathcal{T}|} \left| \sum_{j \in \mathcal{N}} \sum_{t \in \mathcal{T}} \left(\frac{\mathbf{D}_{jt} - \bar{\mathbf{D}}_{jt}}{\epsilon_t \bar{\mathbf{D}}_{jt}} \right)^2 \leq \rho^2 \right. \right\},$$

Equivalently, can write $D_{jt} = \bar{D}_{jt}(1 + \epsilon_t \cdot \mathbf{z}_{jt})$, where $\mathbf{z} \in \mathcal{U} = \{\mathbf{z} \in \mathbb{R}^{|\mathcal{N}| \cdot |\mathcal{T}|} : \|\mathbf{z}\|_2 \leq \rho\}$

A Quick Example: A Facility Location Problem (Baron et al. 2011)

Need to decide where to open facilities, how much capacity to install, and how to assign customer demands over a future planning horizon, in order to maximize profits.

Step 1. Start with a **deterministic** model formulation:

$$\begin{aligned} \max_{X, I, Z, P} \quad & \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} \sum_{j \in \mathcal{N}} (\eta - d_{ij}) X_{ij\tau} D_{j\tau} - \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} c_i P_{i\tau} - \sum_{i \in \mathcal{F}} (C_i Z_i + K_i I_i) \\ \text{s.t.} \quad & \sum_{i \in \mathcal{F}} X_{ij\tau} \leq 1, \quad j \in \mathcal{N}, \tau \in \mathcal{T}, \\ & \sum_{j \in \mathcal{N}} X_{ij\tau} D_{j\tau} \leq P_{i\tau}, \quad i \in \mathcal{F}, \tau \in \mathcal{T}, \\ & X_{ij\tau} \geq 0, \quad i \in \mathcal{F}, j \in \mathcal{N}, \tau \in \mathcal{T} \\ & P_{i\tau} \leq Z_i, \quad Z_i \leq M \cdot I_i, \quad i \in \mathcal{F}, \tau \in \mathcal{T} \\ & I \in \{0, 1\}^{|\mathcal{F}|}, \quad (\text{where } M \text{ is a large enough constant.}) \end{aligned}$$

Step 2. Identify all uncertain parameters and **model** the uncertainty set \mathcal{U} .

Baron et al. 2011 captured uncertain demands:

$$\mathcal{U} = \left\{ \mathbf{D} \in \mathbb{R}^{|\mathcal{N}| \cdot |\mathcal{T}|} \left| \sum_{j \in \mathcal{N}} \sum_{t \in \mathcal{T}} \left(\frac{\mathbf{D}_{jt} - \bar{\mathbf{D}}_{jt}}{\epsilon_t \bar{\mathbf{D}}_{jt}} \right)^2 \leq \rho^2 \right. \right\},$$

Step 3. Derive robust counterpart for the problem. Here, this will be a Conic Quadratic program.

Caution when Building Robust Models:

Equivalent deterministic/nominal models may lead to different robust counterparts!

Caution when Building Robust Models:
 Equivalent deterministic/nominal models may lead to different robust counterparts!

Our initial **deterministic** model:

$$\begin{aligned}
 \max_{X, I, Z, P} \quad & \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} \sum_{j \in \mathcal{N}} (\eta - d_{ij}) X_{ij\tau} D_{j\tau} - \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} c_i P_{i\tau} - \sum_{i \in \mathcal{F}} (C_i Z_i - K_i I_i) \\
 \text{s.t.} \quad & \sum_{i \in \mathcal{F}} X_{ij\tau} \leq 1, \quad j \in \mathcal{N}, \tau \in \mathcal{T}, \\
 & \sum_{j \in \mathcal{N}} X_{ij\tau} D_{j\tau} \leq P_{i\tau}, \quad i \in \mathcal{F}, \tau \in \mathcal{T}, \\
 & X_{ij\tau} \geq 0, \quad i \in \mathcal{F}, j \in \mathcal{N}, \tau \in \mathcal{T} \\
 & P_{i\tau} \leq Z_i, \quad Z_i \leq M \cdot I_i, \quad i \in \mathcal{F}, \tau \in \mathcal{T} \\
 & I_i \in \{0, 1\}^{|\mathcal{T}|}, \quad (\text{where } M \text{ is a large enough constant.})
 \end{aligned}$$

An equivalent **deterministic** model, with decisions for quantities:

$$\begin{aligned}
 \max_{Q, I, Z, P} \quad & \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} \sum_{j \in \mathcal{N}} (\eta - d_{ij}) Y_{ij\tau} - \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} c_i P_{i\tau} - \sum_{i \in \mathcal{F}} (C_i Z_i - K_i I_i) \\
 \text{s.t.} \quad & \sum_{i \in \mathcal{F}} Y_{ij\tau} \leq D_{j\tau}, \quad j \in \mathcal{N}, \tau \in \mathcal{T}, \\
 & \sum_{j \in \mathcal{N}} Y_{ij\tau} \leq P_{i\tau}, \quad i \in \mathcal{F}, \tau \in \mathcal{T}, \\
 & Y_{ij\tau} \geq 0, \quad i \in \mathcal{F}, j \in \mathcal{N}, \tau \in \mathcal{T} \\
 & P_{i\tau} \leq Z_i, \quad Z_i \leq M \cdot I_i, \quad i \in \mathcal{F}, \tau \in \mathcal{T} \\
 & I_i \in \{0, 1\}^{|\mathcal{T}|}, \quad (\text{where } M \text{ is a large enough constant.})
 \end{aligned}$$

Caution when Building Robust Models:
 Equivalent deterministic/nominal models may lead to different robust counterparts!

Our initial **deterministic** model:

$$\begin{aligned}
 \max_{X, I, Z, P} \quad & \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} \sum_{j \in \mathcal{N}} (\eta - d_{ij}) X_{ij\tau} D_{j\tau} - \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} c_i P_{i\tau} - \sum_{i \in \mathcal{F}} (C_i Z_i - K_i I_i) \\
 \text{s.t.} \quad & \sum_{i \in \mathcal{F}} X_{ij\tau} \leq 1, \quad j \in \mathcal{N}, \tau \in \mathcal{T}, \\
 & \sum_{j \in \mathcal{N}} X_{ij\tau} D_{j\tau} \leq P_{i\tau}, \quad i \in \mathcal{F}, \tau \in \mathcal{T}, \\
 & X_{ij\tau} \geq 0, \quad i \in \mathcal{F}, j \in \mathcal{N}, \tau \in \mathcal{T} \\
 & P_{i\tau} \leq Z_i, \quad Z_i \leq M \cdot I_i, \quad i \in \mathcal{F}, \tau \in \mathcal{T} \\
 & I_i \in \{0, 1\}^{|\mathcal{F}|}, \quad (\text{where } M \text{ is a large enough constant.})
 \end{aligned}$$

An equivalent **deterministic** model, with decisions for quantities:

$$\begin{aligned}
 \max_{Q, I, Z, P} \quad & \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} \sum_{j \in \mathcal{N}} (\eta - d_{ij}) Y_{ij\tau} - \sum_{\tau \in \mathcal{T}} \sum_{i \in \mathcal{F}} c_i P_{i\tau} - \sum_{i \in \mathcal{F}} (C_i Z_i - K_i I_i) \\
 \text{s.t.} \quad & \sum_{i \in \mathcal{F}} Y_{ij\tau} \leq D_{j\tau}, \quad j \in \mathcal{N}, \tau \in \mathcal{T}, \\
 & \sum_{j \in \mathcal{N}} Y_{ij\tau} \leq P_{i\tau}, \quad i \in \mathcal{F}, \tau \in \mathcal{T}, \\
 & Y_{ij\tau} \geq 0, \quad i \in \mathcal{F}, j \in \mathcal{N}, \tau \in \mathcal{T} \\
 & P_{i\tau} \leq Z_i, \quad Z_i \leq M \cdot I_i, \quad i \in \mathcal{F}, \tau \in \mathcal{T} \\
 & I_i \in \{0, 1\}^{|\mathcal{F}|}, \quad (\text{where } M \text{ is a large enough constant.})
 \end{aligned}$$

Are the **robust counterparts** of the two formulations **equivalent**? Which do you think will be **more conservative**?

HINT: In which model are future shipping decisions more “flexible,” e.g., allowed to depend on realized demands?

Are Robust Solutions **Pareto-Efficient**?

$$\max_{x \in \mathcal{X}} \min_{u \in \mathcal{U}} u^T x$$

- Feasible set of solutions $\mathcal{X} = \{x \in \mathbb{R}^n : Ax \leq b\}$
- Uncertainty set of objective coefficients $\mathcal{U} = \{u \in \mathbb{R}^n : Du \geq d\}$

Are Robust Solutions **Pareto-Efficient**?

$$\max_{x \in \mathcal{X}} \min_{u \in \mathcal{U}} u^T x$$

- Feasible set of solutions $\mathcal{X} = \{x \in \mathbb{R}^n : Ax \leq b\}$
- Uncertainty set of objective coefficients $\mathcal{U} = \{u \in \mathbb{R}^n : Du \geq d\}$
- Classical RO framework results in
 - Optimal value J_{RO}^*
 - Set of robustly optimal solutions

$$X^{RO} = \left\{ x \in \mathcal{X} : \exists y \geq 0 \text{ such that } D^T y = x, \quad y^T d \geq J_{RO}^* \right\}$$

Set of **Robustly Optimal** Solutions

- $X^{\text{RO}} = \{x \in \mathcal{X} : \exists y \geq 0 \text{ such that } D^T y = x, \quad y^T d \geq J_{\text{RO}}^*\}$
- $x \in X^{\text{RO}}$ guarantees that no other solution exists with higher **worst-case** objective value $u^T x$

Set of Robustly Optimal Solutions

- $X^{\text{RO}} = \{x \in \mathcal{X} : \exists y \geq 0 \text{ such that } D^T y = x, \quad y^T d \geq J_{\text{RO}}^*\}$
- $x \in X^{\text{RO}}$ guarantees that no other solution exists with higher **worst-case** objective value $u^T x$
- What if an uncertainty scenario materializes that does not correspond to the worst-case?
- Are there any guarantees that no other solution \bar{x} exists that, apart from protecting us from worst-case scenarios, also performs better overall, under all possible uncertainty realizations?

$$\max_{x \in \mathcal{X}} \min_{u \in \mathcal{U}} u^T x \quad (3)$$

Definition

A solution x is called a **Pareto Robustly Optimal (PRO) solution** for Problem (3) if

- (a) it is robustly optimal, i.e., $x \in X^{\text{RO}}$, and
- (b) there is no $\bar{x} \in \mathcal{X}$ such that

$$u^T \bar{x} \geq u^T x, \quad \forall u \in \mathcal{U}, \quad \text{and}$$

$$\bar{u}^T \bar{x} > \bar{u}^T x, \quad \text{for some } \bar{u} \in \mathcal{U}.$$

Pareto Robustly Optimal solutions (Iancu & Trichakis 2014)

$$\max_{x \in \mathcal{X}} \min_{u \in \mathcal{U}} u^T x \quad (3)$$

Definition

A solution x is called a **Pareto Robustly Optimal (PRO) solution** for Problem (3) if

- (a) it is robustly optimal, i.e., $x \in X^{\text{RO}}$, and
- (b) there is no $\bar{x} \in \mathcal{X}$ such that

$$u^T \bar{x} \geq u^T x, \quad \forall u \in \mathcal{U}, \quad \text{and}$$

$$\bar{u}^T \bar{x} > \bar{u}^T x, \quad \text{for some } \bar{u} \in \mathcal{U}.$$

- $X^{\text{PRO}} \subseteq X^{\text{RO}}$: set of all PRO solutions

Some questions

- Given a RO solution, is it also PRO?
- How can one find a PRO solution?
- Can we optimize over \mathcal{X}^{PRO} ?
- Can we characterize \mathcal{X}^{PRO} ?
 - Is it non-empty?
 - Is it convex?
 - When is $\mathcal{X}^{\text{PRO}} = \mathcal{X}^{\text{RO}}$?
- How does the notion generalize in other RO formulations?

Theorem

Given a solution $x \in X^{\text{RO}}$ and an arbitrary point $\bar{p} \in \text{ri}(\mathcal{U})$, consider the following linear optimization problem:

$$\begin{array}{ll}\text{maximize} & \bar{p}^\top y \\ \text{subject to} & y \in \mathcal{U}^* \\ & x + y \in \mathcal{X}.\end{array}$$

Then, either

- $\mathcal{U}^* \stackrel{\text{def}}{=} \{y \in \mathbb{R}^n : y^\top u \geq 0, \forall u \in \mathcal{U}\}$ is the dual of \mathcal{U}

Theorem

Given a solution $x \in X^{\text{RO}}$ and an arbitrary point $\bar{p} \in \text{ri}(\mathcal{U})$, consider the following linear optimization problem:

$$\begin{array}{ll}\text{maximize} & \bar{p}^\top y \\ \text{subject to} & y \in \mathcal{U}^* \\ & x + y \in \mathcal{X}.\end{array}$$

Then, either

- the optimal value is zero and $x \in X^{\text{PRO}}$, or
- $\mathcal{U}^* \stackrel{\text{def}}{=} \{y \in \mathbb{R}^n : y^\top u \geq 0, \forall u \in \mathcal{U}\}$ is the dual of \mathcal{U}

Theorem

Given a solution $x \in X^{\text{RO}}$ and an arbitrary point $\bar{p} \in \text{ri}(\mathcal{U})$, consider the following linear optimization problem:

$$\begin{array}{ll}\text{maximize} & \bar{p}^\top y \\ \text{subject to} & y \in \mathcal{U}^* \\ & x + y \in \mathcal{X}.\end{array}$$

Then, either

- the optimal value is zero and $x \in X^{\text{PRO}}$, or
 - the optimal value is strictly positive and $\bar{x} = x + y^* \in X^{\text{PRO}}$, for any optimal y^* .
-
- $\mathcal{U}^* \stackrel{\text{def}}{=} \{y \in \mathbb{R}^n : y^\top u \geq 0, \forall u \in \mathcal{U}\}$ is the dual of \mathcal{U}

- Finding a point $\bar{u} \in \text{ri}(\mathcal{U})$ can be done efficiently using LP techniques
- Testing whether $x \in X^{\text{RO}}$ is no harder than solving the classical RO problem in this setting
- Finding a PRO solution $x \in X^{\text{PRO}}$ is no harder than solving the classical RO problem in this setting

- If $\bar{u} \in \text{ri}(\mathcal{U})$, all optimal solutions to the problem below are PRO:

$$\begin{array}{ll}\text{maximize} & \bar{u}^\top x \\ \text{subject to} & x \in X^{\text{RO}}\end{array}$$

- If $0 \in \text{ri}(\mathcal{U})$, then $X^{\text{PRO}} = X^{\text{RO}}$
- If $\bar{u} \in \text{ri}(\mathcal{U})$, then $X^{\text{PRO}} = X^{\text{RO}}$ if and only if the optimal value of this LP is zero:

$$\begin{array}{ll}\text{maximize} & \bar{u}^\top y \\ \text{subject to} & x \in X^{\text{RO}} \\ & y \in \mathcal{U}^* \\ & x + y \in \mathcal{X}\end{array}$$

Optimizing over / Understanding X^{PRO}

- Secondary objective r : can we solve

$$\begin{array}{ll} \text{maximize} & r^{\top} x \\ \text{subject to} & x \in X^{\text{PRO}}? \end{array}$$

- Interesting case: $X^{\text{RO}} \neq X^{\text{PRO}}$

- Secondary objective r : can we solve

$$\begin{array}{ll} \text{maximize} & r^T x \\ \text{subject to} & x \in X^{\text{PRO}}? \end{array}$$

Proposition

X^{PRO} is not necessarily convex.

- $\mathcal{X} = \{x \in \mathbb{R}_+^4 : x_1 \leq 1, x_2 + x_3 \leq 6, x_3 + x_4 \leq 5, x_2 + x_4 \leq 5\}$
- $\mathcal{U} = \text{conv}(\{e_i, i \in \{1, \dots, 4\}\})$
- $J_{\text{RO}}^* = 1$, and $X^{\text{RO}} = \{x \in \mathcal{X} : x \geq 1\}$
- $x^1 = [1 \ 2 \ 4 \ 1]^T$, $x^2 = [1 \ 4 \ 2 \ 1]^T \in X^{\text{PRO}}$
- $0.5x^1 + 0.5x^2$ is Pareto dominated by $[1 \ 3 \ 3 \ 2]^T \in X^{\text{RO}}$.

- Secondary objective r : can we solve

$$\begin{array}{ll}\text{maximize} & r^T x \\ \text{subject to} & x \in X^{\text{PRO}}?\end{array}$$

Proposition

If $X^{\text{RO}} \neq X^{\text{PRO}}$, then $X^{\text{PRO}} \cap \text{ri}(X^{\text{RO}}) = \emptyset$.

- Whether solution to nominal RO is PRO depends on algorithm used for solving LP
- Simplex **better for RO problems** than interior point methods

What Are The Gains?

Example (Portfolio)

- $n + 1$ assets, with returns r_i
- $r_i = \mu_i + \sigma_i \zeta_i$, $i = 1, \dots, n$, $r_{n+1} = \mu_{n+1}$
- ζ unknown, $\mathcal{U} = \{\zeta \in \mathbb{R}^n : -1 \leq \zeta \leq 1, \mathbf{1}^T \zeta = 0\}$
- Objective: select weights x to maximize worst-case portfolio return

What Are The Gains?

Example (Portfolio)

- $n + 1$ assets, with returns r_i
- $r_i = \mu_i + \sigma_i \zeta_i$, $i = 1, \dots, n$, $r_{n+1} = \mu_{n+1}$
- ζ unknown, $\mathcal{U} = \{\zeta \in \mathbb{R}^n : -1 \leq \zeta \leq 1, \mathbf{1}^T \zeta = 0\}$
- Objective: select weights x to maximize worst-case portfolio return

What Are The Gains?

Example (Portfolio)

- $n + 1$ assets, with returns r_i
- $r_i = \mu_i + \sigma_i \zeta_i$, $i = 1, \dots, n$, $r_{n+1} = \mu_{n+1}$
- ζ unknown, $\mathcal{U} = \{\zeta \in \mathbb{R}^n : -1 \leq \zeta \leq 1, \mathbf{1}^T \zeta = 0\}$
- Objective: select weights x to maximize worst-case portfolio return

Example (Inventory)



- One warehouse, N retailers where uncertain demand is realized
- Transportation, holding costs and profit margins differ for each retailer
- Demand driven by market factors $d_i = d_i^0 + q_i^T z$, $i = 1, \dots, N$
- Market factors z are uncertain

$$z \in \mathcal{U} = \{z \in \mathbb{R}^N : -b \cdot \mathbf{1} \leq z \leq b \cdot \mathbf{1}, -B \leq \mathbf{1}^T z \leq B\}$$

Numerical experiments

Example (Project management)

- A PERT diagram given by directed, acyclic graph $G = (\mathcal{N}, \mathcal{E})$
- \mathcal{N} are project events, \mathcal{E} are project activities / tasks



Example (Project management)

- A PERT diagram given by directed, acyclic graph $G = (\mathcal{N}, \mathcal{E})$
- \mathcal{N} are project events, \mathcal{E} are project activities / tasks
- Task $e \in \mathcal{E}$ has uncertain duration $\tau_e = \tau_e^0 + \delta_e$

$$\delta \in \mathcal{U} := \{\delta \in \mathbb{R}_+^{|\mathcal{E}|} : \delta \leq b \cdot \mathbf{1}, \quad \mathbf{1}^\top \delta_e \leq B\}$$

- Task $e \in \mathcal{E}$ can be expedited by allocating a budgeted resource x_e

$$\begin{aligned}\tau_e &= \tau_e^0 + \delta_e - x_e \\ \mathbf{1}^\top x &\leq C\end{aligned}$$

- Goal: find resource allocation x to minimize worst-case completion time

Results – finance and inventory examples (10K instances)






Figure: TOP: portfolio example. BOTTOM: inventory example. LEFT: performance gains in nominal scenario. RIGHT: maximal performance gains.

Results – two project management networks (10K instances)



Careful To Avoid Naïve Inefficiencies In Robust Models!

How to Calibrate Uncertainty Sets?

We know how to derive RC for “structured” uncertainty sets. **But how to pick ρ, Γ ?**

How to Calibrate Uncertainty Sets?

We know how to derive RC for “structured” uncertainty sets. **But how to pick ρ, Γ ?**

Let's take a **probabilistic** view for a moment:

Suppose \mathbf{z}_i are really random, and we seek ρ, Γ to ensure $\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^\top \mathbf{x} \leq b]$ is “large”, $\forall \mathbb{P}$

How to Calibrate Uncertainty Sets?

We know how to derive RC for “structured” uncertainty sets. **But how to pick ρ, Γ ?**

Let's take a **probabilistic** view for a moment:

Suppose \mathbf{z}_i are really random, and we seek ρ, Γ to ensure $\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^\top \mathbf{x} \leq b]$ is “large”, $\forall \mathbb{P}$

Theorem (High Probability of Constraint Satisfaction)

Suppose z_i are independent r.v. with mean 0 and support on $[-1, 1]$. Then:

How to Calibrate Uncertainty Sets?

We know how to derive RC for “structured” uncertainty sets. **But how to pick ρ, Γ ?**

Let's take a **probabilistic** view for a moment:

Suppose \mathbf{z}_i are really random, and we seek ρ, Γ to ensure $\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b]$ is “large”, $\forall \mathbb{P}$

Theorem (High Probability of Constraint Satisfaction)

Suppose z_i are independent r.v. with mean 0 and support on $[-1, 1]$. Then:

- if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{box}} \stackrel{\text{def}}{=} \{\mathbf{z} : \|\mathbf{z}\|_{\infty} \leq 1\}$, $\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] = 1$.
(no need for independence or 0-mean)

How to Calibrate Uncertainty Sets?

We know how to derive RC for “structured” uncertainty sets. **But how to pick ρ, Γ ?**

Let's take a **probabilistic** view for a moment:

Suppose \mathbf{z}_i are really random, and we seek ρ, Γ to ensure $\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b]$ is “large”, $\forall \mathbb{P}$

Theorem (High Probability of Constraint Satisfaction)

Suppose z_i are independent r.v. with mean 0 and support on $[-1, 1]$. Then:

- if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{box}} \stackrel{\text{def}}{=} \{\mathbf{z} : \|\mathbf{z}\|_{\infty} \leq 1\}$, $\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] = 1$.
(no need for independence or 0-mean)
- if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{ellipsoid}} \stackrel{\text{def}}{=} \{\mathbf{z} : \|\mathbf{z}\|_2 \leq \rho \stackrel{\text{def}}{=} \sqrt{2 \ln(1/\epsilon)}\}$,
$$\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] \geq 1 - \epsilon.$$

How to Calibrate Uncertainty Sets?

We know how to derive RC for “structured” uncertainty sets. **But how to pick ρ, Γ ?**

Let's take a **probabilistic** view for a moment:

Suppose \mathbf{z}_i are really random, and we seek ρ, Γ to ensure $\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b]$ is “large”, $\forall \mathbb{P}$

Theorem (High Probability of Constraint Satisfaction)

Suppose z_i are independent r.v. with mean 0 and support on $[-1, 1]$. Then:

- if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{box}} \stackrel{\text{def}}{=} \{\mathbf{z} : \|\mathbf{z}\|_{\infty} \leq 1\}$, $\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] = 1$.
(no need for independence or 0-mean)
- if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{ellipsoid}} \stackrel{\text{def}}{=} \{\mathbf{z} : \|\mathbf{z}\|_2 \leq \rho \stackrel{\text{def}}{=} \sqrt{2 \ln(1/\epsilon)}\}$,
$$\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] \geq 1 - \epsilon.$$
- if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{ellipsoid-box}} \stackrel{\text{def}}{=} \{\mathbf{z} : \|\mathbf{z}\|_2 \leq \rho \stackrel{\text{def}}{=} \sqrt{2 \ln(1/\epsilon)}, \|\mathbf{z}\|_{\infty} \leq 1\}$,
$$\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] \geq 1 - \epsilon.$$

How to Calibrate Uncertainty Sets?

We know how to derive RC for “structured” uncertainty sets. **But how to pick ρ, Γ ?**

Let's take a **probabilistic** view for a moment:

Suppose \mathbf{z}_i are really random, and we seek ρ, Γ to ensure $\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b]$ is “large”, $\forall \mathbb{P}$

Theorem (High Probability of Constraint Satisfaction)

Suppose \mathbf{z}_i are independent r.v. with mean 0 and support on $[-1, 1]$. Then:

- if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{box}} \stackrel{\text{def}}{=} \{\mathbf{z} : \|\mathbf{z}\|_{\infty} \leq 1\}$, $\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] = 1$.
(no need for independence or 0-mean)
- if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{ellipsoid}} \stackrel{\text{def}}{=} \{\mathbf{z} : \|\mathbf{z}\|_2 \leq \rho \stackrel{\text{def}}{=} \sqrt{2 \ln(1/\epsilon)}\}$,
$$\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] \geq 1 - \epsilon.$$
- if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{ellipsoid-box}} \stackrel{\text{def}}{=} \{\mathbf{z} : \|\mathbf{z}\|_2 \leq \rho \stackrel{\text{def}}{=} \sqrt{2 \ln(1/\epsilon)}, \|\mathbf{z}\|_{\infty} \leq 1\}$,
$$\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] \geq 1 - \epsilon.$$
- if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{budget}} = \{\mathbf{z} \in \mathbb{R}^L : \|\mathbf{z}\|_{\infty} \leq 1, \|\mathbf{z}\|_1 \leq \Gamma = \sqrt{2 \ln(1/\epsilon)} \sqrt{L}\}$,
$$\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] \geq 1 - \epsilon.$$

How to Calibrate Uncertainty Sets?

We know how to derive RC for “structured” uncertainty sets. **But how to pick ρ, Γ ?**

Let's take a **probabilistic** view for a moment:

Suppose \mathbf{z}_i are really random, and we seek ρ, Γ to ensure $\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b]$ is “large”, $\forall \mathbb{P}$

Theorem (High Probability of Constraint Satisfaction)

Suppose \mathbf{z}_i are independent r.v. with mean $\mathbf{0}$ and support on $[-1, 1]$. Then:

- if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{box}} \stackrel{\text{def}}{=} \{\mathbf{z} : \|\mathbf{z}\|_{\infty} \leq 1\}$, $\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] = 1$.
(no need for independence or 0-mean)
 - if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{ellipsoid}} \stackrel{\text{def}}{=} \{\mathbf{z} : \|\mathbf{z}\|_2 \leq \rho \stackrel{\text{def}}{=} \sqrt{2 \ln(1/\epsilon)}\}$,
$$\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] \geq 1 - \epsilon.$$
 - if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{ellipsoid-box}} \stackrel{\text{def}}{=} \{\mathbf{z} : \|\mathbf{z}\|_2 \leq \rho \stackrel{\text{def}}{=} \sqrt{2 \ln(1/\epsilon)}, \|\mathbf{z}\|_{\infty} \leq 1\}$,
$$\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] \geq 1 - \epsilon.$$
 - if \mathbf{x} feasible for RC with $\mathcal{U}_{\text{budget}} = \{\mathbf{z} \in \mathbb{R}^L : \|\mathbf{z}\|_{\infty} \leq 1, \|\mathbf{z}\|_1 \leq \Gamma = \sqrt{2 \ln(1/\epsilon)} \sqrt{L}\}$,
$$\mathbb{P}[(\bar{\mathbf{a}} + \mathbf{P}\mathbf{z})^T \mathbf{x} \leq b] \geq 1 - \epsilon.$$
- Some probabilistic information allows controlling conservatism: **very useful in applications**
 - The budget Γ depends on the dimension of \mathbf{z} (L), whereas ρ does not!
 - Proof based on concentration inequalities

Another Quick Example: A Portfolio Problem (Ben-Tal and Nemirovski)

- 200 risky assets; asset # 200 is cash, with yearly return $r_{200} = 5\%$ and zero risk
- Yearly returns r_i are **independent r.v.** with values in $[\mu_i - \sigma_i, \mu_i + \sigma_i]$ and means μ_i :

$$\mu_i = 1.05 + 0.3 \frac{(200 - i)}{199}, \quad \sigma_i = 0.05 + 0.6 \frac{(200 - i)}{199}, \quad i = 1, \dots, 199.$$

- Goal: distribute \$1 so as to maximize worst-case value-at-risk at level $\epsilon = 0.5\%$:

$$\max_{x,t} \left\{ t : \mathbb{P} \left[\sum_{i=1}^{199} r_i x_i + r_{200} x_{200} \geq t \right] \geq 1 - \epsilon, \forall \mathbb{P}, \sum_{i=1}^{200} x_i = 1, x \geq 0 \right\},$$

Another Quick Example: A Portfolio Problem (Ben-Tal and Nemirovski)

- 200 risky assets; asset # 200 is cash, with yearly return $r_{200} = 5\%$ and zero risk
- Yearly returns r_i are **independent r.v.** with values in $[\mu_i - \sigma_i, \mu_i + \sigma_i]$ and means μ_i :

$$\mu_i = 1.05 + 0.3 \frac{(200 - i)}{199}, \quad \sigma_i = 0.05 + 0.6 \frac{(200 - i)}{199}, \quad i = 1, \dots, 199.$$

- Goal: distribute \$1 so as to maximize worst-case value-at-risk at level $\epsilon = 0.5\%$:

$$\max_{x, t} \left\{ t : \mathbb{P} \left[\sum_{i=1}^{199} r_i x_i + r_{200} x_{200} \geq t \right] \geq 1 - \epsilon, \forall \mathbb{P}, \sum_{i=1}^{200} x_i = 1, x \geq 0 \right\},$$

- With $z_i \stackrel{\text{def}}{=} (r_i - \mu_i)/\sigma_i$, let's consider 3 uncertainty sets:

① $\mathcal{U}_{\text{box}} = \{z : \|z\|_{\infty} \leq 1\}$

② $\mathcal{U}_{\text{ellipsoid-box}} = \{z : \|z\|_{\infty} \leq 1, \|z\|_2 \leq \rho\}$, with $\rho = \sqrt{2 \ln(1/\epsilon)} = 3.255$

③ $\mathcal{U}_{\text{budget}} = \{z : \|z\|_{\infty} \leq 1, \|z\|_1 \leq \Gamma\}$ with $\Gamma = \sqrt{2 \ln(1/\epsilon)} \sqrt{199} = 45.921$.

Another Quick Example: A Portfolio Problem (Ben-Tal and Nemirovski)

- 200 risky assets; asset # 200 is cash, with yearly return $r_{200} = 5\%$ and zero risk
- Yearly returns r_i are **independent r.v.** with values in $[\mu_i - \sigma_i, \mu_i + \sigma_i]$ and means μ_i :

$$\mu_i = 1.05 + 0.3 \frac{(200 - i)}{199}, \quad \sigma_i = 0.05 + 0.6 \frac{(200 - i)}{199}, \quad i = 1, \dots, 199.$$

- Goal: distribute \$1 so as to maximize worst-case value-at-risk at level $\epsilon = 0.5\%$:

$$\max_{x, t} \left\{ t : \mathbb{P} \left[\sum_{i=1}^{199} r_i x_i + r_{200} x_{200} \geq t \right] \geq 1 - \epsilon, \forall \mathbb{P}, \sum_{i=1}^{200} x_i = 1, x \geq 0 \right\},$$

- With $z_i \stackrel{\text{def}}{=} (r_i - \mu_i)/\sigma_i$, let's consider 3 uncertainty sets:

① $\mathcal{U}_{\text{box}} = \{z : \|z\|_{\infty} \leq 1\}$

② $\mathcal{U}_{\text{ellipsoid-box}} = \{z : \|z\|_{\infty} \leq 1, \|z\|_2 \leq \rho\}$, with $\rho = \sqrt{2 \ln(1/\epsilon)} = 3.255$

③ $\mathcal{U}_{\text{budget}} = \{z : \|z\|_{\infty} \leq 1, \|z\|_1 \leq \Gamma\}$ with $\Gamma = \sqrt{2 \ln(1/\epsilon)} \sqrt{199} = 45.921$.

- Results:

- \mathcal{U}_{box} : worst-case returns $r_i = \mu_i - \sigma_i$ yield less than risk-free return of 5%, so optimal to keep all money in cash; robust optimal return 1.05, risk 0
- $\mathcal{U}_{\text{ellipsoid-box}}$: robust optimal value is 1.12, risk 0.5%
- $\mathcal{U}_{\text{budget}}$: robust optimal value is 1.10, risk 0.5%

- \mathcal{U}_{box} can be quite conservative, a tiny bit of risk can go a long way...

Using Concentration Results to Model Uncertainty Sets

- Dimitris Bertsimas: let's use the **implications** of the **Central Limit Theorem**

*"All epistemological value of the theory of probability is based on this: that **large-scale random phenomena** in their collective action create strict, **non-random regularity**" (Gnedenko and Kolmogorov, 1954)*

Using Concentration Results to Model Uncertainty Sets

- Dimitris Bertsimas: let's use the **implications** of the **Central Limit Theorem**

"All epistemological value of the theory of probability is based on this: that large-scale random phenomena in their collective action create strict, non-random regularity" (Gnedenko and Kolmogorov, 1954)

- Suppose we have uncertainties $\{X_i\}_{i=1}^n$, each with mean μ , standard deviation σ

$$\mathcal{U}_{\text{CLT}} \stackrel{\text{def}}{=} \left\{ (x_1, \dots, x_n) : \left| \sum_{i=1}^n x_i - n\mu \right| \leq \Gamma \sigma \sqrt{n} \right\}.$$

- If n large and CLT premises hold, $\Gamma = 2$ (3) would give 95% (99%) coverage

Using Concentration Results to Model Uncertainty Sets

- Dimitris Bertsimas: let's use the **implications** of the **Central Limit Theorem**

*"All epistemological value of the theory of probability is based on this: that **large-scale random phenomena** in their collective action create strict, **non-random regularity**" (Gnedenko and Kolmogorov, 1954)*

- Suppose we have uncertainties $\{X_i\}_{i=1}^n$, each with mean μ , standard deviation σ

$$\mathcal{U}_{\text{CLT}} \stackrel{\text{def}}{=} \left\{ (x_1, \dots, x_n) : \left| \sum_{i=1}^n x_i - n\mu \right| \leq \Gamma \sigma \sqrt{n} \right\}.$$

- ▶ If n large and CLT premises hold, $\Gamma = 2$ (3) would give 95% (99%) coverage

- Many extensions possible

- ▶ Modeling correlations through a factor model:

$$\mathcal{U}_{\text{corr}} \stackrel{\text{def}}{=} \left\{ \mathbf{x} : \mathbf{x} = \mathbf{P}\mathbf{z} + \boldsymbol{\epsilon}, \left| \sum_{i=1}^m z_i - m\mu_y \right| \leq \Gamma \sigma_z \sqrt{m}, \left| \sum_{i=1}^n \epsilon_i \right| \leq \Gamma \sigma_\epsilon \sqrt{n}, \right\}.$$

- ▶ Using stable laws to model heavy-tailed cases where variance is undefined:

$$\mathcal{U}_{\text{HT}} \stackrel{\text{def}}{=} \left\{ (x_1, \dots, x_n) : \left| \sum_{i=1}^n x_i - n\mu \right| \leq \Gamma n^{1/\alpha} \right\}.$$

- ▶ Constructing typical sets: if H_f is the (Shannon) entropy of f ,

$$(i) \mathbb{P}[\tilde{\mathbf{z}} \in \mathcal{U}_{\text{typical}}] \rightarrow 1, \quad (ii) \left| \frac{1}{n} \log f(\tilde{\mathbf{z}} | \tilde{\mathbf{z}} \in \mathcal{U}_{\text{typical}}) + H_f \right| \leq \epsilon_n$$

Using Concentration Results to Model Uncertainty Sets

- Dimitris Bertsimas: let's use the **implications** of the **Central Limit Theorem**

*"All epistemological value of the theory of probability is based on this: that **large-scale random phenomena** in their collective action create strict, **non-random regularity**" (Gnedenko and Kolmogorov, 1954)*

- Suppose we have uncertainties $\{X_i\}_{i=1}^n$, each with mean μ , standard deviation σ

$$\mathcal{U}_{\text{CLT}} \stackrel{\text{def}}{=} \left\{ (x_1, \dots, x_n) : \left| \sum_{i=1}^n x_i - n\mu \right| \leq \Gamma \sigma \sqrt{n} \right\}.$$

- ▶ If n large and CLT premises hold, $\Gamma = 2$ (3) would give 95% (99%) coverage

- Many extensions possible

- ▶ Modeling correlations through a factor model:

$$\mathcal{U}_{\text{corr}} \stackrel{\text{def}}{=} \left\{ \mathbf{x} : \mathbf{x} = \mathbf{P}\mathbf{z} + \boldsymbol{\epsilon}, \left| \sum_{i=1}^m z_i - m\mu_y \right| \leq \Gamma \sigma_z \sqrt{m}, \left| \sum_{i=1}^n \epsilon_i \right| \leq \Gamma \sigma_\epsilon \sqrt{n}, \right\}.$$

- ▶ Using stable laws to model heavy-tailed cases where variance is undefined:

$$\mathcal{U}_{\text{HT}} \stackrel{\text{def}}{=} \left\{ (x_1, \dots, x_n) : \left| \sum_{i=1}^n x_i - n\mu \right| \leq \Gamma n^{1/\alpha} \right\}.$$

- ▶ Constructing typical sets: if H_f is the (Shannon) entropy of f ,

$$(i) \mathbb{P}[\tilde{\mathbf{z}} \in \mathcal{U}_{\text{typical}}] \rightarrow 1, \quad (ii) \left| \frac{1}{n} \log f(\tilde{\mathbf{z}} | \tilde{\mathbf{z}} \in \mathcal{U}_{\text{typical}}) + H_f \right| \leq \epsilon_n$$

- Bertsimas & Bandi used these to derive **robust equivalents** for several classical queueing theory and information theory results

Using Hypothesis Tests to Model Uncertainty Sets

Another powerful idea: derive **data-driven** uncertainty sets from **hypothesis tests**

From Bertsimas, Gupta, Kallus (2017):

Table 1 Summary of data-driven uncertainty sets proposed in this paper. SOC, EC and LMI denote second-order cone representable sets, exponential cone representable sets, and linear matrix inequalities, respectively

Assumptions on \mathbb{P}^*	Hypothesis test	Geometric description	Eqs.	Inner problem
Discrete support	χ^2 -test	SOC	(13, 15)	
Discrete support	G-test	Polyhedral*	(13, 16)	
Independent marginals	KS Test	Polyhedral*	(21)	Line search
Independent marginals	K Test	Polyhedral*	(76)	Line search
Independent marginals	CvM Test	SOC*	(76, 69)	
Independent marginals	W Test	SOC*	(76, 70)	
Independent marginals	AD Test	EC	(76, 71)	
Independent marginals	Chen et al. [23]	SOC	(27)	Closed-form
None	Marginal Samples	Box	(31)	Closed-form
None	Linear Convex Ordering	Polyhedron	(34)	
None	Shawe-Taylor and Cristianini [46]	SOC	(39)	Closed-form
None	Delage and Ye [25]	LMI	(41)	

The additional “*” notation indicates a set of the above type with one additional, relative entropy constraint. *KS*, *K*, *CvM*, *W*, and *AD* denote the Kolmogorov–Smirnov, Kuiper, Cramer-von Mises, Watson and Anderson-Darling goodness of fit tests, respectively. In some cases, we can identify a worst-case realization of \mathbf{u} in (1) for bi-affine f and a candidate \mathbf{x} with a specialized algorithm. In these cases, the column “Inner Problem” roughly describes this algorithm

Distributionally Robust Modeling

- We saw that embedding some more probabilistic information can help

Distributionally Robust Modeling

- We saw that embedding some more probabilistic information can help
- Let's change the paradigm slightly:
 - we know the probability distribution for $\tilde{\mathbf{z}}$ belongs to an *ambiguity set*: $\mathbb{P} \in \mathcal{P}$
 - we model \mathcal{P} , and are interested in robust expected constraint satisfaction:

$$\sup_{\mathbb{P} \in \mathcal{P}} \mathbb{E}_{\mathbb{P}}[f(\mathbf{x}, \tilde{\mathbf{z}})] \leq b$$

- Now, the adversary is choosing \mathbb{P} , instead of \mathbf{z}
 - **Key advantage:** $\mathbb{E}_{\mathbb{P}}[f(\mathbf{x}, \tilde{\mathbf{z}})]$ as an expression of \mathbb{P} is **always linear**, so much of our earlier machinery (e.g., convex duality) can be applied if the set \mathcal{P} is “well-behaved”

Distributionally Robust Modeling

- We saw that embedding some more probabilistic information can help
- Let's change the paradigm slightly:
 - we know the probability distribution for \tilde{z} belongs to an *ambiguity set*: $\mathbb{P} \in \mathcal{P}$
 - we model \mathcal{P} , and are interested in robust expected constraint satisfaction:

$$\sup_{\mathbb{P} \in \mathcal{P}} \mathbb{E}_{\mathbb{P}}[f(\mathbf{x}, \tilde{z})] \leq b$$

- Now, the adversary is choosing \mathbb{P} , instead of \mathbf{z}
 - **Key advantage**: $\mathbb{E}_{\mathbb{P}}[f(\mathbf{x}, \tilde{z})]$ as an expression of \mathbb{P} is **always linear**, so much of our earlier machinery (e.g., convex duality) can be applied if the set \mathcal{P} is “well-behaved”
- Very old idea, dating to the 1950s
 - Scarf (1958) studied a Newsvendor model with mean and variance of demand known
 - Zacks (1966) studied stochastic LPs with knowledge of mean and support
- Recent tutorial by Kuhn, Shafiee, Wiesemann (2024) very nice summary of state-of-the-art; can model:
 - known (**bounds on**) moments, e.g., means, covariance matrix, higher order
 - information about various spread statistics, e.g., absolute mean spread ($\mathbb{E}[X|X > \theta] - \mathbb{E}[X|X < \theta]$), mean absolute deviation ($\mathbb{E}[|X - m|]$), etc.
 - known (**bounds on**) quantiles, e.g., median
 - multiple confidence regions
 - distance from a nominal distribution (e.g., the empirical one)

Connections with Risk Measures

- A lot of effort in math finance since 2000 to characterize the “right” risk measures
- X is a random **loss**; a **risk measure** $\mu(X)$ should capture the “riskiness” in X
 - used to determine capital requirements for banks, so $\mu(X)$ should be “money”

Connections with Risk Measures

- A lot of effort in math finance since 2000 to characterize the “right” risk measures
- X is a random **loss**; a **risk measure** $\mu(X)$ should capture the “riskiness” in X
 - used to determine capital requirements for banks, so $\mu(X)$ should be “money”

Desirable Axioms (Artzner et al 1999)

- **[P1] Monotonicity:** If $X \leq Y$, then $\mu(X) \leq \mu(Y)$.
- **[P2] Influence of cash:** If $m \in \mathbb{R}$, then $\mu(X + m) = \mu(X) + m$.
- **[P3] Diversification:** $\mu(\lambda X + (1 - \lambda)Y) \leq \lambda \mu(X) + (1 - \lambda)\mu(Y)$, for $\lambda \in [0, 1]$.
- **[P4] Leverage:** If $\lambda \geq 0$, then $\mu(\lambda X) = \lambda \mu(X)$.

Connections with Risk Measures

- A lot of effort in math finance since 2000 to characterize the “right” risk measures
- X is a random **loss**; a **risk measure** $\mu(X)$ should capture the “riskiness” in X
 - used to determine capital requirements for banks, so $\mu(X)$ should be “money”

Desirable Axioms (Artzner et al 1999)

- **[P1] Monotonicity:** If $X \leq Y$, then $\mu(X) \leq \mu(Y)$.
 - **[P2] Influence of cash:** If $m \in \mathbb{R}$, then $\mu(X + m) = \mu(X) + m$.
 - **[P3] Diversification:** $\mu(\lambda X + (1 - \lambda)Y) \leq \lambda \mu(X) + (1 - \lambda)\mu(Y)$, for $\lambda \in [0, 1]$.
 - **[P4] Leverage:** If $\lambda \geq 0$, then $\mu(\lambda X) = \lambda \mu(X)$.
-
- E.g., variance does not satisfy **[P1]**, Value-at-Risk (VaR) does not satisfy **P3**
 - Conditional-Value-at-Risk (CVaR) satisfies [P1-P4]

Connections with Risk Measures

- A lot of effort in math finance since 2000 to characterize the “right” risk measures
- X is a random **loss**; a **risk measure** $\mu(X)$ should capture the “riskiness” in X
 - ▶ used to determine capital requirements for banks, so $\mu(X)$ should be “money”

Desirable Axioms (Artzner et al 1999)

- ▶ **[P1] Monotonicity:** If $X \leq Y$, then $\mu(X) \leq \mu(Y)$.
 - ▶ **[P2] Influence of cash:** If $m \in \mathbb{R}$, then $\mu(X + m) = \mu(X) + m$.
 - ▶ **[P3] Diversification:** $\mu(\lambda X + (1 - \lambda)Y) \leq \lambda \mu(X) + (1 - \lambda)\mu(Y)$, for $\lambda \in [0, 1]$.
 - ▶ **[P4] Leverage:** If $\lambda \geq 0$, then $\mu(\lambda X) = \lambda \mu(X)$.
- ▶ E.g., variance does not satisfy **[P1]**, Value-at-Risk (VaR) does not satisfy **P3**
 - ▶ Conditional-Value-at-Risk (CVaR) satisfies [P1-P4]

THM. μ satisfies **[P1-3]** iff there exists a convex function ϕ and

$$\mu(X) = \sup_{\mathbb{P}} [\mathbb{E}_{\mathbb{P}}[X] - \phi(\mathbb{P})]$$

THM. μ satisfies **[P1-4]** iff there exists a convex set of measures \mathcal{P} and

$$\mu(X) = \sup_{\mathbb{P} \in \mathcal{P}} \mathbb{E}_{\mathbb{P}}[X]$$

- Can construct uncertainty sets using risk measures (Bertsimas, Brown, Sim)
- Similar results also in decision theory under ambiguity (Gilboa, Schmeidler, etc.)

References I

- E. Adida and G. Perakis. A robust optimization approach to dynamic pricing and inventory control with no backorders. *Mathematical Programming*, 107: 97–129, 2006.
- O. Baron, J. Milner, and H. Naseraldin. Facility location: A robust optimization approach. *Production and Operations Management*, 20(5):772–785, 2011.
- A. Ben-Tal, A. Goryashko, E. Guslitzer, and A. Nemirovski. Adjustable robust solutions of uncertain linear programs. *Mathematical Programming*, 99(2): 351–376, 2004.
- A. Ben-Tal, B. Golany, A. Nemirovski, and J.-P. Vial. Retailer-supplier flexible commitments contracts: A robust optimization approach. *Manufacturing & Service Operations Management*, 7(3):248–271, 2005.
- A. Ben-Tal, L. El-Ghaoui, and A. Nemirovski. *Robust Optimization*. Princeton Series in Applied Mathematics. Princeton University Press, 2009.
- D. Bertsimas and D. Pachamanova. Robust multiperiod portfolio management in the presence of transaction costs. *Computers and Operations Research*, 35(1):3–17, 2008.
- D. Bertsimas and A. Thiele. A robust optimization approach to inventory theory. *Operations Research*, 54(1):150–168, 2006.
- D. Bienstock and N. Özbay. Computing robust basestock levels. *Discrete Optimization*, 5(2):389 – 414, 2008.
- S. Ceria and R. Stubbs. Incorporating estimation errors into portfolio selection: Robust portfolio construction. *Journal of Asset Management*, 7(2): 109–127, July 2006.
- F. de Ruiter, A. Ben-Tal, R. Brekelmans, and D. den Hertog. Adjustable robust optimization with decision rules based on inexact revealed data. Center discussion paper series no. 2014-003, CentER, 2014.
- D. Goldfarb and G. Iyengar. Robust Portfolio Selection Problems. *Mathematics of Operations Research*, 28(1):1–38, 2003.
- X. Lin, S. Janak, and C. Floudas. A new robust optimization approach for scheduling under uncertainty: I. bounded uncertainty. *Computers and Chemical Engineering*, 28:1069–1085, 2004.
- A. Lorca and A. Sun. Adaptive robust optimization with dynamic uncertainty sets for multi-period economic dispatch under significant wind. *IEEE Transactions on Power Systems*, 30(4):1702–1713, 2015.
- S. Mittal, A. Schulz, and S. Stiller. Robust appointment scheduling. In *APPROX*, 2014.
- G. Perakis and G. Roels. Robust controls for network revenue management. *Manufacturing & Service Operations Management*, 12(1):56–76, 2010.
- M. Pinar and R. Tütüncü. Robust profit opportunities in risky financial portfolios. *Operations Research Letters*, 33(4):331 – 340, 2005.
- A. Shapiro, D. Dentcheva, and A. Ruszczyński. *Lectures on Stochastic Programming*. MPS / SIAM Series on Optimization. SIAM, 2009.
- R. Tütüncü and M. Koenig. Robust asset allocation. *Annals of Operations Research*, 132(1-4):157–187, November 2004.
- W. Wiesemann, D. Kuhn, and B. Rustem. Robust resource allocations in temporal networks. *Mathematical Programming*, 135:437–471, 2012. ISSN 0025-5610.
- D. Yamashita, V. Armentano, and M. Laguna. Robust optimization models for project scheduling with resource availability cost. *Journal of Scheduling*, 10 (1):67–76, 2007.
- C. Zhao, J. Wang, J. Watson, and Y. Guan. Multi-stage robust unit commitment considering wind and demand response uncertainties. *IEEE Transactions on Power Systems*, 28(3):2708–2717, 2013.