

CME 307 / MS&E 311 / OIT 676: Optimization

LP geometry, modeling and solution techniques

Professor Udell

Management Science and Engineering
Stanford

September 29, 2026

First quiz Thursday next week

- ▶ Take paper when you arrive
- ▶ Write your SUNetID¹ and your name.
- ▶ We will project the question.
- ▶ You will have 5 minutes to write your answer on the paper.
- ▶ When time is up, put your pen or pencil down and stop writing.

¹Your SUNetID is your Stanford University Network ID, and consists of letters, **not** numbers. Your email is yourSUNetID@stanford.edu.

First quiz Thursday next week

- ▶ Take paper when you arrive
- ▶ Write your SUNetID¹ and your name.
- ▶ We will project the question.
- ▶ You will have 5 minutes to write your answer on the paper.
- ▶ When time is up, put your pen or pencil down and stop writing.

since a 5 minute quiz is very short, we will grade on a **check** basis.

- ▶ **no check**: no serious attempt
- ▶ **check**: a serious attempt
- ▶ **check+**: correct or nearly correct

¹Your SUNetID is your Stanford University Network ID, and consists of letters, **not** numbers. Your email is yourSUNetID@stanford.edu.

Exit ticket

Please fill out the exit ticket (linked from course schedule spreadsheet) before you leave:

- ▶ Helps us improve the course

We will not grade this, but we will read it!



Course overview

the practice of optimization consists of:

- ▶ modeling (real-world problems $\rightarrow$ optimization problems)
- ▶ analysis (properties of optimization problems)
- ▶ algorithms (how to solve optimization problems)

all three are necessary for success in optimization!

Course overview

the practice of optimization consists of:

- ▶ modeling (real-world problems $\rightarrow$ optimization problems)
- ▶ analysis (properties of optimization problems)
- ▶ algorithms (how to solve optimization problems)

all three are necessary for success in optimization!

- ▶ last class: mostly modeling
- ▶ today: mostly analysis
- ▶ next class: algorithms and some more modeling

Outline

LP standard form

LP inequality form

What kinds of points can be optimal?

Solving LPs

Modeling

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $\pm\infty$

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $\pm\infty$

Q: if $p^* = -\infty$, does a solution exist?

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $\pm\infty$

Q: if $p^* = -\infty$, does a solution exist?

Q: if $p^* = \infty$, does a solution exist?

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $\pm\infty$

Q: if $p^* = -\infty$, does a solution exist?

Q: if $p^* = \infty$, does a solution exist?

henceforth assume $A \in \mathbb{R}^{m \times n}$ has full row rank m

Q: why? how to check?

Linear programming: standard form

standard form linear program (LP)

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

optimal value p^* , solution x^* (if it exists)

- ▶ any x with $Ax = b$ and $x \geq 0$ is called a **feasible point**
- ▶ if problem is infeasible, we say $p^* = \infty$
- ▶ p^* can be finite or $\pm\infty$

Q: if $p^* = -\infty$, does a solution exist?

Q: if $p^* = \infty$, does a solution exist?

henceforth assume $A \in \mathbb{R}^{m \times n}$ has full row rank m

Q: why? how to check?

A: otherwise infeasible or redundant rows; use gaussian elimination to check and remove

LP example: diet problem

We are planning a backpacking trip, and want to minimize the total weight of the food packed subject to nutritional requirements. We have a list of essential nutrients and how much an active person needs of each. We also know the weight of each food, and how much of each nutrient is in each food.

- ▶ x_j servings of food j , $j = 1, \dots, n$
- ▶ c_j weight per serving
- ▶ a_{ij} amount of nutrient i in food j
- ▶ b_i required amount of nutrient i , $i = 1, \dots, m$

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}$$

Geometry of LP: conic hull

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

Geometry of LP: conic hull

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

interpretation: conic hull

- ▶ define the **cone** generated by $A = [A_1, \dots, A_n]$:

$$\{Ax \mid x \geq 0\} = \left\{ \sum_{i=1}^n A_i x_i \mid x \geq 0 \right\} = \mathbf{cone}(A_1, \dots, A_n)$$

Geometry of LP: conic hull

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

interpretation: conic hull

- ▶ define the **cone** generated by $A = [A_1, \dots, A_n]$:

$$\{Ax \mid x \geq 0\} = \left\{ \sum_{i=1}^n A_i x_i \mid x \geq 0 \right\} = \mathbf{cone}(A_1, \dots, A_n)$$

- ▶ LP is feasible if $b \in \mathbf{cone}(A_1, \dots, A_n)$

Geometry of LP: conic hull

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

interpretation: conic hull

- ▶ define the **cone** generated by $A = [A_1, \dots, A_n]$:

$$\{Ax \mid x \geq 0\} = \left\{ \sum_{i=1}^n A_i x_i \mid x \geq 0 \right\} = \mathbf{cone}(A_1, \dots, A_n)$$

- ▶ LP is feasible if $b \in \mathbf{cone}(A_1, \dots, A_n)$

Q: how many columns are needed to represent a feasible $b \in \mathbb{R}^m$?

Geometry of LP: conic hull

the **feasible set** is the set of points $\{x \mid Ax = b, x \geq 0\}$ that satisfy all constraints.

interpretation: conic hull


- define the **cone** generated by $A = [A_1, \dots, A_n]$:

$$\{Ax \mid x \geq 0\} = \left\{ \sum_{i=1}^n A_i x_i \mid x \geq 0 \right\} = \mathbf{cone}(A_1, \dots, A_n)$$

- LP is feasible if $b \in \mathbf{cone}(A_1, \dots, A_n)$

Q: how many columns are needed to represent a feasible $b \in \mathbb{R}^m$?

A: at most m (here, b uses only a_1 and a_4)



Geometry of LP: hyperplane and halfspaces

**interpretation: intersection of
hyperplane and halfspaces**

- ▶ define a **hyperplane** $\{x \mid Ax = b\}$

Geometry of LP: hyperplane and halfspaces

**interpretation: intersection of
hyperplane and halfspaces**

- ▶ define a **hyperplane** $\{x \mid Ax = b\}$
(dimension?)

Geometry of LP: hyperplane and halfspaces

**interpretation: intersection of
hyperplane and halfspaces**

- ▶ define a **hyperplane** $\{x \mid Ax = b\}$
(dimension?)
- ▶ define a **halfspace** $\{x \mid a^T x \geq b\}$

Geometry of LP: hyperplane and halfspaces

**interpretation: intersection of
hyperplane and halfspaces**


- ▶ define a **hyperplane** $\{x \mid Ax = b\}$
(dimension?)
- ▶ define a **halfspace** $\{x \mid a^T x \geq b\}$
- ▶ the **positive orthant** $x \geq 0$ is an
intersection of halfspaces

Geometry of LP: hyperplane and halfspaces

interpretation: intersection of hyperplane and halfspaces

- ▶ define a **hyperplane** $\{x \mid Ax = b\}$ (dimension?)
- ▶ define a **halfspace** $\{x \mid a^T x \geq b\}$
- ▶ the **positive orthant** $x \geq 0$ is an intersection of halfspaces
- ▶ LP is feasible if hyperplane $\{x \mid Ax = b\}$ intersects the positive orthant

example: in $\mathbb{R}^3$, $\mathbf{1}^T x = 1$ cuts the orthant in the triangle $\text{conv}\{e_1, e_2, e_3\}$



Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ define the **convex hull** of a set S :

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1 \right\}$$

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ define the **convex hull** of a set S :

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1 \right\}$$

- ▶ define **polytope**: the convex hull of a finite set: $\text{conv}(\{x_1, \dots, x_k\})$

some useful convex sets:

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ define the **convex hull** of a set S :

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1 \right\}$$

- ▶ define **polytope**: the convex hull of a finite set: $\text{conv}(\{x_1, \dots, x_k\})$

some useful convex sets:

- ▶ a hyperplane is convex

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ define the **convex hull** of a set S :

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1 \right\}$$

- ▶ define **polytope**: the convex hull of a finite set: $\text{conv}(\{x_1, \dots, x_k\})$

some useful convex sets:

- ▶ a hyperplane is convex
- ▶ a halfspace is convex

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ define the **convex hull** of a set S :

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1 \right\}$$

- ▶ define **polytope**: the convex hull of a finite set: $\text{conv}(\{x_1, \dots, x_k\})$

some useful convex sets:

- ▶ a hyperplane is convex
- ▶ a halfspace is convex
- ▶ the intersection of convex sets is convex

Geometry of LP: convexity

- ▶ define **convex combination** of $x, y \in \mathbb{R}^n$: $\theta x + (1 - \theta)y$ for $\theta \in [0, 1]$
- ▶ define **convex set**: C is convex if for any $x, y \in C$,

$$\theta x + (1 - \theta)y \in C, \quad \theta \in [0, 1]$$

- ▶ define the **convex hull** of a set S :

$$\text{conv}(S) = \left\{ \sum_{i=1}^k \theta_i x_i \mid x_i \in S, \theta_i \geq 0, \sum_{i=1}^k \theta_i = 1 \right\}$$

- ▶ define **polytope**: the convex hull of a finite set: $\text{conv}(\{x_1, \dots, x_k\})$

some useful convex sets:

- ▶ a hyperplane is convex
- ▶ a halfspace is convex
- ▶ the intersection of convex sets is convex
- ▶ the feasible set $\{x : Ax = b, x \geq 0\}$ is convex

Outline

LP standard form

LP inequality form

What kinds of points can be optimal?

Solving LPs

Modeling

LP inequality form

another useful form for LP is **inequality form**

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

LP inequality form

another useful form for LP is **inequality form**

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b\end{array}$$

interpretation: halfspaces

- ▶ $a_i^T x \leq b_i$ defines a **halfspace**
- ▶ $Ax \leq b$ defines a **polyhedron**: intersection of halfspaces
- ▶ LP is feasible if polyhedron $\{x \mid Ax \leq b\}$ is nonempty

LP example: production planning

- ▶ x_i units of product i
- ▶ c_i cost per unit
- ▶ a_{ij} amount of resource j used by product i
- ▶ b_j amount of resource j available
- ▶ d_i demand for product i

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & 0 \leq x \leq d\end{array}$$

LP example: production planning

- ▶ x_i units of product i
- ▶ c_i cost per unit
- ▶ a_{ij} amount of resource j used by product i
- ▶ b_j amount of resource j available
- ▶ d_i demand for product i

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & 0 \leq x \leq d\end{array}$$

extensions:

- ▶ fixed cost for producing product i at all?

LP example: production planning

- ▶ x_i units of product i
- ▶ c_i cost per unit
- ▶ a_{ij} amount of resource j used by product i
- ▶ b_j amount of resource j available
- ▶ d_i demand for product i

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & 0 \leq x \leq d\end{array}$$

extensions:

- ▶ fixed cost for producing product i at all?
 $c^T x + f^T z$, $z_i \in \{0, 1\}$, $x_i \leq Mz_i$ for M large

LP inequality form to standard form

standard form to inequality form

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad \rightarrow$$

LP inequality form to standard form

standard form to inequality form

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad \rightarrow \quad \begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & Ax \geq b \\ & -x \leq 0 \end{array}$$

LP inequality form to standard form

standard form to inequality form

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad \rightarrow \quad \begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & Ax \geq b \\ & -x \leq 0 \end{array}$$

inequality form to standard form

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \end{array} \quad \rightarrow$$

LP inequality form to standard form

standard form to inequality form

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad \rightarrow \quad \begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \\ & Ax \geq b \\ & -x \leq 0 \end{array}$$

inequality form to standard form

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax \leq b \end{array} \quad \rightarrow \quad \begin{array}{ll} \text{minimize} & c^T (x_+ - x_-) \\ \text{subject to} & A(x_+ - x_-) + s = b \\ & s, x_+, x_- \geq 0 \end{array}$$

so both forms have the same expressive power, and feasible sets are polyhedra

Active constraints and variables

active constraints. for constraint set $Ax \leq b$, an **active constraint** i at x is one that holds with equality:

$$a_i^T x = b_i$$

Active constraints and variables

active constraints. for constraint set $Ax \leq b$, an **active constraint** i at x is one that holds with equality:

$$a_i^T x = b_i$$

- ▶ the **active set** $S(x)$ at x is the set of indices of active constraints

$$S(x) = \{i \mid a_i^T x = b_i\}$$

Active constraints and variables

active constraints. for constraint set $Ax \leq b$, an **active constraint** i at x is one that holds with equality:

$$a_i^T x = b_i$$

- ▶ the **active set** $S(x)$ at x is the set of indices of active constraints

$$S(x) = \{i \mid a_i^T x = b_i\}$$

active variables. for nonnegative variable $x \geq 0$, variable i is **active** if $x_i > 0$

example: active slack variables are dual to active constraints

$$Ax \leq b \iff Ax + s = b, s \geq 0$$


$$a_i^T x = b_i \iff s_i = 0$$

constraint i is active $\iff$ slack variable s_i is inactive

Active constraints: example

$$F = \left\{ x \in \mathbb{R}^2 \mid \begin{array}{l} x \geq 0, \ x_1 \leq 4, \\ x_2 \leq 3, \ x_1 + x_2 \leq 5 \end{array} \right\}$$

- ▶ corner $v = (4, 1)$: $x_1 \leq 4$ and $x_1 + x_2 \leq 5$ active
- ▶ edge point $e = (3, 2)$: only $x_1 + x_2 \leq 5$ active
- ▶ interior point $p = (2, 1)$: none active



Outline

LP standard form

LP inequality form

What kinds of points can be optimal?

Solving LPs

Modeling

Extreme points

define **extreme point**: $x \in \mathbb{R}^n$ is extreme in $C \subset \mathbb{R}^n$ if it cannot be written as a convex combination of other points in C : for $\theta \in [0, 1]$,

$$x \in C \quad \text{and} \quad x = \theta y + (1 - \theta)z \quad \implies \quad x = y = z$$

Extreme points

define **extreme point**: $x \in \mathbb{R}^n$ is extreme in $C \subset \mathbb{R}^n$ if it cannot be written as a convex combination of other points in C : for $\theta \in [0, 1]$,

$$x \in C \quad \text{and} \quad x = \theta y + (1 - \theta)z \quad \implies \quad x = y = z$$

fact: if x^* is the unique optimal solution of $\text{minimize}_{x \in S} c^T x$, then x^* is extreme in the set S .

Extreme points

define **extreme point**: $x \in \mathbb{R}^n$ is extreme in $C \subset \mathbb{R}^n$ if it cannot be written as a convex combination of other points in C : for $\theta \in [0, 1]$,

$$x \in C \quad \text{and} \quad x = \theta y + (1 - \theta)z \quad \implies \quad x = y = z$$

fact: if x^* is the unique optimal solution of $\text{minimize}_{x \in S} c^T x$, then x^* is extreme in the set S .

proof: suppose by way of contradiction that x^* is not extreme in S :

$$\begin{aligned} x^* &= \theta y + (1 - \theta)z \quad \text{for } y, z \in S, \theta \in (0, 1) \\ p^* := c^T x^* &= \theta c^T y + (1 - \theta)c^T z > \theta p^* + (1 - \theta)p^* = p^* \end{aligned}$$

where the inequality follows from the (unique) optimality of x^* . Contradiction!

Extreme points

define **extreme point**: $x \in \mathbb{R}^n$ is extreme in $C \subset \mathbb{R}^n$ if it cannot be written as a convex combination of other points in C : for $\theta \in [0, 1]$,

$$x \in C \quad \text{and} \quad x = \theta y + (1 - \theta)z \quad \implies \quad x = y = z$$

fact: if x^* is the unique optimal solution of $\text{minimize}_{x \in S} c^T x$,
then x^* is extreme in the set S .

proof: suppose by way of contradiction that x^* is not extreme in S :

$$\begin{aligned} x^* &= \theta y + (1 - \theta)z && \text{for } y, z \in S, \theta \in (0, 1) \\ p^* := c^T x^* &= \theta c^T y + (1 - \theta)c^T z > \theta p^* + (1 - \theta)p^* = p^* \end{aligned}$$

where the inequality follows from the (unique) optimality of x^* . Contradiction!

Q: Example of an LP with a non-extreme solution?

Extreme points

define **extreme point**: $x \in \mathbb{R}^n$ is extreme in $C \subset \mathbb{R}^n$ if it cannot be written as a convex combination of other points in C : for $\theta \in [0, 1]$,

$$x \in C \quad \text{and} \quad x = \theta y + (1 - \theta)z \quad \implies \quad x = y = z$$

fact: if x^* is the unique optimal solution of $\text{minimize}_{x \in S} c^T x$, then x^* is extreme in the set S .

proof: suppose by way of contradiction that x^* is not extreme in S :

$$\begin{aligned} x^* &= \theta y + (1 - \theta)z && \text{for } y, z \in S, \theta \in (0, 1) \\ p^* := c^T x^* &= \theta c^T y + (1 - \theta)c^T z > \theta p^* + (1 - \theta)p^* = p^* \end{aligned}$$

where the inequality follows from the (unique) optimality of x^* . Contradiction!

Q: Example of an LP with a non-extreme solution?

Q: Does there always exist an extreme solution?

Vertices

define **vertex**: $x \in S \subset \mathbb{R}^n$ is a vertex of set S if for some vector $c \in \mathbb{R}^n$,

$$c^T x < c^T y \quad \forall y \in S \setminus \{x\}$$

Vertices

define **vertex**: $x \in S \subset \mathbb{R}^n$ is a vertex of set S if for some vector $c \in \mathbb{R}^n$,

$$c^T x < c^T y \quad \forall y \in S \setminus \{x\}$$

interpretation: $\{z : c^T z = c^T x\}$ is a hyperplane that intersects S only at x .
we say this hyperplane **supports** S at x

Vertices

define **vertex**: $x \in S \subset \mathbb{R}^n$ is a vertex of set S if for some vector $c \in \mathbb{R}^n$,

$$c^T x < c^T y \quad \forall y \in S \setminus \{x\}$$

interpretation: $\{z : c^T z = c^T x\}$ is a hyperplane that intersects S only at x .
we say this hyperplane **supports** S at x

fact: x is a vertex of $S \implies x$ is an extreme point of S

Vertices

define **vertex**: $x \in S \subset \mathbb{R}^n$ is a vertex of set S if for some vector $c \in \mathbb{R}^n$,

$$c^T x < c^T y \quad \forall y \in S \setminus \{x\}$$

interpretation: $\{z : c^T z = c^T x\}$ is a hyperplane that intersects S only at x .
we say this hyperplane **supports** S at x

fact: x is a vertex of $S \implies x$ is an extreme point of S

proof:

Vertices

define **vertex**: $x \in S \subset \mathbb{R}^n$ is a vertex of set S if for some vector $c \in \mathbb{R}^n$,

$$c^T x < c^T y \quad \forall y \in S \setminus \{x\}$$

interpretation: $\{z : c^T z = c^T x\}$ is a hyperplane that intersects S only at x .
we say this hyperplane **supports** S at x

fact: x is a vertex of $S \implies x$ is an extreme point of S

proof: x is a vertex of S . suppose its defining vector is c and consider the optimization problem

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & x \in S \end{array}$$

Vertices

define **vertex**: $x \in S \subset \mathbb{R}^n$ is a vertex of set S if for some vector $c \in \mathbb{R}^n$,

$$c^T x < c^T y \quad \forall y \in S \setminus \{x\}$$

interpretation: $\{z : c^T z = c^T x\}$ is a hyperplane that intersects S only at x .
we say this hyperplane **supports** S at x

fact: x is a vertex of $S \implies x$ is an extreme point of S


proof: x is a vertex of S . suppose its defining vector is c and consider the optimization problem

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & x \in S \end{array}$$

x is the unique optimum of this problem, so the proof of this statement follows from the previous proof.

Extreme points and supporting hyperplanes


$$P = \text{conv}\{(0, 0), (3, 0), (2, 2), (0, 2)\}$$



Extreme points and supporting hyperplanes

$$P = \text{conv}\{(0, 0), (3, 0), (2, 2), (0, 2)\}$$

Q: is y an extreme point? is x^* ?




Extreme points and supporting hyperplanes

$$P = \text{conv}\{(0, 0), (3, 0), (2, 2), (0, 2)\}$$

Q: is y an extreme point? is x^* ?

A: y : no, $y = \frac{1}{2}(0, 2) + \frac{1}{2}(2, 2)$;

x^* : yes



Extreme points and supporting hyperplanes


$$P = \text{conv}\{(0, 0), (3, 0), (2, 2), (0, 2)\}$$

Q: is y an extreme point? is x^* ?

A: y : no, $y = \frac{1}{2}(0, 2) + \frac{1}{2}(2, 2)$;

x^* : yes

Q: is y a vertex? is x^* ?



Extreme points and supporting hyperplanes

$$P = \text{conv}\{(0, 0), (3, 0), (2, 2), (0, 2)\}$$


Q: is y an extreme point? is x^* ?

A: y : no, $y = \frac{1}{2}(0, 2) + \frac{1}{2}(2, 2)$;

x^* : yes

Q: is y a vertex? is x^* ?

A: y : no (not extreme); x^* : yes, it uniquely minimizes $c^T x$ for $c = (-1, -1)$



Extreme points and supporting hyperplanes

$$P = \text{conv}\{(0, 0), (3, 0), (2, 2), (0, 2)\}$$

Q: is y an extreme point? is x^* ?


A: y : no, $y = \frac{1}{2}(0, 2) + \frac{1}{2}(2, 2)$;

x^* : yes

Q: is y a vertex? is x^* ?

A: y : no (not extreme); x^* : yes, it uniquely minimizes $c^T x$ for $c = (-1, -1)$

Q: which other c make x^* the unique minimizer?



Extreme points and supporting hyperplanes

$$P = \text{conv}\{(0, 0), (3, 0), (2, 2), (0, 2)\}$$

Q: is y an extreme point? is x^* ?

A: y : no, $y = \frac{1}{2}(0, 2) + \frac{1}{2}(2, 2)$;


x^* : yes

Q: is y a vertex? is x^* ?

A: y : no (not extreme); x^* : yes, it uniquely minimizes $c^T x$ for $c = (-1, -1)$

Q: which other c make x^* the unique minimizer?

A: any $-c$ strictly between the outward normals $(0, 1)$ and $(2, 1)$ of the two edges at x^*



Basic feasible solution

recall the standard form LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad (\text{LP})$$

Basic feasible solution

recall the standard form LP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}\quad (\text{LP})$$

define: $x \in \mathbb{R}^n$ is a **basic feasible solution** (BFS) of (LP) if there is a set $S \subset \{1, \dots, n\}$ of m columns so that $A_S \in \mathbb{R}^{m \times m}$ is invertible and

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0, \quad x \geq 0.$$

- ▶ $A_S = \{A_{S_1}, \dots, A_{S_m}\} \in \mathbb{R}^{m \times m}$ is submatrix of A with columns in S
- ▶ $\bar{S}$ is the complement of S in $\{1, \dots, n\}$

Basic feasible solution

recall the standard form LP

$$\begin{array}{ll}\text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0\end{array}\quad (\text{LP})$$

define: $x \in \mathbb{R}^n$ is a **basic feasible solution** (BFS) of (LP) if there is a set $S \subset \{1, \dots, n\}$ of m columns so that $A_S \in \mathbb{R}^{m \times m}$ is invertible and

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0, \quad x \geq 0.$$

- ▶ $A_S = \{A_{S_1}, \dots, A_{S_m}\} \in \mathbb{R}^{m \times m}$ is submatrix of A with columns in S
- ▶ $\bar{S}$ is the complement of S in $\{1, \dots, n\}$
- ▶ two BFS with S, S' are neighbors if they share all but one columns:
 $|S \cap S'| = m - 1$

Basic feasible solution

recall the standard form LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad (\text{LP})$$

define: $x \in \mathbb{R}^n$ is a **basic feasible solution** (BFS) of (LP) if there is a set $S \subset \{1, \dots, n\}$ of m columns so that $A_S \in \mathbb{R}^{m \times m}$ is invertible and

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0, \quad x \geq 0.$$

- ▶ $A_S = \{A_{S_1}, \dots, A_{S_m}\} \in \mathbb{R}^{m \times m}$ is submatrix of A with columns in S
- ▶ $\bar{S}$ is the complement of S in $\{1, \dots, n\}$
- ▶ two BFS with S, S' are neighbors if they share all but one columns:
 $|S \cap S'| = m - 1$

Basic feasible solution

recall the standard form LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad (\text{LP})$$

define: $x \in \mathbb{R}^n$ is a **basic feasible solution** (BFS) of (LP) if there is a set $S \subset \{1, \dots, n\}$ of m columns so that $A_S \in \mathbb{R}^{m \times m}$ is invertible and

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0, \quad x \geq 0.$$

- ▶ $A_S = \{A_{S_1}, \dots, A_{S_m}\} \in \mathbb{R}^{m \times m}$ is submatrix of A with columns in S
- ▶ $\bar{S}$ is the complement of S in $\{1, \dots, n\}$
- ▶ two BFS with S, S' are neighbors if they share all but one columns:
 $|S \cap S'| = m - 1$

Q: how to find a BFS?

Basic feasible solution

recall the standard form LP

$$\begin{array}{ll} \text{minimize} & c^T x \\ \text{subject to} & Ax = b \\ & x \geq 0 \end{array} \quad (\text{LP})$$

define: $x \in \mathbb{R}^n$ is a **basic feasible solution** (BFS) of (LP) if there is a set $S \subset \{1, \dots, n\}$ of m columns so that $A_S \in \mathbb{R}^{m \times m}$ is invertible and

$$x_S = A_S^{-1}b, \quad x_{\bar{S}} = 0, \quad x \geq 0.$$



- ▶ $A_S = \{A_{S_1}, \dots, A_{S_m}\} \in \mathbb{R}^{m \times m}$ is submatrix of A with columns in S
- ▶ $\bar{S}$ is the complement of S in $\{1, \dots, n\}$
- ▶ two BFS with S, S' are neighbors if they share all but one columns:
 $|S \cap S'| = m - 1$

Q: how to find a BFS?

A: choose m linearly independent columns of A and set $x = A_S^{-1}b$; check $x \geq 0$.



A basis produces a basic feasible solution

$$A = \begin{bmatrix} 1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 \end{bmatrix}, \quad b = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$



A basis produces a basic feasible solution



$$A = \begin{bmatrix} 1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 \end{bmatrix}, \quad b = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$



Q: choose $S = \{1, 2\}$. what is the BFS?

A basis produces a basic feasible solution

$$A = \begin{bmatrix} 1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 \end{bmatrix}, \quad b = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$





Q: choose $S = \{1, 2\}$. what is the BFS?

A: $x_S = A_S^{-1}b = (3, 2)$, so $x = (3, 2, 0, 0)$: the corner $x_3 = x_4 = 0$

A basis produces a basic feasible solution

$$A = \begin{bmatrix} 1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 \end{bmatrix}, \quad b = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$





Q: choose $S = \{1, 2\}$. what is the BFS?

A: $x_S = A_S^{-1}b = (3, 2)$, so $x = (3, 2, 0, 0)$: the corner $x_3 = x_4 = 0$

Q: is $S = \{1, 4\}$ a basis? does it give a BFS?

A basis produces a basic feasible solution

$$A = \begin{bmatrix} 1 & 0 & 1 & 2 \\ 0 & 1 & 1 & 1 \end{bmatrix}, \quad b = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$



Q: choose $S = \{1, 2\}$. what is the BFS?

A: $x_S = A_S^{-1}b = (3, 2)$, so $x = (3, 2, 0, 0)$: the corner $x_3 = x_4 = 0$

Q: is $S = \{1, 4\}$ a basis? does it give a BFS?

A: a basis, but $x = (-1, 0, 0, 2) \not\geq 0$: basic but **infeasible**

Extreme point $\iff$ vertex $\iff$ BFS

fact. consider the feasible set in standard form $F = \{x \mid Ax = b, x \geq 0\}$ in $\mathbb{R}^n$.
the following are equivalent:

- ▶ x is an extreme point of F
- ▶ x is a vertex of F
- ▶ x is a BFS of F

Extreme point $\iff$ vertex $\iff$ BFS

fact. consider the feasible set in standard form $F = \{x \mid Ax = b, x \geq 0\}$ in $\mathbb{R}^n$.
the following are equivalent:

- ▶ x is an extreme point of F
- ▶ x is a vertex of F
- ▶ x is a BFS of F

implications: since any polyhedron $Ax \leq b$ can be rewritten in standard form,

- ▶ (BFS $\implies$) a polyhedron has a finite number of extreme points
- ▶ (extreme point $\implies$) BFS are independent of the representation of the feasible set

Extreme point $\iff$ vertex $\iff$ BFS

fact. consider the feasible set in standard form $F = \{x \mid Ax = b, x \geq 0\}$ in $\mathbb{R}^n$.
the following are equivalent:

- ▶ x is an extreme point of F
- ▶ x is a vertex of F
- ▶ x is a BFS of F

implications: since any polyhedron $Ax \leq b$ can be rewritten in standard form,

- ▶ (BFS $\implies$) a polyhedron has a finite number of extreme points
- ▶ (extreme point $\implies$) BFS are independent of the representation of the feasible set

we have already shown that vertex $\implies$ extreme point. need to show

- ▶ extreme point $\implies$ BFS
- ▶ BFS $\implies$ vertex

Extreme point $\implies$ BFS

let $x^\star$ be an extreme point of F , and let

$$S = \{i : x_i^\star > 0\}.$$

Extreme point $\implies$ BFS

let x^* be an extreme point of F , and let

$$S = \{i : x_i^* > 0\}.$$

1. The columns in A_S are linearly independent. Otherwise, there is a nonzero $d \in \mathbb{R}^n$, supported on S , with $Ad = 0$. Since $x_i^* > 0$ on S , for sufficiently small $\epsilon > 0$,

$$x^+ = x^* + \epsilon d, \quad x^- = x^* - \epsilon d$$

are nonnegative. They also satisfy $Ax^+ = Ax^- = b$. Hence $x^+, x^- \in F$, but

$$x^* = \frac{1}{2}x^+ + \frac{1}{2}x^-, \quad x^+ \neq x^-,$$

contradicting that x^* is extreme.

Extreme point $\implies$ BFS

let x^* be an extreme point of F , and let

$$S = \{i : x_i^* > 0\}.$$

1. The columns in A_S are linearly independent. Otherwise, there is a nonzero $d \in \mathbb{R}^n$, supported on S , with $Ad = 0$. Since $x_i^* > 0$ on S , for sufficiently small $\epsilon > 0$,

$$x^+ = x^* + \epsilon d, \quad x^- = x^* - \epsilon d$$

are nonnegative. They also satisfy $Ax^+ = Ax^- = b$. Hence $x^+, x^- \in F$, but

$$x^* = \frac{1}{2}x^+ + \frac{1}{2}x^-, \quad x^+ \neq x^-,$$

contradicting that x^* is extreme.

2. Complete A_S to a basis. Since A has rank m , extend S to B with $|B| = m$ and A_B invertible. Then $x_B^* = 0$, and $Ax^* = b$ gives

$$x_B^* = A_B^{-1}b.$$

Thus x^* is a BFS.

BFS $\implies$ vertex

suppose x^* is a BFS of F with active set S and A_S invertible. define $c \in \mathbb{R}^n$ as

$$c_i = \begin{cases} 0 & \text{if } i \in S \\ 1 & \text{otherwise} \end{cases}$$

so $c^T x^* = 0$.

BFS $\implies$ vertex

suppose x^* is a BFS of F with active set S and A_S invertible. define $c \in \mathbb{R}^n$ as

$$c_i = \begin{cases} 0 & \text{if } i \in S \\ 1 & \text{otherwise} \end{cases}$$

so $c^T x^* = 0$.

- ▶ x^* is the only point in F supported on S , as $\text{nullspace}(A_S) = 0$,
- ▶ so any other feasible point $x \in F$ has a positive objective value $c^T x > 0$.

hence x^* is a vertex of F with defining vector c .

Outline

LP standard form

LP inequality form

What kinds of points can be optimal?

Solving LPs

Modeling

Solving LPs

algorithms:

- ▶ enumerate all vertices and check
- ▶ fourier-motzkin elimination
- ▶ simplex method
- ▶ ellipsoid method
- ▶ interior point methods
- ▶ first-order methods
- ▶ ...

Solving LPs

algorithms:

- ▶ enumerate all vertices and check
- ▶ fourier-motzkin elimination
- ▶ simplex method
- ▶ ellipsoid method
- ▶ interior point methods
- ▶ first-order methods
- ▶ ...

remarks:

- ▶ enumeration and elimination are simple but not practical
- ▶ simplex was the first practical algorithm; still used today
- ▶ ellipsoid method is the first polynomial-time algorithm; not practical
- ▶ interior point methods are polynomial-time and practical
- ▶ first-order methods are practical and scale to large problems

Example of Fourier-Motzkin elimination

consider the system of inequalities

$$x_1 + 2x_2 \leq 4$$

$$-x_1 + x_2 \leq 1$$

$$x_1, x_2 \geq 0$$

Example of Fourier-Motzkin elimination

consider the system of inequalities

$$x_1 + 2x_2 \leq 4$$

$$-x_1 + x_2 \leq 1$$

$$x_1, x_2 \geq 0$$

we can collect inequalities on x_1 into those bounding it above and below:

$$\{0, x_2 - 1\} \leq x_1 \leq 4 - 2x_2$$

Example of Fourier-Motzkin elimination

consider the system of inequalities

$$x_1 + 2x_2 \leq 4$$

$$-x_1 + x_2 \leq 1$$

$$x_1, x_2 \geq 0$$

we can collect inequalities on x_1 into those bounding it above and below:

$$\{0, x_2 - 1\} \leq x_1 \leq 4 - 2x_2$$

by appending all pairwise inequalities to existing inequalities on x_2 , we recover the feasible set for x_2 :

$$0 \leq 4 - 2x_2, \quad x_2 - 1 \leq 4 - 2x_2, \quad x_2 \geq 0$$

$$\implies x_2 \in [0, 5/3].$$

Example of Fourier-Motzkin elimination

consider the system of inequalities

$$x_1 + 2x_2 \leq 4$$

$$-x_1 + x_2 \leq 1$$

$$x_1, x_2 \geq 0$$

we can collect inequalities on x_1 into those bounding it above and below:


$$\{0, x_2 - 1\} \leq x_1 \leq 4 - 2x_2$$

by appending all pairwise inequalities to existing inequalities on x_2 , we recover the feasible set for x_2 :

$$0 \leq 4 - 2x_2, \quad x_2 - 1 \leq 4 - 2x_2, \quad x_2 \geq 0$$

$$\implies x_2 \in [0, 5/3].$$

elimination method also shows projection of a polyhedron is a (closed) polyhedron



Enumerate vertices of LP

can generate all extreme points of LP: for each $S \subseteq \{1, \dots, n\}$ with $|S| = m$,

- ▶ $A_S \in \mathbb{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ solve $A_S x_S = b$ for x_S and set $x_{\bar{S}} = 0$
- ▶ if $x_S \geq 0$, then x is a BFS
- ▶ evaluate objective $c^T x$

the best BFS is optimal!

Enumerate vertices of LP

can generate all extreme points of LP: for each $S \subseteq \{1, \dots, n\}$ with $|S| = m$,

- ▶ $A_S \in \mathbb{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ solve $A_S x_S = b$ for x_S and set $x_{\bar{S}} = 0$
- ▶ if $x_S \geq 0$, then x is a BFS
- ▶ evaluate objective $c^T x$

the best BFS is optimal!

problem: how many BFSs are there?

Enumerate vertices of LP

can generate all extreme points of LP: for each $S \subseteq \{1, \dots, n\}$ with $|S| = m$,

- ▶ $A_S \in \mathbb{R}^{m \times m}$, submatrix of A with columns in S , is invertible
- ▶ solve $A_S x_S = b$ for x_S and set $x_{\bar{S}} = 0$
- ▶ if $x_S \geq 0$, then x is a BFS
- ▶ evaluate objective $c^T x$

the best BFS is optimal!

problem: how many BFSs are there?

n choose m is $\binom{n}{m} = \frac{n!}{m!(n-m)!}$ (“exponentially many”)


Simplex algorithm: example

$$\begin{array}{ll}\text{minimize} & -x_1 - 2x_2 \\ \text{subject to} & x_1 + x_2 \leq 4 \\ & x_1 + 3x_2 \leq 6 \\ & x \geq 0\end{array}$$

Simplex algorithm: example

$$\begin{array}{ll}\text{minimize} & -x_1 - 2x_2 \\ \text{subject to} & x_1 + x_2 \leq 4 \\ & x_1 + 3x_2 \leq 6 \\ & x \geq 0\end{array}$$


- start at $x^{(0)} = (0, 0)$: objective 0



Simplex algorithm: example

$$\begin{array}{ll}\text{minimize} & -x_1 - 2x_2 \\ \text{subject to} & x_1 + x_2 \leq 4 \\ & x_1 + 3x_2 \leq 6 \\ & x \geq 0\end{array}$$


- ▶ start at $x^{(0)} = (0, 0)$: objective 0
- ▶ increase x_2 until a constraint binds:
 $x^{(1)} = (0, 2)$, objective -4



Simplex algorithm: example

$$\begin{array}{ll}\text{minimize} & -x_1 - 2x_2 \\ \text{subject to} & x_1 + x_2 \leq 4 \\ & x_1 + 3x_2 \leq 6 \\ & x \geq 0\end{array}$$


- ▶ start at $x^{(0)} = (0, 0)$: objective 0
- ▶ increase x_2 until a constraint binds:
 $x^{(1)} = (0, 2)$, objective -4
- ▶ increase x_1 along the edge:
 $x^{(2)} = (3, 1)$, objective -5



Simplex algorithm: example

$$\begin{array}{ll}\text{minimize} & -x_1 - 2x_2 \\ \text{subject to} & x_1 + x_2 \leq 4 \\ & x_1 + 3x_2 \leq 6 \\ & x \geq 0\end{array}$$

- ▶ start at $x^{(0)} = (0, 0)$: objective 0
- ▶ increase x_2 until a constraint binds:
 $x^{(1)} = (0, 2)$, objective -4
- ▶ increase x_1 along the edge:
 $x^{(2)} = (3, 1)$, objective -5




Simplex algorithm: example

$$\begin{array}{ll}\text{minimize} & -x_1 - 2x_2 \\ \text{subject to} & x_1 + x_2 \leq 4 \\ & x_1 + 3x_2 \leq 6 \\ & x \geq 0\end{array}$$

- ▶ start at $x^{(0)} = (0, 0)$: objective 0
- ▶ increase x_2 until a constraint binds:
 $x^{(1)} = (0, 2)$, objective -4
- ▶ increase x_1 along the edge:
 $x^{(2)} = (3, 1)$, objective -5

Q: why does x_2 stop at 2, not 4?




Simplex algorithm: example

$$\begin{array}{ll}\text{minimize} & -x_1 - 2x_2 \\ \text{subject to} & x_1 + x_2 \leq 4 \\ & x_1 + 3x_2 \leq 6 \\ & x \geq 0\end{array}$$

- ▶ start at $x^{(0)} = (0, 0)$: objective 0
- ▶ increase x_2 until a constraint binds:
 $x^{(1)} = (0, 2)$, objective -4
- ▶ increase x_1 along the edge:
 $x^{(2)} = (3, 1)$, objective -5

Q: why does x_2 stop at 2, not 4?

A: $x_1 + 3x_2 \leq 6$ binds first



Simplex algorithm: example


$$\begin{array}{ll}\text{minimize} & -x_1 - 2x_2 \\ \text{subject to} & x_1 + x_2 \leq 4 \\ & x_1 + 3x_2 \leq 6 \\ & x \geq 0\end{array}$$

- ▶ start at $x^{(0)} = (0, 0)$: objective 0
- ▶ increase x_2 until a constraint binds:
 $x^{(1)} = (0, 2)$, objective -4
- ▶ increase x_1 along the edge:
 $x^{(2)} = (3, 1)$, objective -5

Q: why does x_2 stop at 2, not 4?

A: $x_1 + 3x_2 \leq 6$ binds first

Q: why stop at $x^{(2)}$?



Simplex algorithm: example

$$\begin{array}{ll}\text{minimize} & -x_1 - 2x_2 \\ \text{subject to} & x_1 + x_2 \leq 4 \\ & x_1 + 3x_2 \leq 6 \\ & x \geq 0\end{array}$$


- ▶ start at $x^{(0)} = (0, 0)$: objective 0
- ▶ increase x_2 until a constraint binds:
 $x^{(1)} = (0, 2)$, objective -4
- ▶ increase x_1 along the edge:
 $x^{(2)} = (3, 1)$, objective -5

Q: why does x_2 stop at 2, not 4?

A: $x_1 + 3x_2 \leq 6$ binds first

Q: why stop at $x^{(2)}$?

A: no neighboring vertex improves the objective



Simplex algorithm

basic idea: local search on the vertices of the feasible set

- ▶ start at BFS x and evaluate objective $c^T x$
- ▶ move to a neighboring BFS x' with better objective $c^T x'$
- ▶ repeat until no improvement possible

Simplex algorithm

basic idea: local search on the vertices of the feasible set

- ▶ start at BFS x and evaluate objective $c^T x$
- ▶ move to a neighboring BFS x' with better objective $c^T x'$
- ▶ repeat until no improvement possible

discuss in groups:

- ▶ how to find an initial BFS?
- ▶ how to find a neighboring BFS with better objective?
- ▶ how to prove optimality?

Finding an initial BFS

solve an auxiliary problem for which a BFS is known:

Finding an initial BFS

solve an auxiliary problem for which a BFS is known:

$$\begin{array}{ll}\text{minimize} & \sum_{i=1}^m z_i \\ \text{subject to} & Ax + Dz = b \\ & x, z \geq 0\end{array}$$

where $D \in \mathbb{R}^{m \times m}$ is a diagonal matrix with $D_{ii} = \mathbf{sign}(b_i)$ for $i = 1, \dots, m$.

Finding an initial BFS

solve an auxiliary problem for which a BFS is known:

$$\begin{array}{ll}\text{minimize} & \sum_{i=1}^m z_i \\ \text{subject to} & Ax + Dz = b \\ & x, z \geq 0\end{array}$$

where $D \in \mathbb{R}^{m \times m}$ is a diagonal matrix with $D_{ii} = \mathbf{sign}(b_i)$ for $i = 1, \dots, m$.

► $x = 0, z = |b|$ is a BFS of this problem

Finding an initial BFS

solve an auxiliary problem for which a BFS is known:

$$\begin{array}{ll}\text{minimize} & \sum_{i=1}^m z_i \\ \text{subject to} & Ax + Dz = b \\ & x, z \geq 0\end{array}$$

where $D \in \mathbb{R}^{m \times m}$ is a diagonal matrix with $D_{ii} = \mathbf{sign}(b_i)$ for $i = 1, \dots, m$.

- ▶ $x = 0, z = |b|$ is a BFS of this problem
- ▶ $(x, z) = (x, 0)$ is a BFS of this problem $\iff x$ is a BFS of the original problem

Find a better neighboring BFS

start with BFS x with active set S , $x_S > 0$. (called a **non-degenerate** BFS.)
construct the **j th basic direction** d^j by turning on variable $j \notin S$

$$x^+ \leftarrow x + \theta d^j, \quad \text{stepsize } \theta > 0$$

where $d_j^j = 1$ and $d_i^j = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S^j .

Find a better neighboring BFS

start with BFS x with active set S , $x_S > 0$. (called a **non-degenerate** BFS.)
construct the **j th basic direction** d^j by turning on variable $j \notin S$

$$x^+ \leftarrow x + \theta d^j, \quad \text{stepsize } \theta > 0$$

where $d_j^j = 1$ and $d_i^j = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S^j .

► need to stay feasible wrt equality constraints, so need

$$0 = Ad^j = A_S d_S^j + A_j \implies d_S^j = -A_S^{-1} A_j$$

Find a better neighboring BFS

start with BFS x with active set S , $x_S > 0$. (called a **non-degenerate** BFS.)
construct the **j th basic direction** d^j by turning on variable $j \notin S$

$$x^+ \leftarrow x + \theta d^j, \quad \text{stepsize } \theta > 0$$

where $d_j^j = 1$ and $d_i^j = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S^j .

- ▶ need to stay feasible wrt equality constraints, so need

$$0 = Ad^j = A_S d_S^j + A_j \implies d_S^j = -A_S^{-1} A_j$$

- ▶ as $x_S > 0$ is non-degenerate, $\exists \theta > 0$ st $x^+ \geq 0$

Find a better neighboring BFS

start with BFS x with active set S , $x_S > 0$. (called a **non-degenerate** BFS.)
construct the **j th basic direction** d^j by turning on variable $j \notin S$

$$x^+ \leftarrow x + \theta d^j, \quad \text{stepsize } \theta > 0$$

where $d_j^j = 1$ and $d_i^j = 0$ for $i \notin S \cup \{j\}$. need to solve for d_S^j .

- ▶ need to stay feasible wrt equality constraints, so need

$$0 = A d^j = A_S d_S^j + A_j \implies d_S^j = -A_S^{-1} A_j$$

- ▶ as $x_S > 0$ is non-degenerate, $\exists \theta > 0$ st $x^+ \geq 0$
- ▶ how does objective change if we move to $x^+ = x + \theta d^j$?

$$c^T x^+ - c^T x = \theta c^T d^j = \theta (c_j - c_S^T A_S^{-1} A_j)$$

Reduced cost

define **reduced cost** $\bar{c}_j = c_j - c_S^T A_S^{-1} A_j, j \notin S$

Reduced cost

define **reduced cost** $\bar{c}_j = c_j - c_S^T A_S^{-1} A_j, j \notin S$

facts:

- ▶ if $\bar{c} \geq 0$, x is optimal
- ▶ if x is optimal and nondegenerate ($x_S > 0$), then $\bar{c} \geq 0$

Reduced cost

define **reduced cost** $\bar{c}_j = c_j - c_S^T A_S^{-1} A_j, j \notin S$

facts:

- ▶ if $\bar{c} \geq 0$, x is optimal
- ▶ if x is optimal and nondegenerate ($x_S > 0$), then $\bar{c} \geq 0$

let's show the first fact.

if $\bar{c} \geq 0$, x is optimal

let x be a BFS with basis S , and let x' be any feasible point. Since

$$A_S x_S = b, \quad A_S x'_S + A_{\bar{S}} x'_{\bar{S}} = b,$$

we have

$$x'_S = x_S - A_S^{-1} A_{\bar{S}} x'_{\bar{S}}.$$

if $\bar{c} \geq 0$, x is optimal

let x be a BFS with basis S , and let x' be any feasible point. Since

$$A_S x_S = b, \quad A_S x'_S + A_{\bar{S}} x'_{\bar{S}} = b,$$

we have

$$x'_S = x_S - A_S^{-1} A_{\bar{S}} x'_{\bar{S}}.$$

substitute into the objective:

$$\begin{aligned} c^T x' &= c_S^T x'_S + c_{\bar{S}}^T x'_{\bar{S}} \\ &= c_S^T x_S + \left(c_{\bar{S}}^T - c_S^T A_S^{-1} A_{\bar{S}} \right) x'_{\bar{S}} \\ &= c^T x + \bar{c}_{\bar{S}}^T x'_{\bar{S}} \\ &\geq c^T x, \end{aligned}$$

because $\bar{c}_{\bar{S}} \geq 0$ and $x'_{\bar{S}} \geq 0$.

if $\bar{c} \geq 0$, x is optimal

let x be a BFS with basis S , and let x' be any feasible point. Since

$$A_S x_S = b, \quad A_S x'_S + A_{\bar{S}} x'_{\bar{S}} = b,$$

we have

$$x'_S = x_S - A_S^{-1} A_{\bar{S}} x'_{\bar{S}}.$$

substitute into the objective:

$$\begin{aligned} c^T x' &= c_S^T x'_S + c_{\bar{S}}^T x'_{\bar{S}} \\ &= c_S^T x_S + \left(c_{\bar{S}}^T - c_S^T A_S^{-1} A_{\bar{S}} \right) x'_{\bar{S}} \\ &= c^T x + \bar{c}_{\bar{S}}^T x'_{\bar{S}} \\ &\geq c^T x, \end{aligned}$$

because $\bar{c}_{\bar{S}} \geq 0$ and $x'_{\bar{S}} \geq 0$. therefore no feasible point has lower objective value than x , so x is optimal.

Outline

LP standard form

LP inequality form

What kinds of points can be optimal?

Solving LPs

Modeling

LP solvers

practical solvers for LP and MILP:

- ▶ Gurobi, FICO Xpress, IBM CPLEX, and COPT are leading commercial solvers

LP solvers

practical solvers for LP and MILP:

- ▶ Gurobi, FICO Xpress, IBM CPLEX, and COPT are leading commercial solvers
- ▶ Gurobi is an excellent default when licenses are available
- ▶ HiGHS is the strongest open-source default for LP; it is available in SciPy via `linprog(method="highs")`

LP solvers

practical solvers for LP and MILP:

- ▶ Gurobi, FICO Xpress, IBM CPLEX, and COPT are leading commercial solvers
- ▶ Gurobi is an excellent default when licenses are available
- ▶ HiGHS is the strongest open-source default for LP; it is available in SciPy via `linprog(method="highs")`
- ▶ GLPK and SCIP are also free, but are usually slower for pure LP
- ▶ The fastest solver depends on the problem structure, warm starts, memory limits, and numerical conditioning; benchmark representative instances

LP solvers

practical solvers for LP and MILP:

- ▶ Gurobi, FICO Xpress, IBM CPLEX, and COPT are leading commercial solvers
- ▶ Gurobi is an excellent default when licenses are available
- ▶ HiGHS is the strongest open-source default for LP; it is available in SciPy via `linprog(method="highs")`
- ▶ GLPK and SCIP are also free, but are usually slower for pure LP
- ▶ The fastest solver depends on the problem structure, warm starts, memory limits, and numerical conditioning; benchmark representative instances
- ▶ `gurobipy` is a python interface to Gurobi
- ▶ CVX* (including CVXPY in python) are modeling languages that call solvers with good support for convex problems

LP solvers

practical solvers for LP and MILP:

- ▶ Gurobi, FICO Xpress, IBM CPLEX, and COPT are leading commercial solvers
- ▶ Gurobi is an excellent default when licenses are available
- ▶ HiGHS is the strongest open-source default for LP; it is available in SciPy via `linprog(method="highs")`
- ▶ GLPK and SCIP are also free, but are usually slower for pure LP
- ▶ The fastest solver depends on the problem structure, warm starts, memory limits, and numerical conditioning; benchmark representative instances
- ▶ `gurobipy` is a python interface to Gurobi
- ▶ CVX* (including CVXPY in python) are modeling languages that call solvers with good support for convex problems
- ▶ JuliaOpt/JuMP is a modeling language in Julia that calls solvers and is super speedy for MILP applications

LP solvers

practical solvers for LP and MILP:

- ▶ Gurobi, FICO Xpress, IBM CPLEX, and COPT are leading commercial solvers
- ▶ Gurobi is an excellent default when licenses are available
- ▶ HiGHS is the strongest open-source default for LP; it is available in SciPy via `linprog(method="highs")`
- ▶ GLPK and SCIP are also free, but are usually slower for pure LP
- ▶ The fastest solver depends on the problem structure, warm starts, memory limits, and numerical conditioning; benchmark representative instances
- ▶ `gurobipy` is a python interface to Gurobi
- ▶ CVX* (including CVXPY in python) are modeling languages that call solvers with good support for convex problems
- ▶ JuliaOpt/JuMP is a modeling language in Julia that calls solvers and is super speedy for MILP applications demo:
 - ▶ power systems
https://jump.dev/JuMP.jl/stable/tutorials/applications/power_systems/

Oro Verde case + tutorial

<https://github.com/stanford-cme-307/demos/tree/main/gurobipy>

A modeling workflow

an optimization model describes exactly which decisions are allowed

1. define each decision variable and its bounds
2. translate one requirement at a time
3. test the formulation on boundary cases

A modeling workflow

an optimization model describes exactly which decisions are allowed

1. define each decision variable and its bounds
2. translate one requirement at a time
3. test the formulation on boundary cases

three recurring ideas:

- ▶ introduce an auxiliary variable to expose a linear relation
- ▶ use binary variables to encode yes/no decisions
- ▶ use bounds to turn conditional constraints on and off

Two standard-form reformulations

feature	new variables	substitution
$a^T x \leq b$ in standard form	$s \geq 0$	$a^T x + s = b$
$x_i \in \mathbb{R}$ in standard form	$x_i^+, x_i^- \geq 0$	$x_i = x_i^+ - x_i^-$

Two standard-form reformulations

feature	new variables	substitution
$a^T x \leq b$ in standard form	$s \geq 0$	$a^T x + s = b$
$x_i \in \mathbb{R}$ in standard form	$x_i^+, x_i^- \geq 0$	$x_i = x_i^+ - x_i^-$

substitute these identities everywhere, including the objective

$$c_i x_i \longmapsto c_i x_i^+ - c_i x_i^-.$$

The epigraph trick

the **epigraph** of f contains every point above its graph:

$$\text{epi } f = \{(x, t) : f(x) \leq t\}.$$

The epigraph trick

the **epigraph** of f contains every point above its graph:

$$\text{epi } f = \{(x, t) : f(x) \leq t\}.$$

minimizing the upper bound t makes it tight:

$$\min_x \sum_i |x_i| \quad \Longleftrightarrow \quad \begin{array}{ll} \text{minimize}_{x,t} & \mathbf{1}^T t \\ \text{subject to} & -t \leq x \leq t, \end{array}$$

$$\min_x \max_i f_i(x) \quad \Longleftrightarrow \quad \begin{array}{ll} \text{minimize}_{x,t} & t \\ \text{subject to} & f_i(x) \leq t, \quad \forall i. \end{array}$$

add the original constraints to both sides; the right-hand problems are LPs when each f_i is affine

Exercise: worst-case residual

reformulate as an LP:

$$\underset{x}{\text{minimize}} \quad \max_{i=1,\dots,m} |a_i^T x - b_i|.$$

Exercise: worst-case residual

reformulate as an LP:

$$\underset{x}{\text{minimize}} \quad \max_{i=1,\dots,m} |a_i^T x - b_i|.$$

A:

$$\begin{aligned} &\underset{x,t}{\text{minimize}} \quad t \\ &\text{subject to} \quad -t \leq a_i^T x - b_i \leq t, \quad i = 1, \dots, m. \end{aligned}$$

Each constraint gives $|a_i^T x - b_i| \leq t$; the objective finds the smallest uniform bound.

Binary variables encode propositions

let $z_A, z_B \in \{0, 1\}$ record whether projects A and B are funded

$$\text{"if } A, \text{ then } B\text{"} \quad \Longleftrightarrow \quad z_A \leq z_B$$

Binary variables encode propositions

let $z_A, z_B \in \{0, 1\}$ record whether projects A and B are funded

$$\text{"if } A, \text{ then } B" \iff z_A \leq z_B$$

z_A	z_B	$z_A \leq z_B?$
0	0	yes
0	1	yes
1	0	no
1	1	yes

Binary variables encode propositions

let $z_A, z_B \in \{0, 1\}$ record whether projects A and B are funded

$$\text{"if } A, \text{ then } B" \iff z_A \leq z_B$$

z_A	z_B	$z_A \leq z_B?$
0	0	yes
0	1	yes
1	0	no
1	1	yes

the inequality excludes exactly the forbidden truth assignment; binary variables make the model a mixed-integer LP

Counting binary decisions

for a binary variable, $1 - z_i$ encodes “not i ”; sums count how many propositions are true:

$$\text{at least one} : \sum_i z_i \geq 1,$$

$$\text{at most one} : \sum_i z_i \leq 1,$$

$$\text{exactly } k : \sum_i z_i = k.$$

Counting binary decisions

for a binary variable, $1 - z_i$ encodes “not i ”; sums count how many propositions are true:

$$\text{at least one} : \sum_i z_i \geq 1,$$

$$\text{at most one} : \sum_i z_i \leq 1,$$

$$\text{exactly } k : \sum_i z_i = k.$$

an implication between binary variables is also a counting constraint:

$$z_A = 1 \implies z_B = 1 \quad \iff \quad z_A \leq z_B$$

Exercise: project logic

write linear constraints for the following rules:

1. fund A only if E is funded
2. if A is not funded, then fund G
3. fund exactly one of B and C
4. funding A or B requires funding H

Exercise: project logic

write linear constraints for the following rules:

1. fund A only if E is funded
2. if A is not funded, then fund G
3. fund exactly one of B and C
4. funding A or B requires funding H

$$z_A \leq z_E$$

Exercise: project logic

write linear constraints for the following rules:

1. fund A only if E is funded
2. if A is not funded, then fund G
3. fund exactly one of B and C
4. funding A or B requires funding H

$$z_A \leq z_E$$

$$1 - z_A \leq z_G$$

Exercise: project logic

write linear constraints for the following rules:

1. fund A only if E is funded
2. if A is not funded, then fund G
3. fund exactly one of B and C
4. funding A or B requires funding H

$$z_A \leq z_E$$

$$1 - z_A \leq z_G$$

$$z_B + z_C = 1$$

Exercise: project logic

write linear constraints for the following rules:

1. fund A only if E is funded

$$z_A \leq z_E$$

2. if A is not funded, then fund G

$$1 - z_A \leq z_G$$

3. fund exactly one of B and C

$$z_B + z_C = 1$$

4. funding A or B requires funding H

$$z_A \leq z_H, \quad z_B \leq z_H$$

Exercise: project logic

write linear constraints for the following rules:

1. fund A only if E is funded

$$z_A \leq z_E$$

2. if A is not funded, then fund G

$$1 - z_A \leq z_G$$

3. fund exactly one of B and C

$$z_B + z_C = 1$$

4. funding A or B requires funding H

$$z_A \leq z_H, \quad z_B \leq z_H$$

Q: Why is $z_A + z_B \leq z_H$ too strong?

Exercise: project logic

write linear constraints for the following rules:

1. fund A only if E is funded

$$z_A \leq z_E$$

2. if A is not funded, then fund G

$$1 - z_A \leq z_G$$

3. fund exactly one of B and C

$$z_B + z_C = 1$$

4. funding A or B requires funding H

$$z_A \leq z_H, \quad z_B \leq z_H$$

Q: Why is $z_A + z_B \leq z_H$ too strong?

A: It forbids funding both A and B , even when H is funded.

Conditional linear constraints

suppose $z \in \{0, 1\}$ and we require

$$z = 1 \implies a^T x \leq b.$$

Conditional linear constraints

suppose $z \in \{0, 1\}$ and we require

$$z = 1 \implies a^T x \leq b.$$

make the constraint nonbinding when $z = 0$:

$$a^T x \leq b + M(1 - z).$$

Conditional linear constraints

suppose $z \in \{0, 1\}$ and we require

$$z = 1 \implies a^T x \leq b.$$

make the constraint nonbinding when $z = 0$:

$$a^T x \leq b + M(1 - z).$$

z	resulting constraint	desired effect
1	$a^T x \leq b$	constraint on
0	$a^T x \leq b + M$	constraint off

Conditional linear constraints

suppose $z \in \{0, 1\}$ and we require

$$z = 1 \implies a^T x \leq b.$$

make the constraint nonbinding when $z = 0$:

$$a^T x \leq b + M(1 - z).$$

z	resulting constraint	desired effect
1	$a^T x \leq b$	constraint on
0	$a^T x \leq b + M$	constraint off

M must cover the largest possible violation of $a^T x \leq b$

Bounds determine a valid big- M

if x already lies in a known set X , the tightest nonnegative choice is

$$M = \max \left\{ 0, \max_{x \in X} (a^T x - b) \right\}.$$

Bounds determine a valid big- M

if x already lies in a known set X , the tightest nonnegative choice is

$$M = \max \left\{ 0, \max_{x \in X} (a^T x - b) \right\}.$$

classroom capacity. $X_{ij} = 1$ means class i uses room j . Suppose enrollment is a decision with $0 \leq p_i \leq \bar{p}_i$.

$$X_{ij} = 1 \implies p_i \leq c_j$$

becomes

$$p_i \leq c_j + \underbrace{\max\{0, \bar{p}_i - c_j\}}_{M_{ij}}(1 - X_{ij}).$$

Bounds determine a valid big- M

if x already lies in a known set X , the tightest nonnegative choice is

$$M = \max \left\{ 0, \max_{x \in X} (a^T x - b) \right\}.$$

classroom capacity. $X_{ij} = 1$ means class i uses room j . Suppose enrollment is a decision with $0 \leq p_i \leq \bar{p}_i$.

$$X_{ij} = 1 \implies p_i \leq c_j$$

becomes

$$p_i \leq c_j + \underbrace{\max\{0, \bar{p}_i - c_j\}}_{M_{ij}}(1 - X_{ij}).$$

if p_i is fixed data, preprocess data to remove ineligible assignments; no need for big- M

Exercise: a plant that can be on or off

the decisions are production $x \geq 0$ and operating status $z \in \{0, 1\}$.

the data are:

- ▶ $0 < L \leq U$: minimum and maximum production when the plant operates
- ▶ $c \geq 0$: production cost per unit
- ▶ $f \geq 0$: fixed cost whenever the plant operates

Exercise: a plant that can be on or off

the decisions are production $x \geq 0$ and operating status $z \in \{0, 1\}$.

the data are:

- ▶ $0 < L \leq U$: minimum and maximum production when the plant operates
- ▶ $c \geq 0$: production cost per unit
- ▶ $f \geq 0$: fixed cost whenever the plant operates

Q: Model the feasible production levels and the production cost.

Exercise: a plant that can be on or off

the decisions are production $x \geq 0$ and operating status $z \in \{0, 1\}$.

the data are:

- ▶ $0 < L \leq U$: minimum and maximum production when the plant operates
- ▶ $c \geq 0$: production cost per unit
- ▶ $f \geq 0$: fixed cost whenever the plant operates

Q: Model the feasible production levels and the production cost.

A:

$$Lz \leq x \leq Uz, \quad \text{cost} = cx + fz.$$

If $z = 0$, then $x = 0$; if $z = 1$, then $L \leq x \leq U$.

An equivalence needs two implications

let $q \in \{0, 1, \dots, U\}$ and define

$$z = 1 \iff q \geq k.$$

An equivalence needs two implications

let $q \in \{0, 1, \dots, U\}$ and define

$$z = 1 \iff q \geq k.$$

the two implications have an exact linear formulation:

$$q \geq kz, \quad q \leq (k - 1) + (U - k + 1)z.$$

An equivalence needs two implications

let $q \in \{0, 1, \dots, U\}$ and define

$$z = 1 \iff q \geq k.$$

the two implications have an exact linear formulation:

$$q \geq kz, \quad q \leq (k-1) + (U-k+1)z.$$

$$\begin{array}{l|l} z = 0 & q \leq k-1 \\ z = 1 & q \geq k \end{array}$$

An equivalence needs two implications

let $q \in \{0, 1, \dots, U\}$ and define

$$z = 1 \iff q \geq k.$$

the two implications have an exact linear formulation:

$$q \geq kz, \quad q \leq (k-1) + (U-k+1)z.$$

$$\begin{array}{l|l} z = 0 & q \leq k-1 \\ z = 1 & q \geq k \end{array}$$

for continuous q , the reverse implication introduces a strict boundary; use a nonzero tolerance, or allow either value of z at $q = k$

Exercise: an indicator for a count

five projects have selection variables $x_1, \dots, x_5 \in \{0, 1\}$. Let $z = 1$ exactly when at least three projects are selected.

Exercise: an indicator for a count

five projects have selection variables $x_1, \dots, x_5 \in \{0, 1\}$. Let $z = 1$ exactly when at least three projects are selected.

Q: Find two linear constraints that define z .

Exercise: an indicator for a count

five projects have selection variables $x_1, \dots, x_5 \in \{0, 1\}$. Let $z = 1$ exactly when at least three projects are selected.

Q: Find two linear constraints that define z .

A: Set $q = \sum_{i=1}^5 x_i$, $U = 5$, and $k = 3$:

$$\sum_{i=1}^5 x_i \geq 3z, \quad \sum_{i=1}^5 x_i \leq 2 + 3z.$$

The first implication forces the count up when $z = 1$; the second forces $z = 1$ when the count reaches three.

Logical modeling: the short version

- ▶ encode propositions with binary variables

Logical modeling: the short version

- ▶ encode propositions with binary variables
- ▶ use sums of binaries to count choices

Logical modeling: the short version

- ▶ encode propositions with binary variables
- ▶ use sums of binaries to count choices
- ▶ translate each implication separately

Logical modeling: the short version

- ▶ encode propositions with binary variables
- ▶ use sums of binaries to count choices
- ▶ translate each implication separately
- ▶ deactivate a linear constraint using a bound-derived constant

Logical modeling: the short version

- ▶ encode propositions with binary variables
- ▶ use sums of binaries to count choices
- ▶ translate each implication separately
- ▶ deactivate a linear constraint using a bound-derived constant
- ▶ test every formulation at $z = 0$ and $z = 1$

Logical modeling: the short version

- ▶ encode propositions with binary variables
- ▶ use sums of binaries to count choices
- ▶ translate each implication separately
- ▶ deactivate a linear constraint using a bound-derived constant
- ▶ test every formulation at $z = 0$ and $z = 1$

solver indicator constraints express implications directly, but good variable bounds still matter

Exit ticket

Please fill out the exit ticket (linked from course schedule spreadsheet) before you leave:

- ▶ Helps us improve the course

We will not grade this, but we will read it!

