


# MS&E 228: Inference with Modern Non-Linear Prediction

Vasilis Syrgkanis

MS&E, Stanford







# Recap of Last Lecture

# Causal Inference Pipeline

Theory




Practice




# Causal Inference Pipeline

Theory



Practice



# Goals for Today

- Methods for Confidence Intervals for ATE with non-linear models
  - General Neyman Orthogonality Framework (Double/Debiased ML)
  - Methods for Confidence Intervals for ATE in a partially-linear model
  - Sample-splitting and cross-fitting
- 
- Proof sketch of main theorem\*

# The Example Problem



# Identification under Conditional Ignorability

- Once we condition on enough variables  $X$  that affect treatment assignment, remnant variation in  $D$  is exogenous (as-if trial)

$$Y^{(d)} \perp\!\!\!\perp D \mid X \quad (\text{conditional ignorability})$$

- Why useful:

$$\begin{aligned} E[Y \mid D = d, X] &= E[Y^{(D)} \mid D = d, X] \\ &= E[Y^{(d)} \mid D = d, X] = E[Y^{(d)} \mid X] \end{aligned}$$

- Average treatment effect is “identified” as (g-formula):

$$\begin{aligned} \theta_0 &= E[Y^{(1)} - Y^{(0)}] = E[E[Y^{(1)} - Y^{(0)} \mid X]] \\ &= E[E[Y \mid D = 1, X] - E[Y \mid D = 0, X]] \end{aligned}$$

# Let's take it to data

- We observe  $n$  samples  $Z_1, \dots, Z_n$  where  $Z_i = (X_i, D_i, Y_i)$

- Want to estimate average effect  $\theta_0$ , which satisfies:

$$\theta_0 = E[g_0(1, X) - g_0(0, X)]$$

- Where:

$$g_0(D, X) := E[Y \mid D, X]$$

- We want to be able to use ML to learn regression function  $g_0$ !

# What do we want from $\hat{\theta}$ ?

- **Ideally** parametric rates for  $\theta_0$  even when we have slower rates for  $g_0$
- **Ideally** construction of confidence intervals for  $\theta_0$

- **One approach.** Asymptotic normality

$$\sqrt{n}(\hat{\theta} - \theta_0) \rightarrow_d N(0, \sigma^2)$$

- Implies construction of approximately correct confidence intervals

$$\text{with prob. } \approx 95\%: \theta_0 \in [\hat{\theta} \pm 1.96\hat{\sigma}/\sqrt{n}]$$

# Natural Estimation Algorithm

- Estimate  $\hat{g}$  of  $g_0$  from data
- Calculate empirical plug-in average:

$$\hat{\theta} := \frac{1}{n} \sum_{i=1}^n \hat{g}(1, X) - \hat{g}(0, X)$$

# Natural Algorithm Gone Wrong


```
def est(X, D, y): # direct non-orthogonal estimator of average effect
    est = RandomForestRegressor(min_samples_leaf=20)
    est.fit(np.hstack([D.reshape(-1, 1), X]), y)
    ones = np.hstack([np.ones((X.shape[0], 1)), X])
    zeros = np.hstack([np.zeros((X.shape[0], 1)), X])
    preds = est.predict(ones) - est.predict(zeros)
    return np.mean(preds), np.std(preds)/np.sqrt(X.shape[0])
```

## Simple Example

$$X \sim N(0, I_{20})$$

$$D \sim \text{Binomial}(0.5 + \text{clip}(X_0, -0.4, 0.4))$$

$$y \sim \theta_0 D + X_0 + X_1 + N(0, 1)$$



# Natural Estimation Algorithm (Draft 2)

- Split the data in half  $S_1, S_2$
- On first half  $S_1$ , estimate  $\hat{g}$  of  $g_0$
- Calculate empirical plug-in average on second half  $S_2$ :

$$\hat{\theta} := \frac{1}{|S_2|} \sum_{i \in S_2} \hat{g}(1, X) - \hat{g}(0, X)$$

# Natural Estimation Algorithm (Draft 3)

- Split data in  $K$  parts,  $S_1, \dots, S_K$
- For each part  $k$ , estimate  $\hat{g}_k$  using data from all parts except  $S_k$
- Calculate average over all data:

$$\hat{\theta} = \frac{1}{n} \sum_{k=1}^K \sum_{i \in S_k} \hat{g}_k(1, X) - \hat{g}_k(0, X)$$

# Natural Algorithm (Draft 3) Gone Wrong


```
def est2(X, D, y): # direct non-orthogonal estimator with sample splitting
    effects = np.zeros(X.shape[0])
    for train, test in KFold(n_splits=3).split(X):
        est = RandomForestRegressor(min_samples_leaf=20)
        est.fit(np.hstack([D[train].reshape(-1, 1), X[train]]), y[train])
        ones = np.hstack([np.ones((X[test].shape[0], 1)), X[test]])
        zeros = np.hstack([np.zeros((X[test].shape[0], 1)), X[test]])
        effects[test] = est.predict(ones) - est.predict(zeros)
    return np.mean(effects), np.std(effects)/np.sqrt(X.shape[0])
```

## Simple Example

$$X \sim N(0, I_{20})$$

$$D \sim \text{Binomial}(0.5 + \text{clip}(X_0, -0.4, 0.4))$$

$$y \sim \theta_0 D + X_0 + X_1 + N(0, 1)$$





When is estimate  $\hat{\theta}$   $\sqrt{n}$ -asymptotically normal?

When is estimate  $\hat{\theta}$   $\sqrt{n}$ -asymptotically normal?  
We need to change the moment we use

# Debiased Machine Learning

# Average Causal Effect Example

- We observe  $n$  samples  $Z_1, \dots, Z_n$  where  $Z_i = (X_i, D_i, Y_i)$
- Want to estimate average effect  $\theta_0$ , which satisfies:

$$\theta_0 := E[g_0(1, X) - g_0(0, X)]$$

- Where:

$$g_0(D, X) := E[Y \mid D, X]$$

- The identification formula for  $\theta_0$  is sensitive to variations in  $g$
- Any bias or error in  $g$  propagates to bias or error in moment and  $\hat{\theta}$
- Can we add a correction that corrects the biases of  $\hat{g}$

# Better Formula for ATE

Regression residual is a proxy that  $g$  is biased

- **Key Idea.** Add a debiasing correction

$$M(g, a) = E[g(1, X) - g(0, X)] + E[a(D, X) (Y - g(D, X))]$$

- What is  $a_0$ ?
- Insensitivity: Take derivative with respect to  $g$  at  $\theta_0, g_0, a_0$  in any direction  $v \in G$

$$\left. \frac{\partial}{\partial t} M(g_0 + t v, a_0) \right|_{t=0} = E[v(1, X) - v(0, X)] - E[a(D, X) v(D, X)] = 0$$

- If this holds then if  $g$  is very wrong but  $a$  is correct:

$$\begin{aligned} \theta &= E[a_0(D, X)Y] = E[a_0(D, X)E[Y \mid D, X]] \\ &= E[a_0(D, X)g(D, X)] = E[g(1, X) - g(0, X)] \end{aligned}$$

# Inverse Propensity Weighting (IPW)

- The following works: inverse propensity scoring

$$a_0(D, X) = \frac{D}{\Pr[D = 1|X]} - \frac{1 - D}{\Pr[D = 0|X]}$$

- Sketch:

$$\begin{aligned} E \left[ \frac{D}{\Pr[D = 1|X]} g(D, X) \right] &= E \left[ \frac{D}{\Pr[D = 1|X]} g(1, X) \right] \\ &= E \left[ \frac{E[D|X]}{\Pr[D = 1|X]} g(1, X) \right] \\ &= E[g(1, X)] \end{aligned}$$

# New Formula is Insensitive

$$M(g, a) = E[g(1, X) - g(0, X)] + E[a(D, X) (Y - g(D, X))]$$

- Take derivative with respect to  $g$  at  $g_0, a_0$  in any direction  $v \in G$

$$\left. \frac{\partial}{\partial t} M(g_0 + t v, a_0) \right|_{t=0} = E[v(1, X) - v(0, X)] - E[a(D, X) v(D, X)] = 0$$

Take derivative with respect to  $a$  at  $g_0, a_0$  in any direction  $v \in A$

$$\left. \frac{\partial}{\partial t} M(g_0, a_0 + t v) \right|_{t=0} = E[v(D, X) (Y - g_0(D, X))] = 0$$

# Asymptotic Normality of De-biased Estimate

$$\hat{\theta} := E_n[\hat{g}(1, X) - \hat{g}(0, X) + \hat{a}(D, X) \cdot (Y - \hat{g}(D, X))]$$

- Assume that propensities are bounded away from 0 and 1 (strict overlap)
- Assume  $\hat{g}, \hat{a}$  estimated on separate sample (or cross-fitting), are consistent and:

$$\sqrt{n} E[(a_0(D, X) - \hat{a}(D, X))(\hat{g}(D, X) - g_0(D, X))] \rightarrow_p 0$$

- Assume random variables  $Y, a(D, X), g(D, X)$  have bounded fourth moments
- Then:

$$\sqrt{n}(\hat{\theta} - \theta_0) \rightarrow_d N(0, \sigma^2), \quad \hat{\sigma}^2 = Var_n(\hat{g}(1, X) - \hat{g}(0, X) + \hat{a}(X) \cdot (Y - \hat{g}(X)))$$



# Python Pseudocode

```
cv = KFold(n_splits=nfolds, shuffle=True, random_state=123)
yhat0, yhat1 = np.zeros(y.shape), np.zeros(y.shape)
# we will fit a model  $E[Y | D, X]$  by fitting a separate model for  $D=0$ 
# and a separate model for  $D=1$ .
for train, test in cv.split(X, y):
    # train a model on training data that received zero and predict on all test data
    yhat0[test] = modely.fit(X[train][D[train]==0], y[train][D[train]==0]).predict(X[test])
    # train a model on training data that received one and predict on all test data
    yhat1[test] = modely.fit(X[train][D[train]==1], y[train][D[train]==1]).predict(X[test])
# prediction for observed treatment
yhat = yhat0 * (1 - D) + yhat1 * D
# propensity scores
Dhat = cross_val_predict(modeld, X, D, cv=cv, method='predict_proba', n_jobs=-1)[:, 1]
Dhat = np.clip(Dhat, trimming, 1 - trimming)
# doubly robust quantity for every sample
drhat = yhat1 - yhat0 + (y - yhat) * (D/Dhat - (1 - D)/(1 - Dhat))
point = np.mean(drhat)
var = np.var(drhat)
stderr = np.sqrt(var / X.shape[0])
return point, stderr, yhat, Dhat, y - yhat, D - Dhat, drhat
```

# Continuous Treatments under Partial Linearity

# Partially Linear Model

- Relevant in many applications: dose-response curve in healthcare, effect of price on demand, return-on-investment
- Assume conditional exogeneity

$$Y^{(d)} \perp\!\!\!\perp D \mid X$$

- Assume partially linear response

$$Y = \theta_0 D + f_0(X) + \epsilon, \quad E[\epsilon \mid D, X] = 0$$

- Equivalently, a partial linearity condition on the conditional expectation function

$$g_0(D, X) = E[Y \mid D, X] = \theta_0 D + f_0(X)$$

- Parameter of interest  $\theta_0$  is constant marginal effect of treatment

# Generalization of FWL Theorem

- Let's define a slight variant of residualization

$$\tilde{V} = V - E[V|X]$$

- Generalization of FWL theorem to partially linear models

$$\tilde{Y} = \theta_0 \tilde{D} + \epsilon, \quad E[\epsilon|\tilde{D}] = 0$$

- Let's consider the residual outcome


$$\begin{aligned}\tilde{Y} &= Y - E[Y|X] \\ &= \theta_0 D + f_0(X) + \epsilon - E[\theta_0 D + f_0(X) + \epsilon|X] \\ &= \theta_0 D + f_0(X) + \epsilon - \theta_0 E[D|X] - f_0(X) \\ &= \theta_0 (D - E[D|X]) + \epsilon\end{aligned}$$

Regression model  $h_0(X)$  predicting  
the outcome from the controls

Regression model  $p_0(X)$  predicting  
the treatment from the controls


# Orthogonal Method: Double ML

- **Double ML.** Split samples in half
  - Regress  $Y \sim X$  with ML on first half, to get estimate  $\hat{h}(S)$  of  $E[Y|X]$
  - Regress  $D \sim X$  with ML on first half, to get estimate  $\hat{p}(S)$  of  $E[D|X]$



# Orthogonal Method: Double ML

- **Double ML.** Split samples in half
  - Regress  $Y \sim X$  with ML on first half, to get estimate  $\hat{h}(S)$  of  $E[Y|X]$
  - Regress  $D \sim X$  with ML on first half, to get estimate  $\hat{p}(S)$  of  $E[D|X]$
  - Construct residuals on other half,  $\tilde{D} := D - \hat{p}(X)$  and  $\tilde{Y} := Y - \hat{h}(X)$
  - Run OLS on residuals:  $\tilde{Y} \sim \tilde{D}$  to get  $\hat{\theta}$
- Final OLS, *in population limit*, equivalent to solving normal equation:
$$E[(\tilde{Y} - \theta \tilde{D})\tilde{D}] = 0$$
- Define the *formula*:
$$M(\theta; h, p) = E \left[ \left( Y - h(X) - \theta (D - p(X)) \right) (D - p(X)) \right]$$
- Final OLS, *in population limit*, equivalent to solving for  $\theta$ :
$$M(\theta; h_0, p_0) = 0$$



# Insensitivity of Double ML Method

- The *formula*  $M$  is insensitive to the nuisance functions  $h, p$

$$M(\theta, h, p) = E \left[ \left( Y - h(X) - \theta (D - p(X)) \right) (D - p(X)) \right]$$

- Directional derivative with respect to  $h$

$$\left. \frac{\partial}{\partial t} M(\theta_0; h_0, p_0 + t v) \right|_{t=0} = -E[v(X)(D - p_0(X))] = E[v(X)E[D - p_0(X) | X]] = 0$$

- Directional derivative with respect to  $p$

$$\left. \frac{\partial}{\partial t} M(\theta_0; h_0, p_0 + t v) \right|_{t=0} = E[v(X)(D - p_0(X))] - E \left[ \left( Y - h_0(X) - \theta_0(D - p_0(X)) \right) v(X) \right] = 0$$

# Asymptotic Normality of DoubleML Estimate

$$E_n[(\hat{Y} - \theta \hat{D})\hat{D}] = 0 \Leftrightarrow \hat{\theta} = \frac{E_n[\hat{Y}\hat{D}]}{E_n[\hat{D}^2]}$$

- Assume that  $E[\tilde{D}^2] = E[\text{Var}(D | X)] > 0$  (average overlap)
- Assume  $\hat{h}, \hat{p}$  estimated on separate sample (or cross-fitting), are consistent and:

$$\sqrt{n} \left( \text{RMSE}(\hat{h}) \cdot \text{RMSE}(\hat{p}) + \text{RMSE}(\hat{p})^2 \right) \rightarrow_p 0$$

- Assume random variables  $Y, D, X$  have bounded fourth moments

$$\sqrt{n}(\hat{\theta} - \theta_0) \rightarrow_d N(0, \sigma^2),$$

Estimate asymptotically normal

$$\sigma^2 := \frac{E[(\tilde{Y} - \theta_0 \tilde{D})^2 \tilde{D}^2]}{E[\tilde{D}^2]^2},$$

Asymptotic variance


$$\hat{\sigma}^2 = \frac{E_n[(\hat{Y} - \hat{\theta} \hat{D})^2 \hat{D}^2]}{E_n[\hat{D}^2]^2}$$

Estimate of variance  $\Rightarrow$  95% CI  $\left[ \theta \pm 1.96 \frac{\hat{\sigma}}{\sqrt{n}} \right]$



# Natural Algorithm (Draft 3) Gone Right


```
def dml2(X, D, y): # orthogonal dml with sample-splitting
    est_y = RandomForestRegressor(min_samples_leaf=20)
    yres = y - cross_val_predict(est_y, X, y, cv=3)
    est_t = RandomForestRegressor(min_samples_leaf=20)
    Dres = D - cross_val_predict(est_t, X, D, cv=3)
    theta = np.mean(yres * Dres) / np.mean(Dres**2)
    var = np.mean((Dres**2) * (yres - theta*Dres)**2) / np.mean(Dres**2)
    stderr = np.sqrt(var / X.shape[0])
    return theta, stderr
```



# Natural Algorithm (Draft 3) Gone Right

```
from econml.dml import LinearDML

dml = LinearDML(model_y=RandomForestRegressor(min_samples_leaf=20),
                 model_t=RandomForestRegressor(min_samples_leaf=20))
est.fit(y, D, W=X).effect_inference()
```



# Proof sketch

$$\begin{aligned}\sqrt{n}(\hat{\theta} - \theta_0) &= \sqrt{n} \left( \frac{E_n[\hat{Y}\hat{D}]}{E_n[\hat{D}^2]} - \theta_0 \right) \\ &\approx \sqrt{n} \left( \frac{E_n[\tilde{Y}\tilde{D}]}{E_n[\tilde{D}^2]} - \theta_0 \right) \\ &= \sqrt{n} \left( \frac{E_n[\tilde{Y}\tilde{D}]}{E_n[\tilde{D}^2]} - \frac{E_n[\tilde{D}^2]}{E_n[\tilde{D}^2]} \theta_0 \right) \\ &= \sqrt{n} \left( \frac{E_n[(\tilde{Y} - \theta_0 \tilde{D})\tilde{D}]}{E_n[\tilde{D}^2]} \right)\end{aligned}$$

Due to insensitivity of the formula and fast enough rates of the nuisance models, we can ignore the error in the residuals

$$\approx \sqrt{n} \left( \frac{E_n[(\tilde{Y} - \theta_0 \tilde{D})\tilde{D}]}{E[\tilde{D}^2]} \right) = \sqrt{n} E_n \left[ \frac{(\tilde{Y} - \theta_0 \tilde{D})\tilde{D}}{E[\tilde{D}^2]} \right] \rightarrow_d N \left( 0, \text{Var} \left( \frac{(\tilde{Y} - \theta_0 \tilde{D})\tilde{D}}{E[\tilde{D}^2]} \right) \right)$$

Central Limit Theorem

By law of large numbers, the denominator can be replaced by the expectation,  $E_n[\tilde{D}^2] \approx E[\tilde{D}^2]$

# (Extended) Partially Linear Model

- Relevant in many applications: dose-response curve in healthcare, effect of price on demand, return-on-investment
- Assume conditional exogeneity

$$Y^{(d)} \perp\!\!\!\perp D \mid X$$

- Assume partially linear response, for a known feature map  $\phi$ 
$$Y = \theta_0' \phi(D, X) + f_0(X) + \epsilon, \quad E[\epsilon \mid D, X] = 0$$
- Parameter of interest  $\theta_0$
- **Example:** in pricing applications  $\phi(D, X) = (D, D^2, X \cdot D, X \cdot D^2)$

# Orthogonal Method: (Extended) Double ML

- **Double ML.** Split samples in half

- Regress  $Y \sim X$  with ML on first half, to get estimate  $\hat{h}(S)$  of  $E[Y|X]$
- Regress  $\phi(D, X) \sim X$  with ML on first half, to get estimate  $\hat{p}(S)$  of  $E[\phi(D, X)|X]$
- Construct residuals on other half,  $\tilde{D} := \phi(D, X) - \hat{p}(X)$  and  $\tilde{Y} := Y - \hat{h}(X)$
- Run OLS on residuals:  $\tilde{Y} \sim \tilde{D}$  to get  $\hat{\theta}$

- Final OLS, in population limit, equivalent to solving normal equation:

$$E[(\tilde{Y} - \theta' \tilde{D}) \tilde{D}] = 0$$

- Define the *formula*:

$$M(\theta, h, p) = E \left[ \left( Y - h(X) - \theta' (\phi(D, X) - p(X)) \right) (\phi(D, X) - p(X)) \right]$$

- OLS is equivalent to solving with respect to  $\theta$ :

$$M(\theta, h_0, p_0) = 0$$

# Asymptotic Normality of DoubleML Estimate

$$E_n[(\hat{Y} - \theta \hat{D})\hat{D}] = 0 \Leftrightarrow \hat{\theta} = E_n[\hat{D}\hat{D}']^{-1} E_n[\hat{D}\hat{Y}]$$

- Assume that  $E[\tilde{D}\tilde{D}'] \succcurlyeq 0$  (minimum eigenvalue bounded away)
- Assume  $\hat{h}, \hat{p}$  estimated on separate sample (or cross-fitting), are consistent and:

$$\sqrt{n} \left( \text{RMSE}(\hat{h}) \cdot \text{RMSE}(\hat{p}) + \text{RMSE}(\hat{p})^2 \right) \rightarrow_p 0$$

- Assume random variables  $Y, D, X$  have bounded fourth moments

$$\sqrt{n}(\hat{\theta} - \theta_0) \rightarrow_d N(0, V), \quad \hat{V} := E_n[\hat{D}\hat{D}']^{-1} E_n[(\hat{Y} - \hat{\theta}\hat{D})^2 \hat{D}\hat{D}'] E_n[\hat{D}\hat{D}']^{-1}$$

*Sandwich formula* for heteroskedasticity-robust  
standard errors in OLS packages (HCO)

General Theory

# Estimation from Moment Restrictions

- Observe samples  $Z_1, \dots, Z_n$  i.i.d. from data distribution  $D$
- Parameter of interest  $\theta_0 \in \mathbb{R}^d$  is identified as the solution to a set of *population formulas*

$$M(\theta_0; g_0) := E_{Z \sim D}[m(Z; \theta_0, g_0)] = 0$$

*Vector of moment restrictions* that distribution  $D$  needs to satisfy and which have a unique solution with respect to  $\theta$

- $g_0 \in G$  a function we don't care (*nuisance function*) but need to estimate from data

Examples.

$$m(Z; \theta, g, a) = g(1, X) - g(0, X) - a(D, X)(Y - g(D, X)) - \theta \quad (\text{ATE for binary treatment})$$

$$m(Z; \theta, h, p) = (Y - h(X) - \theta(D - p(X))) \cdot (D - p(X)) \quad (\text{ATE under PLR model})$$



# Cross-fitting Estimation Algorithm

Split the data in half

- On first half, estimate  $\hat{g}_1$  of  $g_0$  and predict on second half
- On second half, estimate  $\hat{g}_2$  of  $g_0$  and predict on first half

On all data, solution  $\hat{\theta}$  to empirical plug-in moment equation:

$$M_n(\hat{\theta}, \hat{g}) := \frac{1}{n} \sum_{i \in S_1} m(Z_i; \hat{\theta}, \hat{g}_2) + \frac{1}{n} \sum_{i \in S_2} m(Z_i; \hat{\theta}, \hat{g}_1) = 0$$

In practice use  $K \approx 3$  to 5 folds: for each fold, train on all other folds and predict on that fold

# Neyman Orthogonality (Insensitivity)

Moment  $M(\theta, g)$  is **Neyman orthogonal** if for any  $v \in G - g_0$ :

$$\left. \frac{\partial}{\partial t} M(\theta_0, g_0 + t v) \right|_{t=0} = 0$$

# Main Theorem

If moment is Neyman orthogonal and RMSE of  $\hat{g}$  is  $o_p(n^{-1/4})$ , plus regularity conditions

$$\sqrt{n} (\hat{\theta} - \theta_0) \rightarrow N \left( 0, J_0^{-1} \Sigma (J_0^{-1})^\top \right)$$

where  $J_0 := \nabla_\theta M(\theta_0, g_0)$  and  $\Sigma := E[m(Z; \theta_0, g_0) m(Z; \theta_0, g_0)^\top]$

# Practical Variants of Cross-Fitting

# Stacking and Model Selection

- If we want to choose among many models or perform stacking, we can just use a stacked or automl model in place of each ML model

# Stacking ML Models

```
def dml2(X, D, y): # orthogonal dml with sample-splitting
    est_y = StackingRegressor([rf, nnet, gbf, lasso])
    yres = y - cross_val_predict(est_y, X, y, cv=3)
    est_t = StackingRegressor([rf, nnet, gbf, lasso])
    Dres = D - cross_val_predict(est_t, X, D, cv=3)
    theta = np.mean(yres * Dres) / np.mean(Dres**2)
    var = np.mean((Dres**2) * (yres - theta*Dres)**2) / np.mean(Dres**2)
    stderr = np.sqrt(var / X.shape[0])
    return theta, stderr
```

# AutoML Models

```
from flaml import AutoML

def dml2(X, D, y): # orthogonal dml with sample-splitting
    est_y = AutoML()
    yres = y - cross_val_predict(est_y, X, y, cv=3)
    est_t = AutoML()
    Dres = D - cross_val_predict(est_t, X, D, cv=3)
    theta = np.mean(yres * Dres) / np.mean(Dres**2)
    var = np.mean((Dres**2) * (yres - theta*Dres)**2) / np.mean(Dres**2)
    stderr = np.sqrt(var / X.shape[0])
    return theta, stderr
```

# Stacking and Model Selection

- If we want to choose among many models or perform stacking, we can just use a stacked or automl model in place of each ML model
- Model selection or stacking done many times within each training fold
- Computationally expensive and statistically lossy
- Can we use all the data to at least select among models?



# Semi-Crossfitting Estimation Algorithm

Split the data in half (*in practice  $K$  folds*)

- On first half, estimate  $\hat{g}_1^{(1)}, \dots, \hat{g}_1^{(L)}$  of  $g_0$  and predict on second half
- On second half, estimate  $\hat{g}_2^{(1)}, \dots, \hat{g}_2^{(L)}$  of  $g_0$  and predict on first half
- Choose the model  $\ell \in \{1, \dots, L\}$  that optimizes out-of-sample RMSE

- On all data, solution  $\hat{\theta}$  to empirical plug-in moment equation:

$$M_n(\hat{\theta}, \hat{g}) := \frac{1}{n} \sum_{i \in S_1} m(Z_i; \hat{\theta}, \hat{g}_2^{(\ell)}) + \frac{1}{n} \sum_{i \in S_2} m(Z_i; \hat{\theta}, \hat{g}_1^{(\ell)}) = 0$$

# Semi-Crossfitting

```
def dml2(X, D, y): # orthogonal dml with semi-crossfitting
    # cross val predict with many models
    est_y = [rf, gbf, lasso]
    yres = np.array([y - cross_val_predict(est, X, y, cv=3) for est in est_y])
    est_d = [rf, gbf, lasso]
    Dres = np.array([D - cross_val_predict(est, X, D, cv=3) for est in est_d])
    # select models with best out of fold performance
    best_y = np.argmin(np.mean(yres**2, axis=1))
    best_d = np.argmin(np.mean(Dres**2, axis=1))
    yres = yres[best_y]
    Dres = Dres[best_d]
    # go with their corresponding residuals
    theta = np.mean(yres * Dres) / np.mean(Dres**2)
    var = np.mean((Dres**2) * (yres - theta*Dres)**2) / np.mean(Dres**2)
    stderr = np.sqrt(var / X.shape[0])
    return theta, stderr
```

# Semi-Crossfitting

- If the number of models  $L$  is small, then “spillover” is ok and approach still works. For practical purposes  $L$  should be thought as constant.
- Under further regularity, provably asymptotic normality holds if
$$\sqrt{\log(L)} = o(n^{1/4})$$

# Semi-Crossfitting with Stacking

Split the data in half (*in practice  $K$  folds*)

- On first half, estimate  $\hat{g}_1^{(1)}, \dots, \hat{g}_1^{(L)}$  of  $g_0$  and predict on second half
- On second half, estimate  $\hat{g}_2^{(1)}, \dots, \hat{g}_2^{(L)}$  of  $g_0$  and predict on first half
- Construct weights  $\alpha = (\alpha_1, \dots, \alpha_L)$  on the models using all the data (stacking)
- Define stacked models  $\hat{g}_k^{(\alpha)} = \sum_{j=1}^L \alpha_j \cdot \hat{g}_k^{(j)}$
- On all data, solution  $\hat{\theta}$  to empirical plug-in moment equation:

$$M_n(\hat{\theta}, \hat{g}) := \frac{1}{n} \sum_{i \in S_1} m(Z_i; \hat{\theta}, \hat{g}_2^{(\alpha)}) + \frac{1}{n} \sum_{i \in S_2} m(Z_i; \hat{\theta}, \hat{g}_1^{(\alpha)}) = 0$$

# Semi-Crossfitting with Stacking

```
def dml2(X, D, y): # orthogonal dml with semi-crossfitting and stacking
    # cross val predict with many models
    est_y = [rf, gbf, lasso]
    ypreds = np.array([cross_val_predict(est, X, y, cv=3) for est in est_y]).T
    est_d = [rf, gbf, lasso]
    Dpreds = np.array([cross_val_predict(est, X, D, cv=3) for est in est_d]).T
    # calculate stacked residuals by finding optimal coefficients
    # and weighing out-of-sample predictions by these coefficients
    yres = y - LinearRegression().fit(ypreds, y).predict(ypreds)
    Dres = D - LinearRegression().fit(Dpreds, D).predict(Dpreds)
    # go with the stacked residuals
    theta = np.mean(yres * Dres) / np.mean(Dres**2)
    var = np.mean((Dres**2) * (yres - theta*Dres)**2) / np.mean(Dres**2)
    stderr = np.sqrt(var / X.shape[0])
    return theta, stderr
```

# Semi-Crossfitting

- If the number of models  $L$  is small, then “spillover” is ok and approach still works. For practical purposes  $L$  should be thought as constant.
- Under further regularity, provably asymptotic normality holds if
$$\sqrt{L} = o(n^{1/4})$$

Equivalent view of cross-fitting with stacking (lens of FWL theorem)

- Construct out of fold predictions based on many ML models
- Use these predictions as engineered features  $X$  in a simple OLS regression on  $D, X$
- Use the coefficient and standard error of  $D$  from this final OLS

# Appendix: General Theory (Expanded)

# Main Theorem (expanded) Define RMSE: $\|h\|_{L^2} = \sqrt{E[h(X)^2]}$

- If moment is **Neyman orthogonal** and RMSE of  $\hat{g}$  goes down at rate  $n^{1/4}$ , plus regularity conditions

$$n^{1/4} \|\hat{g} - g_0\|_{L^2} \approx 0$$

Jacobian of moments with respect to parameter; relates to identification strength

- Then the estimate  $\hat{\theta}$  is **asymptotically linear**

$$\sqrt{n}(\hat{\theta} - \theta_0) \approx \sqrt{n} E_n[\phi_0(Z)], \quad \phi_0(Z) = -J_0^{-1} m(Z; \theta_0, g_0), \quad J_0 := \partial_{\theta} E[m(Z; \theta_0, g_0)]$$

influence function

influence is a linear transformation of the moment; transforming from

- Consequently, it is **asymptotically normal**

$$\sqrt{n}(\hat{\theta} - \theta_0) \sim_a N(0, V), \quad V = E[\phi_0(Z)\phi_0(Z)']$$

Covariance of the influence function

- Confidence intervals** for any projection based on estimate of variance are asymptotically valid

$$\ell' \theta \in \left[ \ell' \hat{\theta} \pm c \sqrt{\frac{\ell' \hat{V} \ell}{n}} \right],$$

$$\hat{V} = \text{Var}_n(\hat{\phi}(Z)),$$

Empirical variance of approximate influence

$$\hat{\phi}(Z) := -\hat{J}^{-1} m(Z; \hat{\theta}, \hat{g}),$$

Approximate influence function

$$\hat{J} = \partial_{\theta} E_n[m(Z; \hat{\theta}, \hat{g})]$$

Empirical average of jacobian



# Python Pseudocode

```
# General DML pseudocode
```

```
def general_dml(Z, ell, nfolds, moment, jacobian, nuisance
```

```
    # construct out-of-fold predictions from the nuisance
```

```
    ghat = cross_val_predict(nuisance_estimator, Z, cv=nfold
```

```
    # use these predictions to define the empirical moment
```

```
    with respect to theta
```

```
    avg_moment = lambda theta: np.mean(moment(Z, theta, ghat), axis=0)
```

```
    # solve for the empirical moment equation equals zero
```

```
    thetahat = fsolve(avg_moment)
```

```
    # calculate empirical jacobian with respect to theta, evaluated
```

```
    # at the estimates thetahat and ghat
```

```
    Jhat = np.mean(jacobian(Z, thetahat, ghat), axis=0)
```

```
    # construct approximate influence function for each sample
```

```
    phihat = - moment(Z, thetahat, ghat) @ np.linalg.pinv(Jhat).T
```

```
    # variance estimate
```

```
    var = phihat.T @ phihat / phihat.shape[0]
```

```
    # estimate and standard error for any projection ell of the parameters
```

```
    point = ell @ thetahat
```

```
    stderr_ell = np.sqrt(ell.T @ var @ ell / phihat.shape[0])
```

```
    return point, stderr_ell
```

Estimate  $\hat{g}$  out of fold and predict

Construct cross-fitted moment function  
 $\theta \rightarrow E_n[m(Z; \theta, \hat{g})]$

Solve that  
moment = 0  
wrt  $\theta$

Calculate  
 $\hat{J} := \partial_{\theta} E_n[m(Z; \hat{\theta}, \hat{g})]$

Approximate influence  
function  
 $Z \rightarrow \hat{\phi}(Z) := \hat{J}^{-1} m(Z; \hat{\theta}, \hat{g})$

Empirical covariance of  
estimate  
 $\hat{V} = E_n[\hat{\phi}(Z) \hat{\phi}(Z)']$

# Main Theorem (linear moments)

- If moments are linear

$$m(Z; \theta, g) = v(Z; g) - a(Z; g)\theta$$

- Estimate is closed form:

$$\hat{\theta} = \hat{J}^{-1} E_n[v(Z; g)], \quad \hat{J} = E_n[a(Z; g)]$$

- Then the estimate  $\hat{\theta}$  is *asymptotically linear*

$$\sqrt{n}(\hat{\theta} - \theta_0) \approx \sqrt{n} E_n[\phi_0(Z)], \quad \phi_0(Z) = -J_0^{-1} m(Z; \theta_0, g_0), \quad J_0 := E[a(Z; g_0)]$$

- Consequently, it is *asymptotically normal*

$$\sqrt{n}(\hat{\theta} - \theta_0) \sim_a N(0, V), \quad V := E[\phi_0(Z)\phi_0(Z)']$$

- *Confidence intervals* for any projection based on estimate of variance are asymptotically valid

$$\ell' \theta \in \left[ \ell' \hat{\theta} \pm c \sqrt{\frac{\ell' \hat{V} \ell}{n}} \right], \quad \hat{V} = \text{Var}_n(\hat{\phi}(Z)), \quad \hat{\phi}(Z) := -\hat{J}^{-1} m(Z; \hat{\theta}, \hat{g}), \quad \hat{J} = E_n[a(Z; \hat{g})]$$

# Python Pseudocode

```
# General DML pseudocode for linear moments: m(Z; theta, g)
def general_dml_linear_moment(Z, ell, nfolds, alpha, nu, jacob):
    # construct out-of-fold predictions from the nuisance estimator
    ghat = cross_val_predict(nuisance_estimator, Z, cv=nfolds)
    # use these predictions to define the empirical moment equation
    # and solve explicitly with respect to theta
    Jhat = np.mean(alpha(Z, ghat), axis=0)
    avg_nu = np.mean(nu(Z, ghat), axis=0)
    # solve for the empirical moment equation equals zero
    invJhat = np.linalg.pinv(Jhat)
    thetahat = invJhat @ avg_nu
    # construct approximate influence function for each sample
    phihat = - (nu(Z, ghat) - alpha(Z, ghat) @ thetahat) @ invJhat.T
    # variance estimate
    var = phihat.T @ phihat / phihat.shape[0]
    # estimate and standard error for any projection ell of the parameter
    point = ell @ thetahat
    stderr_ell = np.sqrt(ell.T @ var @ ell / phihat.shape[0])
    return point, stderr_ell
```

Estimate  $\hat{g}$  out of fold and predict

cross-fitted jacobian  $\hat{J} := E_n[a(Z; \hat{g})]$

cross-fitted offset  $E_n[v(Z; \hat{g})]$

Closed form solution:  $\theta = \hat{J}^{-1} E_n[v(Z; \hat{g})]$

Approximate influence function  
 $Z \rightarrow \hat{\phi}(Z) := \hat{J}^{-1} m(Z; \hat{\theta}, \hat{g})$

Empirical covariance of estimate  
 $\hat{V} = E_n[\hat{\phi}(Z) \hat{\phi}(Z)']$

# Proving the Main Theorem

# Linear in $\theta$ Moments

- We will restrict attention to a broad class that simplifies proof
- Moment is linear in target parameter

$$m(Z; \theta, g) = a(Z; g)' \theta + v(Z; g)$$


- Expected moment also linear in  $\theta$

$$M(\theta, g) = A(g)' \theta + V(g)$$

# Proof Ingredients: Linear in $\theta$ Moments

- Since  $M_n(\hat{\theta}, \hat{g}) = 0$  we expect by concentration and sample splitting  $M(\hat{\theta}, \hat{g}) \approx n^{-1/2}$
- Since  $M(\theta_0, g_0) = 0$  we expect by Neyman orthogonality  $M(\theta_0, \hat{g}) \approx RMSE(\hat{g})^2 = o(n^{-1/2})$
- Since moment is linear in  $\theta$ :  $A(\hat{g})(\hat{\theta} - \theta_0) = M(\hat{\theta}, \hat{g}) - M(\theta_0, \hat{g})$
- Since  $A$  is Lipschitz and  $\hat{g} \rightarrow g_0$ :  $A(g_0)(\hat{\theta} - \theta_0) = M(\hat{\theta}, \hat{g}) - M(\theta_0, \hat{g}) + o_p(\|\hat{\theta} - \theta_0\|)$
- Since  $A(g_0)$  is invertible:  $\|\hat{\theta} - \theta_0\| = O(\|M(\hat{\theta}, \hat{g})\| + \|M(\theta_0, \hat{g})\|) = O_p(n^{-1/2})$
- More fine-grained analysis of  $M(\hat{\theta}, \hat{g})$  term, shows:  $\sqrt{n}M(\hat{\theta}, \hat{g}) \rightarrow N(0, V)$

# Proof of Main Theorem (visually)



# Proof of Main Theorem (algebraically)

- Since moment  $M(\theta, g)$  is linear in  $\theta$  and  $M_n(\hat{\theta}, \hat{g}) = 0$  and  $M(\theta_0, g_0) = 0$   

$$A(\hat{g}) (\hat{\theta} - \theta_0) = M(\hat{\theta}, \hat{g}) - M(\theta_0, \hat{g})$$

$$= M(\hat{\theta}, \hat{g}) - M_n(\hat{\theta}, \hat{g}) + M(\theta_0, g_0) - M(\theta_0, \hat{g})$$

- Since  $\text{RMSE}(\hat{g}) = \|\hat{g} - g_0\| = o_p(1)$

$$A(g_0) (\hat{\theta} - \theta_0) = A(\hat{g}) (\hat{\theta} - \theta_0) + o_p(\|\hat{\theta} - \theta_0\|)$$

- Thus

$$A(g_0) (\hat{\theta} - \theta_0) = \underbrace{M(\hat{\theta}, \hat{g}) - M_n(\hat{\theta}, \hat{g})}_{\rightarrow N(0, V)} + \underbrace{M(\theta_0, g_0) - M(\theta_0, \hat{g})}_{= o_p(n^{-1/2})} + o_p(\|\hat{\theta} - \theta_0\|)$$

$\rightarrow N(0, V)$   
 via CLT  
 + sample-splitting  
 + concentration  
 +  $\hat{g} \rightarrow g_0$

$= o_p(n^{-1/2})$   
 via orthogonality  
 +  $\|\hat{g} - g_0\| = o_p(n^{-1/4})$



# Proof of Main Theorem: Orthogonality

- By Neyman orthogonality and bounded second derivative of  $M(\theta_0, g)$  w.r.t.  $g$   
 $M(\theta_0, g_0) - M(\theta_0, \hat{g}) = D_g M(\theta_0, g_0)[g_0 - g] + O(\|\hat{g} - g_0\|^2) = o_p(n^{-1/2})$

- Thus

$$A(g_0) (\hat{\theta} - \theta_0) = \underbrace{M(\hat{\theta}, \hat{g}) - M_n(\hat{\theta}, \hat{g})}_{G_n(\hat{\theta}, \hat{g})} + o_p(n^{-1/2} + \|\hat{\theta} - \theta_0\|)$$

$G_n(\hat{\theta}, \hat{g})$

# Proof of Main Theorem: Sample-Splitting (1)

- Let  $G_n(\theta, g) = M(\theta, g) - M_n(\theta, g)$   
$$G_n(\hat{\theta}, \hat{g}) = G_n(\theta_0, g_0) + G_n(\hat{\theta}, \hat{g}) - G_n(\theta_0, \hat{g}) + G_n(\theta_0, \hat{g}) - G_n(\theta_0, g_0)$$
- Linearity of moment + (sample-splitting and concentration  $\Rightarrow \|A(\hat{g}) - A_n(\hat{g})\| = o_p(1)$ ):  
$$G_n(\hat{\theta}, \hat{g}) - G_n(\theta_0, \hat{g}) = (A(\hat{g}) - A_n(\hat{g})) (\hat{\theta} - \theta_0) = o_p(\|\hat{\theta} - \theta_0\|)$$
- Thus  
$$A(g_0) (\hat{\theta} - \theta_0) = G_n(\theta_0, g_0) + G_n(\theta_0, \hat{g}) - G_n(\theta_0, g_0) + o_p(n^{-1/2} + \|\hat{\theta} - \theta_0\|)$$

# Proof of Main Theorem: Sample-Splitting (2)

- Note for  $X_i = m(Z_i; \theta_0, \hat{g}) - m(Z_i; \theta_0, \hat{g}) - E[m(Z_i; \theta_0, \hat{g}) - m(Z_i; \theta_0, \hat{g})]$

$$G_n(\theta_0, \hat{g}) - G_n(\theta_0, g_0) = \frac{1}{n_2} \sum_{i \in S_2} X_i$$

- By sample splitting,  $X_i$  are i.i.d. with  $E[X_i] = 0$ . By variance decomposition (concentration)

$$\left\| \frac{1}{n_2} \sum_{i \in S_2} X_i \right\|_{L_2} \leq \sqrt{\frac{E[X_i^2]}{n}}$$

- Thus

$$\|G_n(\theta_0, \hat{g}) - G_n(\theta_0, g_0)\|_{L_2} \leq \frac{1}{\sqrt{n}} \sqrt{E \left[ \left( m(Z_i; \theta_0, \hat{g}) - m(Z_i; \theta_0, \hat{g}) \right)^2 \right]} = O \left( \frac{\|\hat{g} - g_0\|}{\sqrt{n}} \right) = o_p(n^{-1/2})$$

# Concluding

- So far

$$A(g_0) (\hat{\theta} - \theta_0) = G_n(\theta_0, g_0) + o_p(n^{-1/2} + \|\hat{\theta} - \theta_0\|)$$

- Since  $A(g_0)$  is invertible and  $G_n(\theta_0, g_0) = O_p(n^{-1/2})$  by concentration  
 $\|\hat{\theta} - \theta_0\| = O_p(n^{-1/2})$

- Thus, we have asymptotic linearity

$$\sqrt{n} (\hat{\theta} - \theta_0) = \sqrt{n} A(g_0)^{-1} G_n(\theta_0, g_0) + o_p(1) = \frac{1}{\sqrt{n_2}} \sum_{i \in S_2} A(g_0)^{-1} m(Z_i; \theta_0, g_0) + o_p(1)$$

- By CLT we get the theorem

# Main Theorem

- If moment is Neyman orthogonal and RMSE of  $\hat{g}$  is  $o_p(n^{-1/4})$  \*

$$\sqrt{n} (\hat{\theta} - \theta_0) \rightarrow N \left( 0, A^{-1} \Sigma (A^{-1})^\top \right)$$

- $A = \nabla_{\theta} M(\theta_0, g_0)$  and  $\Sigma = E[m(Z; \theta_0, g_0) m(Z; \theta_0, g_0)^\top]$

\*plus regularity conditions