

MS&E 233

Game Theory, Data Science and AI

Lecture 13

Vasilis Syrgkanis

Assistant Professor

Management Science and Engineering

(by courtesy) Computer Science and Electrical Engineering

Institute for Computational and Mathematical Engineering

Computational Game Theory for Complex Games

- 1
 - Basics of game theory and zero-sum games (T)
 - Basics of online learning theory (T)
 - Solving zero-sum games via online learning (T)
 - *HW1: implement simple algorithms to solve zero-sum games*
 - Applications to ML and AI (T+A)
 - *HW2: implement boosting as solving a zero-sum game*

- 2
 - Basics of extensive-form games
 - Solving extensive-form games via online learning (T)
 - *HW3: implement agents to solve very simple variants of poker*

- 3
 - General games, equilibria and online learning (T)
 - Online learning in general games
 - *HW4: implement no-regret algorithms that converge to correlated equilibria in general games*

Data Science for Auctions and Mechanisms

- 4
 - Basics and applications of auction theory (T+A)
 - Basic Auctions and Learning to bid in auctions (T)
 - *HW5: implement bandit algorithms to bid in ad auctions*

- 5
 - Optimal auctions and mechanisms (T)
 - **Simple vs optimal mechanisms (T)**
 - *HW6: implement simple and optimal auctions, analyze revenue empirically*

- 6
 - Basics of Statistical Learning Theory (T)
 - Optimizing Mechanisms from Samples (T)
 - *HW7: implement procedures to learn approximately optimal auctions from historical samples and in an online manner*

Further Topics

- 7
 - Econometrics in games and auctions (T+A)
 - A/B testing in markets (T+A)
 - *HW8: implement procedure to estimate values from bids in an auction, empirically analyze inaccuracy of A/B tests in markets*

Guest Lectures

- Mechanism Design and LLMs, Song Zuo, Google Research
- A/B testing in auction markets, Okke Schrijvers, Central Applied Science, Meta

Summarizing Last Lecture

Myerson's Theorem. When valuations are independently distributed, for any BIC, NNT and IR mechanism (and any BNE of a non-truthful mechanism), the payment contribution of each player is their expected virtual value

$$E[\hat{p}_i(v_i)] = E[\hat{x}_i(v_i) \cdot \phi_i(v_i)], \quad \phi_i(v_i) = v_i - \frac{1 - F_i(v_i)}{f_i(v_i)}$$

Corollary. When valuations are independently distributed, for any Bayes-Nash equilibrium of any non-truthful mechanism, the payment contribution of each player is their expected virtual value

$$E[\hat{p}_i(v)] = E[\hat{x}_i(v_i) \cdot \phi_i(v_i)], \quad \phi_i(v_i) = v_i - \frac{1 - F_i(v_i)}{f_i(v_i)}$$

Myerson's Optimal Auction. Assuming that virtual value functions are monotone non-decreasing, the mechanism that maximizes virtual welfare, achieves the largest possible revenue among all possible mechanisms and Bayes-Nash

$$x(v) = \operatorname{argmax}_{x \in X} \sum_i x \cdot \phi_i(v_i), \quad p_i(v) = v_i x_i(v) - \int_0^{v_i} x_i(z, v_{-i}) dz$$

$$\text{Rev} = E \left[\max_{x \in X} \sum_i x \cdot \phi_i(v_i) \right]$$

Dissecting Myerson's Optimal Auction

Identically Distributed Bidders

- Single-item setting, with all bidder values are from same distribution $v_i \sim F$
- Virtual value function is the same for all bidders

$$\phi(v) = v - \frac{1 - F(v)}{f(v)}$$

- Assume that $\phi(v)$ is monotone non-decreasing (F is regular)
- Allocating to highest virtual value \equiv allocating to highest value
- **Optimal auction.** Allocate to highest value, as long as $\phi(v_{(1)}) \geq 0$
- **Optimal auction.** allocate to highest value, as long as $v_1 \geq r_*$

$$r_*: r - \frac{1 - F(r)}{f(r)} = 0, \quad (\text{monopoly reserve price})$$

When bidders are independently and identically distributed according to a regular distribution, then the optimal single-item auction among all auctions is a Second-Price Auction with a Monopoly Reserve Price

Monopoly Reserve Price

- What if we had only one bidder (monopoly)
- Then optimal thing to do is post a reserve price r_*
- The revenue from that single bidder if we post a reserve r is

$$E[r \mathbf{1}\{v \geq r\}] = r (1 - F(r))$$

- The optimal reserve price is given by the first order condition

$$r_*: (1 - F(r)) - r f(r) = 0 \Rightarrow r - \frac{1 - F(r)}{f(r)} = 0$$

- Same as reserve price that we should be using with many bidders

Non-Identically Distributed Bidders

- What if you know ahead of time that one bidder tends to have higher values than the other bidder?
- Shouldn't you treat these bidders differently (price discrimination)?
- Shouldn't you try to extract more revenue from the bidder that tends to have a higher value?

Google digital advertising

All Images News Videos Shopping More

About 6,620,000,000 results (0.44 seconds)

Sponsored

Reddit
https://www.redditforbusiness.com

Advertise on Reddit

Reach over 100K communities — Connect with passionate communities that deliver results for brands across all industries. Create impact & own top communities in your target category for 24 hours. Try Reddit ads.

1

Reddit
https://www.redditforbusiness.com
Advertise on Reddit

$$v_1 \sim U[0,1]$$

Sponsored

Microsoft
https://about.ads.microsoft.com › advertising › start-now

Microsoft Advertising® | Get a \$500 Advertising Credit

$$v_2 \sim U[0,100]$$

You are selling a single item to two bidders. One has values drawn $U[0,1]$ the other $U[0,100]$. What is the optimal auction?

Second-price with a reserve price

Second-price where each bidder has a different reserve price

First-price where each bidder has a different reserve price

None of the above

You are selling a single item to two bidders. One has values drawn $U[0,1]$ the other $U[0,100]$. What is the optimal auction?

Second-price with a reserve price

0%

Second-price where each bidder has a different reserve price

0%

First-price where each bidder has a different reserve price

0%

None of the above

0%

You are selling a single item to two bidders. One has values drawn $U[0,1]$ the other $U[0,100]$. What is the optimal auction?

Second-price with a reserve price

0%

Second-price where each bidder has a different reserve price

0%

First-price where each bidder has a different reserve price

0%

None of the above

0%

Non-Identically Distributed Bidders

- Suppose we have two bidders, $v_1 \sim U[0,1]$, $v_2 \sim U[0,100]$
- Virtual value function for each bidder
- We should allocate to the bidder with the highest virtual value (if positive)!

Non-Identically Distributed Bidders

- Suppose we have two bidders, $v_1 \sim U[0,1]$, $v_2 \sim U[0,100]$

- Virtual value function for each bidder

$$\phi_1(v) = v - \frac{1 - F_1(v)}{f_1(v)} = 2v - 1, \quad \phi_2(v) = v - \frac{1 - \frac{v}{100}}{\frac{1}{100}} = 2v - 100$$

- We should allocate to the bidder with the highest virtual value (if positive)!

$$\operatorname{argmax}\{0, \phi_1(v_1), \phi_2(v_2)\} = \operatorname{argmax}\{0, 2v_1 - 1, 2v_2 - 100\}$$

Non-Identically Distributed Bidders

- Allocate to highest virtual value (if positive)!

$$\operatorname{argmax}\{0, 2v_1 - 1, 2v_2 - 100\}$$

Google


digital advertising

All Images News Videos Shopping More

About 6,620,000,000 results (0.44 seconds)

Sponsored

1



$$v_1 \sim U[0,1]$$



$$v_1 =$$



$$\phi_1 =$$

Sponsored



Microsoft

<https://about.ads.microsoft.com> > advertising > start-now

Microsoft Advertising® | Get a \$500 Advertising Credit

$$v_2 \sim U[0,100]$$



$$v_2 =$$



$$\phi_2 =$$

Non-Identically Distributed Bidders

- Allocate to highest virtual value (if positive)!

$$\operatorname{argmax}\{0, 2v_1 - 1, 2v_2 - 100\}$$

Google


digital advertising

All Images News Videos Shopping More

About 6,620,000,000 results (0.44 seconds)

Sponsored

1



$$v_1 \sim U[0,1]$$



$$v_1 = 1$$



$$\phi_1 = 1$$

Sponsored



Microsoft
<https://about.ads.microsoft.com> > advertising > start-now

Microsoft Advertising® | Get a \$500 Advertising Credit

$$v_2 \sim U[0,100]$$



$$v_2 = 20$$



$$\phi_2 = -60$$

Non-Identically Distributed Bidders

- Allocate to highest virtual value (if positive)!

$$\operatorname{argmax}\{0, 2v_1 - 1, 2v_2 - 100\}$$

Google

digital advertising

All Images News Videos Shopping More

About 6,620,000,000 results (0.44 seconds)

Sponsored



Reddit

<https://www.redditforbusiness.com>

Advertise on Reddit

1



Reddit

<https://www.redditforbusiness.com>

Advertise on Reddit

$$v_1 \sim U[0,1]$$



$$v_1 = 1$$



$$\phi_1 = 1$$

Sponsored



Microsoft

<https://about.ads.microsoft.com> > advertising > start-now

Microsoft Advertising® | Get a \$500 Advertising Credit

$$v_2 \sim U[0,100]$$



$$v_2 = 20$$



$$\phi_2 = -60$$

Non-Identically Distributed Bidders

- Allocate to highest virtual value (if positive)!

$$\operatorname{argmax}\{0, 2v_1 - 1, 2v_2 - 100\}$$

Google

digital advertising

All Images News Videos Shopping More

About 6,620,000,000 results (0.44 seconds)


Sponsored

Sponsored

Microsoft
<https://about.ads.microsoft.com> > advertising > start-now

Microsoft Advertising® | Get a \$500 Advertising Credit

1



$$v_1 \sim U[0,1]$$



$$v_1 = 1$$



$$\phi_1 = 1$$

Sponsored



Microsoft
<https://about.ads.microsoft.com> > advertising > start-now

Microsoft Advertising® | Get a \$500 Advertising Credit

$$v_2 \sim U[0,100]$$



$$v_2 = 51$$



$$\phi_2 = 2$$

Non-Identically Distributed Bidders

- Allocate to highest virtual value (if positive)!

$$\operatorname{argmax}\{0, 2v_1 - 1, 2v_2 - 100\}$$

Google

digital advertising


All Images News Videos Shopping More

About 6,620,000,000 results (0.44 seconds)

Sponsored



1




$$v_1 \sim U[0,1]$$



$$v_1 = .49$$



$$\phi_1 = -.02$$



$$v_2 \sim U[0,100]$$



$$v_2 = 49$$





$$\phi_2 = -2$$

Non-Identically Distributed Bidders

- Allocate to highest virtual value (if positive)!

$$\operatorname{argmax}\{0, 2v_1 - 1, 2v_2 - 100\}$$

Bidder 1 wins if:




Non-Identically Distributed Bidders

- Allocate to highest virtual value (if positive)!

$$\operatorname{argmax}\{0, 2v_1 - 1, 2v_2 - 100\}$$

Bidder 1 wins if: $2v_1 - 1 \geq 2v_2 - 100 \Rightarrow v_1 \geq v_2 - \frac{99}{2}$




Non-Identically Distributed Bidders

- Allocate to highest virtual value (if positive)!

$$\operatorname{argmax}\{0, 2v_1 - 1, 2v_2 - 100\}$$

Optimal auction rules




Non-Identically Distributed Bidders

- Allocate to highest virtual value (if positive)!

$$\operatorname{argmax}\{0, 2v_1 - 1, 2v_2 - 100\}$$

Optimal auction rules

- If $v_1 > .5$, $v_2 < 50$, allocate to 1, charge .5
- If $v_1 < .5$, $v_2 > 50$, allocate to 2, charge 50
- If $.5 \leq v_1 < v_2 - 49.5$, allocate to 2, charge $v_1 + 49.5$
- If $50 \leq v_2 < v_1 + 49.5$, allocate to 1, charge $v_2 - 49.5$



At the optimal auction, we are giving a huge advantage to the weaker bidder! We roughly add 49.5\$ to their bid!

We expect more from stronger bidders and make it harder for them to win, to incentivize them to pay more.

Optimal auction is

1) cumbersome, 2) hard to understand, 3) hard to explain, 4) does not always allocate to the highest value player, 5) discriminates a lot, 6) is many times counter-intuitive, 7) can seem unfair!

Are there simpler auctions that
always achieve almost as good
revenue?

Simple vs. Optimal Auctions

Second-Price with Player-Specific Reserves

- What if we simply run a second price auction but have different reserves for each bidder
- Each bidder i has a reserve price r_i
- Reject all bidders with bid below the reserve
- Among all bidders with value $v_i \geq r_i$, allocate to highest bidder
- Charge winner max of their reserve and the next highest surviving bid

Theorem. There exist personalized reserve prices such that the above auction achieves at least $\frac{1}{2}$ of the optimal auction revenue!

Second-Price with Player-Specific Reserves

Theorem. There exist personalized reserve prices such that the above auction achieves at least $\frac{1}{2}$ of the optimal auction revenue!

- Revenue of the optimal auction is the maximum virtual welfare

$$\text{OPT} = E \left[\max_i \phi_i^+(v_i) \right], \quad \phi_i^+(v_i) = \max\{0, \phi_i(v_i)\}$$

- Assume that reserve prices are at least the monopoly reserves
- Revenue of the second-price with player specific reserves (SP-r)

$$\text{Rev} = E \left[\sum_i x_i(v) \phi_i^+(v_i) \right]$$

- Can we guarantee that the auction collects a $\phi_i^+(v_i)$ that, in expectation, is at least half of the maximum $\phi_i^+(v_i)$?

Second-Price with Player-Specific Reserves

Theorem. There exist personalized reserve prices such that the above auction achieves at least $\frac{1}{2}$ of the optimal auction revenue!

- Can we guarantee that the auction collects a $\phi_i^+(v_i)$ that, in expectation, is at least half of the maximum $\phi_i^+(v_i)$?
- Since the auction allocates to some player with $v_i \geq r_i$
- Since ϕ_i^+ are monotone: to some player with $\phi_i^+(v_i) > \theta_i$
- We can think of $\phi_i^+(v_i)$ as non-negative prizes Π_i

Second-Price with Player-Specific Reserves

Theorem. There exist personalized reserve prices such that the above auction achieves at least $\frac{1}{2}$ of the optimal auction revenue!

- We can think of $\phi_i^+(v_i)$ as non-negative prizes Π_i
- The optimal auction gets revenue that corresponds to the expected maximum prize $E[\max_i \Pi_i]$
- The SP-r auction gets revenue that corresponds to some price Π_τ that satisfies that it is above some threshold θ_τ
- Is there a threshold rule for collecting prizes that guarantees at least half of the expected maximum prize?

Parenthesis: (Optimal Stopping Problems)

- There are n stages
- In each stage i , we are offered a prize $\Pi_i \sim G_i$
- Distributions G_i are known ahead of time
- Realized prize Π_i only revealed at stage i
- At each stage, we can choose to accept Π_i and end the game or discard the prize and continue opening prizes

Question. Is there a strategy to play the game that guarantees at least half of what an oracle who knows all the prizes ahead of time would achieve?

Parenthesis: (Optimal Stopping Problems)

Question. Is there a strategy to play the game that guarantees at least half of what an “prophet” who knows all the prizes ahead of time would achieve?

Theorem (Prophet Inequality). There exists a threshold strategy APX that accepts the first prize that passes a threshold θ , such that:

$$E[\Pi_\tau] \geq \frac{1}{2} E \left[\max_i \Pi_i \right]$$

τ is the random stopping time induced by the threshold policy.

Parenthesis: (Proof of Prophet Inequality)

- Let's be generous with the optimal benchmark

$$E[\Pi_*] = E\left[\max_i \Pi_i\right] \leq E[\theta + [\Pi_* - \theta]_+] \leq \theta + \sum_i \overbrace{E[[\Pi_i - \theta]_+]}^{A_i}$$

- APX gets θ if there exists some prize above, i.e., $\Pi_* \geq \theta$
- On top of that, APX also collects some **excess** $[\Pi_\tau - \theta]_+$
- **Excess** is A_i , when all rewards other than i is $\leq \theta$

$$\text{Excess} \geq \sum_i A_i \Pr(\forall j \neq i: \Pi_j < \theta) \geq \sum_i A_i \Pr(\Pi_* < \theta)$$

Overall: $\text{APX} \geq \theta \Pr(\Pi_* \geq \theta) + \Pr(\Pi_* < \theta) \sum_i A_i$

Choosing $\Pr(\Pi_* \geq \theta) = 1/2$: $\text{APX} \geq \frac{1}{2} (\theta + \sum_i A_i) \geq \frac{1}{2} E[\Pi_*]$

Parenthesis: (Optimal Stopping Problems)

Question. Is there a strategy to play the game that guarantees at least half of what an “prophet” who knows all the prizes ahead of time would achieve?

Theorem (Prophet Inequality). There exists a threshold strategy APX that accepts the first prize that passes a threshold θ , such that:

$$E[\Pi_\tau] \geq \frac{1}{2} E \left[\max_i \Pi_i \right]$$

τ is the random stopping time induced by the threshold policy.

Policy. Simply choose θ such that $\Pr \left(\max_i \Pi_i \geq \theta \right) = 1/2$

Second-Price with Player-Specific Reserves

Theorem. There exist personalized reserve prices such that the above auction achieves at least $1/2$ of the optimal auction revenue!

- Choose θ such that:

$$\Pr\left(\max_i \phi_i^+(v_i) \geq \theta\right) = 1/2$$

- Then set personalized reserve prices implied by:

$$\phi_i^+(v_i) \geq \theta \Leftrightarrow v_i \geq r_i$$

All these designs required knowledge
of distributions of values F_i !

What can we do if we only have
data from F_i ?

Learning Auctions from Samples

Learning from Samples

- We are given a set S of m samples of value profiles

$$S = \left\{ v^j = \left(v_1^j, \dots, v_n^j \right) \right\}_{j=1}^m$$

- Each sample is drawn i.i.d. from the distribution of values

$$v_i^j \sim F_i, \quad v^j \sim \mathbf{F} \stackrel{\text{def}}{=} F_1 \times \dots \times F_n$$

- Samples can be collected from historical runs of truthful auction
- Bids of each bidder in each of the m historical runs of the auction