

Problems for the mini-course “Probability theory”

Kabatyansky G.A., Gasnikov A.V. gasnikov@yandex.ru

Problem 1 (Random walk, 1.5). There is a point on one-dimensional integer lattice \mathbb{Z} . With probability 0.5 the point moves to the right on distance 1 and with probability 0.5 moves to the left on the distance 1. Show that with probability 1 the point will return to the starting point during infinitely time period.

Problem 2 (Hoeffding’s Lemma, 1). Show that for any random variable x with $|x| \leq 1$ and $E[x]=0$ the following inequality is true $E[\exp(tx)] \leq \exp(t^2/2)$.

Problem 3 (Concentration of measure, 1). Consider two unit balls of radius 1 in R^n . Let the center of first ball is at the point $(0, \dots, 0)^T$ and the center of the second ball is at $(1, \dots, 0)^T$. Show that the volume of intersection of these two balls divided by the volume of a unit ball tends to zero with n tending to infinity.

Problem 4 (Concentration of measure, 1). Let a random vector X has a uniform distribution on the unit ball in R^n with center at the point $(0, \dots, 0)^T$. Show that for any $a > 0$ the probability of the event $\|X\|_2 > 1 - a$ tends to 1 with n tending to infinity.

Problem 5 (Concentration of measure, 1). Choose two independent random vectors X and Y with the uniform distribution on the unit ball in R^n with center at the point $(0, \dots, 0)^T$. Proof that with high probability $\langle X, Y \rangle \approx 0$ with n tending to infinity.

Literature

Ball K. An elementary introduction to modern convex geometry // Flavors of geometry. MSRI publication. 1997. V. 31. P. 1–58.

DasGupta A. Asymptotic theory of statistic and probability. Springer, 2008. 687 p.

Levin D.A., Peres Y., Wilmer E.L. Markov chain and mixing times. Providence: AMS, 2009. 387 p.

Durrett R. Probability: Theory and Examples. Cambridge Univ. Press, 2010.