

## PARTIAL DIFFERENTIAL EQUATIONS

### Quiz 3, Time 2 hours

Instructor: James Vickers

Date: Friday 22nd November

---

#### Bonus Question

The forced heat equation

$$\frac{\partial y}{\partial t} - \frac{\partial^2 y}{\partial x^2} = F(x) \quad (1)$$

is defined on the domain  $x \in [-\pi, \pi]$  and for  $t > 0$  and is subject to the Neumann boundary conditions

$$\left. \frac{\partial y}{\partial x} \right|_{x=-\pi} = 0, \quad \text{and} \quad \left. \frac{\partial y}{\partial x} \right|_{x=\pi} = 0. \quad (2)$$

4(a) [15 MARKS] Assuming that the forcing term  $F(x)$  is given by

$$F(x) = \begin{cases} \pi + x & x < 0, \\ \pi - x & x > 0 \end{cases}$$

show that the general solution of the heat equation (1) is

$$y(x, t) = \frac{\pi t}{2} + C_0 + \sum_{n=1}^{\infty} \left[ \frac{2}{\pi n^4} (1 - (-1)^n) + C_n e^{-n^2 t} \right] \cos(nx). \quad (3)$$

4(b) [10 MARKS] If in addition the initial conditions are

$$y(x, 0) = 0,$$

show that the solution can be written as

$$y(x, t) = \frac{\pi t}{2} + \sum_{k=1}^{\infty} \frac{4}{\pi(2k-1)^4} \left( 1 - e^{-(2k-1)^2 t} \right) \cos((2k-1)x).$$

**Solution:**

We immediately note that the problem is reflection symmetric about  $x = 0$  and the boundary conditions are such that, when expressed in the standard Fourier Series like form, we expect a cosine series. Therefore we make the ansatz that

$$y(x, t) = \frac{1}{2}T_0(t) + \sum_{n=1}^{\infty} T_n(t) \cos(nx).$$

In addition we make the ansatz

$$F(x) = \frac{1}{2}F_0 + \sum_{n=1}^{\infty} F_n \cos(nx).$$

We then use the Euler formulas, noting that  $F$  is even and so using the half range series, to compute

$$F_0 = \frac{2}{\pi} \int_0^{\pi} (\pi - x) dx = \pi$$

and (more lengthily, but straightforwardly)

$$F_n = \frac{2}{\pi} \int_0^{\pi} (\pi - x) \cos(nx) dx = \frac{2}{\pi n^2} (1 - (-1)^n).$$

Substituting our ansatzes in to the heat equation gives

$$\frac{1}{2}\dot{T}_0 + \sum \dot{T}_n \cos(nx) = \sum -T_n n^2 \cos(nx) + \frac{\pi}{2} + \sum F_n \cos(nx).$$

Noting that this must hold for all values of  $t$  we note that this is a Fourier series at any fixed value of  $t$ , which allows us to use orthogonality to get

$$\begin{aligned} \dot{T}_0 &= \pi, \\ \dot{T}_n + n^2 T_n &= F_n \end{aligned}$$

which immediately implies (from the linear equation for  $T_n$ ) that

$$\begin{aligned} T_0 &= \pi t + C_0, \\ T_n &= \frac{F_n}{n^2} + C_n e^{-n^2 t} \end{aligned}$$

where the  $C_n$  are integration constants.

Plugging these results back in gives the claimed general solution. We set  $t = 0$  to give

$$0 = 0 + \sum \left[ \frac{2}{\pi n^4} (1 - (-1)^n) + C_n \right] \cos(nx).$$

We can immediately use orthogonality or the uniqueness of Fourier series to note that the term in square brackets must vanish. This implies that  $C_n$  is given by

$$C_n = -\frac{2}{\pi n^4} (1 - (-1)^n)$$

from which it follows that

$$y(t, x) = \frac{\pi t}{2} + \sum_{n=1}^{\infty} \frac{2}{\pi n^4} (1 - (-1)^n) \left( 1 - e^{-n^2 t} \right) \cos(nx).$$

We note that this vanishes for even  $n$ ; setting  $n = 2k - 1$  gives the stated result.