



Minimum Target Age of Child	Type of Vaccine	Dosage	Route of Administration	Site
At Birth	BCG	0.05ml	Intra dermal	Left Upper Arm
	*OPV0	2 drops	Oral	Mouth
	**Hep B0 birth	0.5ml	Intramuscular	Anterolateral aspect of Right thigh
6 Weeks	Pentavalent (DPT, Hep B and Hib) 1	0.5ml	Intramuscular	Anterolateral aspect of Left thigh
	Pneumococcal Conjugate Vaccine1	0.5ml	Intramuscular	Anterolateral aspect of right thigh
	*OPV1	2 drops	Oral	Mouth
	***Rota 1	5 drops	Oral	Mouth
	IPV1	0.5ml	Intramuscular	Anterolateral aspect of right thigh (2.5cm apart from PCV)
10 Weeks	Pentavalent (DPT, Hep B and Hib)2	0.5ml	Intramuscular	Anterolateral aspect of left thigh
	Pneumococcal Conjugate Vaccine2	0.5ml	Intramuscular	Anterolateral aspect of right thigh
	*OPV 2	2 drops	Oral	Mouth
	***Rota 2	5 drops	Oral	Mouth
14 Weeks	Pentavalent 3 (DPT, Hep B and Hib)	0.5ml	Intramuscular	Anterolateral aspect of Left thigh
	Pneumococcal Conjugate Vaccine3	0.5ml	Intramuscular	Anterolateral aspect of right thigh
	*OPV 3	2 drops	Oral	Mouth
	***Rota 3	5 drops	Oral	Mouth
	IPV2	0.5ml	Intramuscular	Anterolateral aspect of right thigh (2.5cm apart from PCV)
6 Months	Vitamin A 1st dose	100,000 IU	Oral	Mouth
9 Months	Measles 1st dose	0.5ml	Subcutaneous	Left upper arm
	Yellow Fever	0.5ml	Subcutaneous	Right upper arm
	Meningitis Vaccine	0.5ml	Intramuscular	Anterolateral aspect of Left thigh
12 Months	Vitamin A 2nd dose	200,000 IU	Oral	Mouth
15 Months	Measles 2nd dose (MCV2)	0.5ml	Subcutaneous	Left upper arm
9-13 years	***HPV 6 months interval (2 doses)	0.5ml	Intramuscular	Deltoid muscle (upper arm)