

Every intervention moved the readout; none moved the behaviour


Base Llama-3.1-8B-Instruct, deceived route, target burn-blister-pop, N = 8 runs per arm. Left: the internal readout each intervention targeted. Right: what the behaviour did.
EXPLORATORY (D-023). One target: the detectable behavioural gap at this N is one run in eight (0.125 of a rate).

the readout moved

no random text arm exists; the band is the random floor of the displacement


self-blame text (vs act-blame)

S4 cells B vs A



the behaviour moved



coherence: not in S4_table.md



injected norm 2.92 in both C and D; C's readout is the injection, not a state change; no CI in table

guilt-arrow steering +4 $\cdot\sigma$ (guilt_clean) at L16



S4 cell C vs B; D random



1.878 $\rightarrow$ 1.018, -46 %; injected norm 0.888 in Q1 and Q3; no CI in table

persona-axis steering -4 $\cdot\sigma$ (persona) at L16



S5c Q1 vs Q0; Q3 random



injected norm 2.65 in P1 and P2 (`into` position: -0.037 $\rightarrow$ -1.583); no CI in table

anti-susceptibility steering -4 $\cdot\sigma$ (nn) at L16, every turn

S5b P1 vs P0; P2 random



○ comparison arm ● intervention arm ● norm-matched random arm ■ random floor (randctl seeds 0-9) — 95 % CI where the table prints one