

DRIVER APPLICATION FORM

COMPANY DETAILS

Company	Hull Ross Associates	Phone	Cameron and Hinton Traders
Email	vubi@mailinator.com	City	Pennington Mays Plc

APPLICANT DETAILS

Full Name	Indigo Swanson	DOB	2003-03-23
Phone	Explicabo Deserunt	Email	kunyhi@mailinator.com
Address	Nisi sequi reprehend		

LICENSE INFORMATION

License No	272	State	Qui unde sunt commo
Issue Date	1971-07-17	Expiry	1989-10-26

DRIVING EXPERIENCE

Equipment	Experience	From	To	Miles
Straight Truck	yes	Sun Nov 23 1975 05:30:00 GMT+0530 (India Standard Time)	Wed Jul 23 1997 05:30:00 GMT+0530 (India Standard Time)	76
Straight Truck	yes	Sun Nov 23 1975 05:30:00 GMT+0530 (India Standard Time)	Wed Jul 23 1997 05:30:00 GMT+0530 (India Standard Time)	76
Truck Tractor	yes	Sat Feb 01 2003 05:30:00 GMT+0530 (India Standard Time)	Thu Apr 22 1993 05:30:00 GMT+0530 (India Standard Time)	28
Semi Trailer	no	-	-	-
Double Triple	yes	Mon Jun 08 1981 05:30:00 GMT+0530 (India Standard Time)	Sat Mar 18 2000 05:30:00 GMT+0530 (India Standard Time)	11
Flatbed	no	-	-	-
Bus	no	-	-	-
Other	no	-	-	-

LICENSE IMAGES

Front Based on molecular characterization, SARS-CoV-2 is considered a new <i>Betacoronavirus</i> belonging to the subgenus <i>Sarbecovirus</i> (3). A few other critical zoonotic viruses (MERS-related CoV and SARS-related CoV) belong to the same genus. However, SARS-CoV-2 was identified as a distinct virus based on the percent identity with other <i>Betacoronavirus</i>; conserved open reading frame 1a/b (ORF1a/b) is below 90% identity (3). An overall 80% nucleotide identity was observed between SARS-CoV-2 and the original SARS-CoV, along with 89% identity with ZC45 and ZXC21 SARS-related CoVs of bats (2, 31, 36). In addition, 82% identity has been observed between SARS-CoV-2 and human SARS-CoV Tor2 and human SARS-CoV BJ01 2003 (31). A sequence identity of only 51.8% was observed between MERS-related CoV and the recently emerged SARS-CoV-2 (37). Phylogenetic analysis of the structural genes also revealed that SARS-CoV-2 is closer to bat SARS-related CoV. Therefore, SARS-CoV-2 might have originated from bats, while other amplifier hosts might have played a role in disease transmission to humans (31). Of note, the other two zoonotic CoVs (MERS-related CoV and SARS-related CoV) also originated from bats (38, 39). Nevertheless, for SARS and MERS, civet	Back Based on molecular characterization, SARS-CoV-2 is considered a new <i>Betacoronavirus</i> belonging to the subgenus <i>Sarbecovirus</i> (3). A few other critical zoonotic viruses (MERS-related CoV and SARS-related CoV) belong to the same genus. However, SARS-CoV-2 was identified as a distinct virus based on the percent identity with other <i>Betacoronavirus</i>; conserved open reading frame 1a/b (ORF1a/b) is below 90% identity (3). An overall 80% nucleotide identity was observed between SARS-CoV-2 and the original SARS-CoV, along with 89% identity with ZC45 and ZXC21 SARS-related CoVs of bats (2, 31, 36). In addition, 82% identity has been observed between SARS-CoV-2 and human SARS-CoV Tor2 and human SARS-CoV BJ01 2003 (31). A sequence identity of only 51.8% was observed between MERS-related CoV and the recently emerged SARS-CoV-2 (37). Phylogenetic analysis of the structural genes also revealed that SARS-CoV-2 is closer to bat SARS-related CoV. Therefore, SARS-CoV-2 might have originated from bats, while other amplifier hosts might have played a role in disease transmission to humans (31). Of note, the other two zoonotic CoVs (MERS-related CoV and SARS-related CoV) also originated from bats (38, 39). Nevertheless, for SARS and MERS, civet
--	---

SIGNATURE

