

DRIVER APPLICATION FORM

COMPANY DETAILS

Company	Burnett and Mclean Trading	Phone	Mcbride and Ewing Inc
Email	sukeqiq@mailinator.com	City	Flores and Hartman Plc

APPLICANT DETAILS

Full Name	Jared Kelly	DOB	2024-04-18
Phone	Dolore corporis at a	Email	tileh@mailinator.com
Address	Deleniti neque conse		

LICENSE INFORMATION

License No	84	State	Porro ullamco aliqui
Issue Date	2017-12-23	Expiry	1983-06-29

DRIVING EXPERIENCE

Equipment	Experience	From	To	Miles
Straight Truck	yes	Thu May 06 1999 05:30:00 GMT+0530 (India Standard Time)	Mon Jul 17 2017 05:30:00 GMT+0530 (India Standard Time)	70
Straight Truck	yes	Thu May 06 1999 05:30:00 GMT+0530 (India Standard Time)	Mon Jul 17 2017 05:30:00 GMT+0530 (India Standard Time)	70
Truck Tractor	yes	Thu Apr 12 1979 05:30:00 GMT+0530 (India Standard Time)	Sun Feb 23 1997 05:30:00 GMT+0530 (India Standard Time)	8
Semi Trailer	yes	Sat Dec 20 2008 05:30:00 GMT+0530 (India Standard Time)	Wed Apr 24 2019 05:30:00 GMT+0530 (India Standard Time)	34
Double Triple	yes	Wed May 13 1970 05:30:00 GMT+0530 (India Standard Time)	Mon Nov 21 1983 05:30:00 GMT+0530 (India Standard Time)	9
Flatbed	yes	Wed Jun 25 1997 05:30:00 GMT+0530 (India Standard Time)	Tue Jul 07 1987 05:30:00 GMT+0530 (India Standard Time)	100
Bus	no	-	-	-
Other	no	-	-	-

LICENSE IMAGES

Front	Back
We assessed the nucleotide percent similarity using the MegAlign software program, where the similarity between the novel SARS-CoV-2 isolates was in the range of 99.4% to 100%. Among the other <i>Sarbecovirus</i> CoV sequences, the novel SARS-CoV-2 sequences revealed the highest similarity to bat-SL-CoV, with nucleotide percent identity ranges between 88.12 and 89.65%. Meanwhile, earlier reported SARS-CoVs showed 70.6 to 74.9% similarity to SARS-CoV-2 at the nucleotide level. Further, the nucleotide percent similarity was 55.4%, 45.5% to 47.9%, 46.2% to 46.6%, and 45.0% to 46.3% to the other four subgenera, namely, <i>Hibecovirus</i>, <i>Nobecovirus</i>, <i>Merbecovirus</i>, and <i>Embecovirus</i>, respectively. The percent similarity index of current outbreak isolates indicates a close relationship between SARS-CoV-2 isolates and bat-SL-CoV, indicating a common origin. However, particular pieces of evidence based on further complete genomic analysis of current isolates are necessary to draw any conclusions, although it was ascertained that the current novel SARS-CoV-2 isolates belong to the subgenus <i>Sarbecovirus</i> in the diverse range of betacoronaviruses. Their possible ancestor was hypothesized to be from bat CoV strains, wherein bats might have played a crucial role in harboring this class of viruses.	cat and camels, respectively, act as amplifier hosts (40, 41). Coronavirus genomes and subgenomes encode six ORFs (31). The majority of the 5' end is occupied by ORF1a/b, which produces 16 nsps. The two polyproteins, pp1a and pp1ab, are initially produced from ORF1a/b by a -1 frameshift between ORF1a and ORF1b (32). The virus-encoded proteases cleave polyproteins into individual nsps (main protease [Mpro], chymotrypsin-like protease [3CLpro], and papain-like proteases [PLPs]) (42). SARS-CoV-2 also encodes these nsps, and their functions have been elucidated recently (31). Remarkably, a difference between SARS-CoV-2 and other CoVs is the identification of a novel short putative protein within the ORF3 band, a secreted protein with an alpha helix and beta-sheet with six strands encoded by ORF8 (31). Coronaviruses encode four major structural proteins, namely, spike (S), membrane (M), envelope (E), and nucleocapsid (N), which are described in detail below. S Glycoprotein Coronavirus S protein is a large, multifunctional class I viral transmembrane protein. The size of this

SIGNATURE

