Production Solutions of

Pet feeding attractant

SFP-MUL02

Description

The special enzyme preparation for production of pet feeding attractant, specially designed for the efficient hydrolysis of animal visceral raw materials (such as liver, intestinal tissue, fish), they can effectively promote the fine decomposition of large molecular proteins, into small molecular forms that are easy to be absorbed by the digestive system of pets, to ensure the efficient use of nutrition. In this process, we not only managed to retain the rich natural flavor of the raw material, but also minimized the loss of heat-sensitive nutrients by using a gentle enzymatic hydrolysis process, which significantly improved the bioavailability of proteins for pets.

Characteristics

Items	SFP-MUL02
pH	Effective range: 5.0-10.0 Optimum range: 6.0-8.0
Temperature	Effective range:25℃-60℃ Optimum range: 45℃-60℃
Appearance	White to light yellow solid powder

Sunson Industry Group Co., Ltd.: No. 126 Kangdi Road, Jinfeng Industrial Park, Yinchuan, China;
jerry.jiang@sunsonzymes.com www.sunsonenzyme.com

Application


Case 1

Raw Material: Fish meal


Recommend dosage: 0.15% of **SFP-MUL02**

Reaction: 55℃ for 3.5 hours at natural pH.


Inactivate the enzymes: 90℃ for 20mins.



Degree of Hydrolysis



Amino acid nitrogen content in hydrolysate



Distribution of peptide molecular weight in hydrolysate

Case 2


Raw material: Chicken Liver

Recommend dosage: 0.5% of **SFP-MUL02**


Reaction: 55℃ for 3 hours at natural pH.

Inactivate the enzymes: 90℃ for 20mins.


Enzymatic Date of **SFP-MUL02**



Degree of Hydrolysis



Amino acid nitrogen content in hydrolysate



Distribution of peptide molecular weight in hydrolysate

Sunson Industry Group Co., Ltd.: No. 126 Kangdi Road, Jinfeng Industrial Park, Yinchuan, China;
jerry.jiang@sunsonzymes.com www.sunsonenzyme.com


Case 3

Raw Material: Chicken Intestinal


Recommend dosage: 0.5% of **SFP-MUL02**

Reaction: 58℃ for 3 hours at natural pH.


Inactivate the enzymes: 90℃ for 20mins.




Degree of Hydrolysis



Amino acid nitrogen content in hydrolysate



Distribution of peptide molecular weight in hydrolysate



Viscosity of material during treatment