

1) Provide Heavy-Flavor Transport Coefficients ($\mu_B=0$)

- (a) Current best estimate of $D_s(2\pi T)$ as function of T over available T -range (both charm and bottom, if available).
- (b) Normalized momentum dependence of friction coefficient, $A(p;T)/A(p=0;T)$, for current best estimate.
- (c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ (in steps of $dp=0.2\text{GeV}$) and $T=0.16-0.6\text{GeV}$ (steps $dT=0.02\text{GeV}$) for $\mu_B=0$. The idea is to run them through a Langevin simulation in a common hydrodynamic medium evolution.


2) Assess Hadronization and Hadronic Phase (test case: 30-50% 5TeV PbPb collisions)

- (a) Compute $H_{AA}(p_T;T_H) = R_{AA}^{H_Q}(p_T;T_H) / R_{AA}^Q(p_T;T_H)$, the ratio of the R_{AA} of the heavy meson (H_Q) just after hadronization to the R_{AA} of the heavy quark (Q) just before hadronization, for $H_Q=D, \Lambda_c$ (as available) and $Q=c$.
- (b) The same as (a) but for the elliptic flow, v_2 : $H_{v2}(p_T;T_H) = v_2^{H_Q}(p_T;T_H) / v_2^Q(p_T;T_H)$.
- (c) Compute H_{AA} and H_{v2} ratios for D-meson spectra at kinetic freezeout over those right after hadronization (if applicable).


3) Transport Simulations with Imposed Coefficients

- (a) Renormalize the charm-quark transport coefficients with a temperature-dependent but momentum-independent K factor, $K(T)$, as to obtain a temperature-independent value of $D_s(2\pi T) \approx 4$ (for Langevin approaches, $D_s = T / [m_Q A(p=0)]$); then compute R_{AA} and v_2 of charm quarks right before hadronization for 30-50% 5TeV PbPb collisions within your model.
- (b) As an optional assignment (time permitting), to compare transport coefficients from different models: Renormalize current charm-quark transport coefficient, $A(p;T)$, \hat{q}/T^3 for a common R_{AA} in a fixed brick problem (as in Fig. 7 in Phys. Rev. C99 (2019) 054907); then compute R_{AA} and v_2 of charm quarks right before hadronization for 30-50% 5TeV PbPb collisions within your model.


Q1(a) Current best estimate of $D_s(2\pi T)$ as function of T over available T -range (both charm and bottom, if available)




Q1(a) Current best estimate of $D_s(2\pi T)$ as function of T over available T -range (both charm and bottom, if available)




Q1(b) Normalized momentum dependence of friction coefficient,
 $A(p;T)/A(p=0;T)$, for current best estimate




Q1(b) Normalized momentum dependence of friction coefficient, $A(p;T)/A(p=0;T)$, for current best estimate




Q1(b) Normalized momentum dependence of friction coefficient,
 $A(p;T)/A(p=0;T)$, for current best estimate




Q1(b) Normalized momentum dependence of friction coefficient,
 $A(p;T)/A(p=0;T)$, for current best estimate




Q1(b) Normalized momentum dependence of friction coefficient,
 $A(p;T)/A(p=0;T)$, for current best estimate




Q1(b) Normalized momentum dependence of friction coefficient,
 $A(p;T)/A(p=0;T)$, for current best estimate




Q1(b) Normalized momentum dependence of friction coefficient,
 $A(p;T)/A(p=0;T)$, for current best estimate




Q1(b) Normalized momentum dependence of friction coefficient,
 $A(p;T)/A(p=0;T)$, for current best estimate




Q1(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ and $T=0.16-0.6\text{GeV}$ for $\mu_B=0$.




Q1(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ and $T=0.16-0.6\text{GeV}$ for $\mu_B=0$.




Q1(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ and $T=0.16-0.6\text{GeV}$ for $\mu_B=0$.




Q1(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ and $T=0.16-0.6\text{GeV}$ for $\mu_B=0$.




Q1(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ and $T=0.16-0.6\text{GeV}$ for $\mu_B=0$.




Q1(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ and $T=0.16-0.6\text{GeV}$ for $\mu_B=0$.




Q1(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ and $T=0.16-0.6\text{GeV}$ for $\mu_B=0$.




Q1(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ and $T=0.16-0.6\text{GeV}$ for $\mu_B=0$.




Q1(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ and $T=0.16-0.6\text{GeV}$ for $\mu_B=0$.




Q1(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ and $T=0.16-0.6\text{GeV}$ for $\mu_B=0$.




Q1(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ and $T=0.16-0.6\text{GeV}$ for $\mu_B=0$.




Q1(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for $p=0-40\text{GeV}$ and $T=0.16-0.6\text{GeV}$ for $\mu_B=0$.




Q2(a) Compute $H_{AA}(p_T; T_H) = R_{AA}^H(p_T; T_H) / R_{AA}^Q(p_T; T_H)$, the ratio of the R_{AA} of the heavy meson (H_Q) just after hadronization to the R_{AA} of the heavy quark (Q) just before hadronization, for $H=Q=D, \Lambda, \bar{c}$ (as available) and $Q=c$.




Q2(a) Compute $H_{AA}(p_T; T_H) = R_{AA}^H(p_T; T_H) / R_{AA}^Q(p_T; T_H)$, the ratio of the R_{AA} of the heavy meson (H_Q) just after hadronization to the R_{AA} of the heavy quark (Q) just before hadronization, for $H = \Lambda_c, D, \Lambda, \Lambda_b, \Lambda_c, \Lambda_b, \Lambda_c, \Lambda_b$ (as available) and $Q = c, b$.




Q2(b) The same as (a) but for the elliptic flow, v_2 : $H_{v_2}(p_T; T_H) = v_2^H Q(p_T; T_H) / v_2^Q(p_T; T_H)$.




Q2(b) The same as (a) but for the elliptic flow, v_2 : $H_{v_2}(p_T; T_H) = v_2^H Q_{v_2}(p_T; T_H) / v_2^Q(p_T; T_H)$.




Q2(c) Compute H_{AA} and H_{v2} ratios for D-meson spectra at kinetic freezeout over those right after hadronization (if applicable).



Q2(c) Compute H_{AA} and H_{v2} ratios for D-meson spectra at kinetic freezeout over those right after hadronization (if applicable).



Q3(a) Renormalize the charm-quark transport coefficients with a temperature-dependent but momentum-independent K factor, $K(T)$, as to obtain a temperature-independent value of $D_s(2\pi T) = 4$ (for Langevin approaches, $D_s = T / [m_Q A(p=0)]$); then compute R_{AA} and v_2 of charm quarks right before hadronization for 30-50% 5TeV PbPb collisions within your model.



Q3(a) Renormalize the charm-quark transport coefficients with a temperature-dependent but momentum-independent K factor, $K(T)$, as to obtain a temperature-independent value of $D_s(2\pi T) \approx 4$ (for Langevin approaches, $D_s = T / [m_Q A(p=0)]$); then compute R_{AA} and v_2 of charm quarks right before hadronization for 30-50% 5TeV PbPb collisions within your model.

