

**ECT\* Feb.2020, Workshop on Heavy-flavor transport in QCD matter**  
**Pre-Meeting Calculation Requests**

1) Provide Heavy-Flavor Transport Coefficients ( $\mu_B=0$ )

(a) Current best estimate of  $D_s(2\pi T)$  as function of  $T$  over available  $T$ -range (both charm and bottom, if available).

(b) Normalized momentum dependence of friction coefficient,  $A(p;T)/A(p=0;T)$ , for current best estimate.

(c) Table of current best estimates of charm friction and momentum-diffusion coefficients for  $p=0-40\text{GeV}$  (in steps of  $dp=0.2\text{GeV}$ ) and  $T=0.16-0.6\text{GeV}$  (steps  $dT=0.02\text{GeV}$ ) for  $\mu_B=0$ . The idea is to run them through a Langevin simulation in a common hydrodynamic medium evolution.

2) Assess Hadronization and Hadronic Phase (test case: 30-50% 5TeV PbPb collisions)

(a) Compute  $H_{AA}(p_T;T_H) = R_{AA}^{H_Q}(p_T;T_H) / R_{AA}^Q(p_T;T_H)$ , the ratio of the  $R_{AA}$  of the heavy meson ( $H_Q$ ) just after hadronization to the  $R_{AA}$  of the heavy quark ( $Q$ ) just before hadronization, for  $H_Q=D, \Lambda_c$  (as available) and  $Q=c$ .

(b) The same as (a) but for the elliptic flow,  $v_2$ :  $H_{v_2}(p_T;T_H) = v_2^{H_Q}(p_T;T_H) / v_2^Q(p_T;T_H)$ .

(c) Compute  $H_{AA}$  and  $H_{v_2}$  ratios for D-meson spectra at kinetic freezeout over those right after hadronization (if applicable).

3) Transport Simulations with Imposed Coefficients

(a) Renormalize the charm-quark transport coefficients with a temperature-dependent but momentum-independent  $K$  factor,  $K(T)$ , as to obtain a temperature-independent value of  $D_s(2\pi T) = 4$  (for Langevin approaches,  $D_s = T/[m_Q A(p=0)]$ ); then compute  $R_{AA}$  and  $v_2$  of charm quarks right before hadronization for 30-50% 5TeV PbPb collisions within your model.

(b) As an optional assignment (time permitting), to compare transport coefficients from different models: Renormalize current charm-quark transport coefficient,  $A(p;T)$ ,  $\hat{q}/T^3$  for a common  $R_{AA}$  in a fixed brick problem (as in Fig. 7 in Phys. Rev. C99 (2019) 054907); then compute  $R_{AA}$  and  $v_2$  of charm quarks right before hadronization for 30-50% 5TeV PbPb collisions within your model.