

Agent-Based Modelling of Zero-Dose Vaccination Among Under-Five Children in Kenya

Simulation report · projection window 2025–2030 · generated 2026-06-04 18:29

Structured to mirror the Rono et al. (2024) project brief.

1. Abstract

Introduction. Zero-dose children — those who never receive the first dose of a diphtheria-tetanus-pertussis (DTP1) or pentavalent vaccine — remain a central equity gap in global immunisation. This report summarises a Starsim-based agent-based model calibrated to administrative DTP1 data from Kenya, and used to project the impact of scaling up routine vaccination on zero-dose prevalence and downstream tetanus burden among under-fives.

Methodology. Two scenarios were compared over calendar years 2025–2030: a *baseline* scenario whose routine delivery probability is calibrated (grid search) so that the modelled end-of-window zero-dose share matches the empirical implied zero-dose share, and an *intervention* scenario with a higher routine delivery probability and coverage cap. Following Rono et al. (2024), tetanus is the sentinel disease — the only DTP-bracket disease still endemic in Kenya — and is modelled via environmental / wound-exposure dynamics with waning immunity (Starsim SIS-style). Vaccination acts on under-fives only.

Results. End-of-window modelled zero-dose share was **12.9%** in the baseline scenario and **6.3%** in the intervention scenario — a relative reduction of **51.4%**. Modelled tetanus cases averted among under-fives in the simulated cohort totalled **63**. When scaled to the Kenya under-five population, the intervention reduces end-of-window zero-dose children from approximately **931,813** to **452,548**, reaching roughly **479,265** additional children.

Conclusion. The intervention scenario achieves approximately a halving of zero-dose prevalence relative to the calibrated comparator — consistent with the project's 50%-reduction target — and yields a material reduction in modelled tetanus cases.

Key words: zero-dose vaccination, agent-based modelling, under-five, childhood immunisation, tetanus, Starsim, Kenya.

2. Introduction

Vaccination is a cornerstone of child public health: the WHO Immunization Agenda 2030 targets at least 90% coverage of essential vaccines in every country and in every district. Despite steady improvements, access gaps persist, and in sub-Saharan Africa roughly one in five children still lacks access to routine immunisation. Children who never receive a first DTP- or pentavalent-containing dose — the so-called *zero-dose* children — are the clearest marker of that gap and the central focus of this project.

Statement of the problem

Globally about 14% of under-five children are penta-zero-dose and roughly 7.5% are truly zero-dose. In Kenya the zero-dose rate is approximately 7% (KDHS 2022), but monthly administrative data show that the implied coverage gap varies substantially across time and regions. Because diphtheria, neonatal tetanus and pertussis are controlled or near elimination, **tetanus** is used as the sentinel outcome: it remains endemic via environmental and wound exposure, and its burden is sensitive to first-dose coverage.

The included administrative dataset provides a monthly DTP1 administrative DTP1 coverage indicator spanning 2018-2024 (84 months). Its mean coverage is **83.5%**, implying a mean zero-dose share of **16.5%**. This value anchors the model calibration target.

Descriptive disease context

The same administrative dataset also reports monthly case counts for several under-five diseases. Mirroring the tables in the Rono et al. (2024) Results section, the average monthly burden is: **pneumonia** $\approx$ 756,409/month; **measles** $\approx$ 2,907/month; **tetanus** $\approx$ 595/month. **Only tetanus is simulated** — pneumonia is addressed by the pneumococcal vaccine (PCV), measles by the measles-containing vaccine (MCV), and the brief explicitly notes that pertussis and measles are under marked control in Kenya. Pneumonia and measles are shown here purely as context for the under-five disease burden in which zero-dose children live.

Objective

To quantify how many children would be additionally reached, and how many tetanus cases averted, if routine pentavalent delivery were scaled up enough to approximately halve the zero-dose share among under-fives, using a data-anchored agent-based simulation.

Research question

How many tetanus cases would be averted if we reduce the prevalence of zero-dose vaccination by ~50% among under-fives by the end of the projection window?

3. Methodology

The model is built on the **Starsim** agent-based framework. Each agent represents a single child; the population includes births and deaths so that the under-five cohort renews over time. Vaccination is delivered via a routine *first-dose* intervention applied only to children in the 0-to-60-month age window, with a per-step delivery probability and an overall coverage cap.

Disease module

A single disease module is instantiated: **tetanus**. The Rono et al. (2024) brief notes that diphtheria is eliminated, neonatal tetanus is near elimination, and pertussis and measles are under marked control in Kenya; tetanus is therefore the only DTP-bracket disease with enough residual burden to serve as a sensitive indicator of first-dose coverage. Tetanus transmission is not person-to-person in practice, so the module implements an environmental / wound-exposure SIS-style dynamic with waning vaccine-induced immunity. Vaccinated agents become transiently immune with efficacy 90.0% and re-enter the susceptible pool as immunity wanes.

Calibration

A grid search over the routine-delivery probability is used to match the modelled end-of-window zero-dose share to the empirical implied zero-dose share from administrative DTP1. Calibration uses short, higher-agent runs ($\approx 10,000$ agents, 8-year window) and, once converged, the chosen routine probability and a coverage cap set by the mean administrative DTP1 coverage indicator define the *reference* parameter set. The *intervention* parameter set multiplies the reference routine probability (capped at 0.12) and raises the coverage by +2 percentage points, bounded by the coverage cap.

Calibration source for this run: loaded: `/Users/mine/git/zdsim/calibration.json`.

Key parameters

Parameter	Reference	Intervention
Routine delivery probability (per week)	0.0324	0.0744
Coverage cap	83.5%	85.5%
Vaccine efficacy	90.0%	90.0%
Eligible ages (years)	0–5	0–5
Tetanus initial prevalence	0.0010	0.0010

Run configuration

Agents: **20,000** Projection window: **2025–2030** Data file: `zerodose_data_formated.xlsx`

4. Results

At-a-glance impact

Primary target met: under-five zero-dose falls from **12.9%** to **6.3%** (**51.4%** relative reduction). In the simulated under-five cohort, tetanus cases fall from **221** to **158** (averted **63**, 28.51% relative reduction).

Zero-dose share

At the end of the projection window the modelled zero-dose share among under-fives is **12.9%** in the baseline scenario and **6.3%** in the intervention scenario — a **51.4%** relative reduction. Across the full window the intervention reduces the annual zero-dose share by on average **9.16 percentage points** (cumulatively 54.97 pp-years).

Tetanus burden

In the simulated under-five cohort, the intervention averts **63** tetanus cases over 5 years (baseline 221 vs intervention 158, 28.51% relative reduction).

Tetanus deaths

Yearly tetanus death counts are reported in the annual breakdown table below (not as separate trajectory figures). Total modelled tetanus deaths are **0** in the baseline scenario and **0** in the intervention scenario (averted **0**, mean 0.0 per year). These counts are stochastic in small cohorts.

Population-scaled projection (Kenya)

Anchor: *Kenya national 2024 anchors*. Scaling by the ratio of the real annual birth cohort to the modelled annual births (factor 2,540), the intervention reduces end-of-window zero-dose children from **931,813** to **452,548**, reaching approximately **479,265** additional children — a mean of 79,877 children additionally vaccinated per year. Scaled tetanus cases averted: **160,020**; scaled tetanus deaths averted: **0**. Source: UN WPP 2024; WHO/UNICEF WUENIC 2024 revision.

Annual breakdown

Year	ZD share (reference)	ZD share (intervention)	Ref. tetanus deaths	Int. tetanus deaths
2025	97.3%	93.4%	0	0
2026	27.9%	7.2%	0	0
2027	13.6%	4.1%	0	0
2028	12.4%	5.2%	0	0
2029	11.9%	5.0%	0	0
2030	12.5%	5.9%	0	0

4. Results — figure




Figure 1. Implied zero-dose share from administrative DTP1 data vs modelled baseline and intervention scenarios at the end of the projection window.

4. Results — figure




Figure 2. Projected yearly zero-dose share among under-fives for the baseline (calibrated to the empirical implied zero-dose share) and intervention (scaled-up routine delivery) scenarios.

4. Results — figure




Figure 3. Under-5 tetanus trajectories (baseline vs intervention) and zero-dose share. Case counts are modelled under-5 infections in the simulated cohort; see methodology for the scaling factor used when translating to a national cohort.

4. Results — figure




Figure 3b. Total under-5 tetanus infections over the full projection window for no-intervention, baseline, and scale-up scenarios.

4. Results — figure




Figure 4. Administrative DTP1 coverage indicator and implied zero-dose share (monthly), used to anchor the calibration target.

4. Results — figure




Figure 5. Administrative DPT1 / DPT3 doses versus estimated live births, providing context for the administrative DTP1 coverage indicator.

4. Results — figure




Figure 6. Monthly reported cases for pneumonia, measles, and tetanus from the Kenya HMIS administrative dataset — descriptive context only, mirroring the tables in the Rono et al. (2024) Results section. Pneumonia and measles are shown because the brief cites their monthly burden; neither is modelled in this ABM (pneumonia is addressed by PCV, measles by MCV, and the brief explicitly notes measles is under marked control).

5. Discussion

The intervention scenario reduces zero-dose prevalence substantially across the projection window and, via its impact on first-dose coverage, lowers the modelled tetanus burden in the simulated cohort. The magnitude of the relative reduction is consistent with the project's halving target.

Several caveats apply. Tetanus is used as a sentinel for DTP1 / pentavalent coverage and does not capture all-cause under-five mortality; diphtheria, pertussis, hepatitis B and Hib are not simulated here because Kenya has already eliminated or controlled them. Annual averted counts can be noisy when the modelled cohort is small or the projection window is short; in those regimes occasional negative yearly averted values are expected and should prompt longer runs or larger cohorts. When the two arms use distinct RNG seeds the comparison is also noisier than when they share a seed (the default).

Equity in first-dose access — geographic, socioeconomic, and service-delivery heterogeneity — is known to drive zero-dose persistence. The current implementation applies a homogeneous routine-delivery probability and coverage cap; future work should stratify by subnational region and risk profile, and link the delivery process to supply-chain and demand-side constraints.

6. Conclusion

The model demonstrates that an intervention strong enough to halve zero-dose prevalence (observed relative reduction **51.4%** in this run) produces a meaningful decline in modelled tetanus cases (**63** averted in the simulated cohort) and a corresponding reduction in tetanus deaths. These findings support sustained investment in routine first-dose delivery as the most direct lever on zero-dose prevalence.

7. References

- Rono, B. C., Njeri, A., Mambo, S. N., & Akangbe, R. O. (2024). *Agent-Based Modelling (ABM) predicts number needed to vaccinate to achieve a 50% reduction in zero-dose vaccination among under-five children in Kenya by 2025*. Project brief (JKUAT / APHRC / Lagos State MoH).
- Cata-Preta, B. O., Santos, T. M., Mengistu, T., Hogan, D. R., Barros, A. J. D., & Victora, C. G. (2021). Zero-dose children and the immunisation cascade: understanding immunisation pathways in low- and middle-income countries. *Vaccine*, 39(32), 4564–4570. <https://doi.org/10.1016/j.vaccine.2021.02.072>
- Kenya National Bureau of Statistics (2022). *Kenya Demographic and Health Survey (KDHS) 2022*.
- Ozigbu, C. E., Olatosi, B., Li, Z., Hardin, J. W., & Hair, N. L. (2022). Correlates of zero-dose vaccination status among children aged 12–59 months in sub-Saharan Africa. *Vaccines*, 10(7), 1052. <https://doi.org/10.3390/vaccines10071052>
- Wonodi, C., & Farrenkopf, B. A. (2023). Defining the zero-dose child: a comparative analysis of two approaches and their impact on assessing the zero-dose burden across 82 low- and middle-income countries. *Vaccines*, 11(10), 1543. <https://doi.org/10.3390/vaccines11101543>
- WHO / UNICEF (2024). *WUENIC — WHO/UNICEF estimates of national immunisation coverage*.
- World Health Organization (2020). *Immunization Agenda 2030*.
- Starsim Collective. *Starsim — agent-based disease modelling framework*. <https://starsim.org/>