

0.1 Exact equations of motion for spin-1

$$\mathcal{L} = -\frac{1}{2}(D^\mu W^\nu - D^\nu W^\mu)^\dagger (D_\mu W_\nu - D_\nu W_\mu) + m^2 W^{\mu\dagger} W_\mu - i\lambda e W^{\mu\dagger} W^\nu F_{\mu\nu} \quad (0.1)$$

where as usual D is the long derivative $D^\mu = \partial^\mu + ieA^\mu$.

Obtain the equations of motion from the Euler-Lagrange equations:

$$\frac{\partial \mathcal{L}}{\partial W^{\dagger\alpha}} - \partial_\mu \frac{\partial \mathcal{L}}{\partial [\partial_\mu W^{\dagger\alpha}]} = 0$$

Or equivalently,

$$\frac{\partial \mathcal{L}}{\partial W^{\dagger\alpha}} - D_\mu \frac{\partial \mathcal{L}}{\partial [D_\mu W^{\dagger\alpha}]} = 0$$

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial W^{\dagger\alpha}} &= \frac{\partial}{\partial W^{\dagger\alpha}} \left(m^2 W^{\mu\dagger} W_\mu - i\lambda e W^{\mu\dagger} W^\nu F_{\mu\nu} \right) \\ &= m^2 W_\alpha - ie\lambda W^\nu F_{\alpha\nu} \end{aligned}$$

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial [D_\gamma W^{\dagger\alpha}]} &= -\frac{1}{2} \frac{\partial}{\partial [D_\gamma W^{\dagger\alpha}]} (D^\mu W^\nu - D^\nu W^\mu)^\dagger (D_\mu W_\nu - D_\nu W_\mu) \\ &= -\frac{1}{2} (g^{\mu\gamma} g_\alpha^\nu - g^{\nu\gamma} g_\alpha^\mu) (D_\mu W_\nu - D_\nu W_\mu) \\ &= D^\gamma W_\alpha - D_\alpha W^\gamma \end{aligned}$$

So the complete equation from Euler-Lagrange is

$$m^2 W_\alpha - ie\lambda W^\nu F_{\alpha\nu} - D_\mu (D^\mu W_\alpha - D_\alpha W^\mu) = 0 \quad (0.2)$$

This is a set of four coupled second order equations for the field W . We rewrite as a set of first order equations by introducing a field $W_{\mu\nu} = D_\mu W_\nu - D_\nu W_\mu$. So (0.2) becomes

$$m^2 W_\alpha - ie\lambda W^\mu F_{\alpha\mu} - D^\mu W_{\mu\alpha} = 0 \quad (0.3)$$

$W^{\mu\nu}$ is antisymmetric and so has six degrees of freedom, corresponding to six independent fields. Together with W^μ this represents a total of ten fields. However, upon examination only some of these fields are dynamic. The fields W^{0i} and W^i appear in the equations with time derivatives, while the fields W^{ij} and W^0 never do. So it is only necessary to consider the former six fields. So that these six fields all have the same dimension, we will define $\frac{W^{i0}}{m} = i\eta^i$.

We will now eliminate the extraneous fields and solve for $iD_0 W^i$, $iD_0 \eta^i$.

$$m^2 W_\alpha - ie\lambda W^\nu F_{\alpha\nu} - D_\mu (D^\mu W_\alpha - D_\alpha W^\mu) = 0 \quad (0.4)$$

Define $W_{\mu\nu} = D_\mu W_\nu - D_\nu W_\mu$. Then

$$m^2 W_\alpha - ie\lambda W^\nu F_{\alpha\nu} - D^\mu W_{\mu\alpha} = 0 \quad (0.5)$$

To get the exact Hamiltonian of a bispinor, we can eliminate nondynamic fields.

First consider (0.5) with $\alpha = 0$.

$$m^2 W_0 - ie\lambda W^\nu F_{0\nu} - D^\mu W_{\mu 0} \quad (0.6)$$

$$m^2 W_0 - ie\lambda W^j F_{0j} - D^j W_{j0} \quad (0.7)$$

Solve this for W_0

$$W_0 = \frac{1}{m^2} (ie\lambda W^j F_{0j} + D^j W_{j0}) \quad (0.8)$$

Now, consider (0.5) with $\alpha = i$

$$m^2 W_i - ie\lambda W^\mu F_{i\mu} - D^\mu W_{\mu i} = 0 \quad (0.9)$$

$$m^2 W_i - ie\lambda W^0 F_{i0} - D^0 W_{0i} - ie\lambda W^j F_{ij} - D^j W_{ji} = 0 \quad (0.10)$$

Using (0.8) we can replace W_0

$$m^2 W_i - \frac{ie\lambda}{m^2} (ie\lambda W^\nu F_{0\nu} + D^j W_{j0}) F_{i0} - D^0 W_{0i} - ie\lambda W^j F_{ij} - D^j W_{ji} = 0 \quad (0.11)$$

Using $W_{ji} = D_j W_i - D_i W_j$

$$m^2 W_i - \frac{ie\lambda}{m^2} (ie\lambda W^\nu F_{0\nu} + D^j W_{j0}) F_{i0} - D^0 W_{0i} - ie\lambda W^j F_{ij} - D^j (D_j W_i - D_i W_j) = 0 \quad (0.12)$$

Solve this for $D^0 W_{0i}$:

$$D^0 W_{0i} = m^2 W_i - \frac{ie\lambda}{m^2} (ie\lambda W^\nu F_{0\nu} + D^j W_{j0}) F_{i0} - ie\lambda W^j F_{ij} - D^j (D_j W_i - D_i W_j) \quad (0.13)$$

To get a similar equation for W_i , consider

$$W_{i0} = D_i W_0 - D_0 W_i \quad (0.14)$$

Then

$$D^0 W_i = D_i W_0 - W_{i0} \quad (0.15)$$

$$D^0 W_i = D_i \frac{1}{m^2} (ie\lambda W^j F_{0j} + D^j W_{j0}) - W_{i0} \quad (0.16)$$

We now have equations that tell us the time evolution of a total of six fields: W_i and W_{i0} . We want to treat these as the components of some sort of bispinor. To that end, first define $\eta_i = -i/mW_{i0}$ so that we have a pair of fields with the same mass dimension and hermiticity. (Since $W_{i0} = D_i W_0 - D_0 W_i$ would pick up another minus sign under complex conjugation compared to W_i .) Going the other way $W_{i0} = im\eta_i$.

Since eventually we want a nonrelativistic expression, we should write the spatial components of four vectors as three vectors. To that end also write the components of the tensor $F_{\mu\nu}$ in terms of three vectors.

$$F_{0i} = E^i, \quad F_{ij} = -\epsilon_{ijk} B^k$$

Regular spatial vectors are “naturally raised” while D_i is “naturally lowered”. Then we can rewrite (0.13).

$$-imD_0\eta_i = m^2 W_i + \frac{ie\lambda}{m^2} (ie\lambda W^j E^j + D^j W_{j0}) E^i - ie\lambda W^j \epsilon_{ijk} B^k - D^j (D_j W_i - D_i W_j) \quad (0.17)$$

$$iD_0\eta^i = -mW^i + \frac{ie\lambda}{m^3} (ie\lambda W^j E^j + D^j W_{j0}) E^i - \frac{ie\lambda}{m} W^j \epsilon_{ijk} B^k - \frac{1}{m} D_j (D_j W^i - D_i W^j) \quad (0.18)$$

And likewise (0.16) becomes

$$D^0 W_i = D_i \frac{1}{m^2} (-ie\lambda W^j E^j + imD^j \eta_j) - im\eta_i \quad (0.19)$$

$$iD^0 W^i = -iD_i \frac{1}{m^2} (-ie\lambda W^j E^j + imD_j \eta^j) + m\eta^i \quad (0.20)$$

$$iD^0 W^i = -\frac{1}{m^2} D_i e\lambda \mathbf{W} \cdot \mathbf{E} + \frac{1}{m} D_i \mathbf{D} \cdot \boldsymbol{\eta} + m\eta^i \quad (0.21)$$

0.1.1 Spin identities

To obtain some Shrodinger like equation for a bispinor, we need to express the time derivative of the bispinor in terms of other operators which can be interpreted as the Hamiltonian: $i\partial_0\Psi = \hat{H}\Psi$. We have expressions for the time derivatives of W_i and η_i , but some mixing of the indices is involved. To write the Hamiltonian in a block form we can introduce spin matrices whose action will mix the components of the fields.

The spin matrix for a spin one particle can be represented as:

$$(S^k)_{ij} = -i\epsilon_{ijk} \quad (0.22)$$

which leads to the following identities:

$$(\mathbf{a} \times \mathbf{v})_i = -i(\mathbf{S} \cdot \mathbf{a})_{ij}v_j \quad (0.23)$$

$$\{\mathbf{a} \times (\mathbf{a} \times \mathbf{v})\}_i = -(\{\mathbf{S} \cdot \mathbf{a}\}^2)_{ij}v_j \quad (0.24)$$

$$a_i(\mathbf{b} \cdot \mathbf{v}) = \{\mathbf{a} \cdot \mathbf{b} \delta_{ij} - (S^k S^\ell)_{ij} a^\ell b^k\} v_j \quad (0.25)$$

Using these identities

$$iD^0\mathbf{W} = -\frac{e\lambda}{m^2}\mathbf{D}(\mathbf{W} \cdot \mathbf{E}) + \frac{1}{m}\mathbf{D}(\mathbf{D} \cdot \boldsymbol{\eta}) + m\boldsymbol{\eta} \quad (0.26)$$

$$iD^0\mathbf{W} = -\frac{1}{m^2}\{\mathbf{D} \cdot \mathbf{E} - (\mathbf{S} \cdot \mathbf{E})(\mathbf{S} \cdot \mathbf{D})\}\mathbf{W} + \frac{1}{m}\{\mathbf{D}^2 - (\mathbf{S} \cdot \mathbf{D})^2\}\boldsymbol{\eta} + m\boldsymbol{\eta} \quad (0.27)$$

and for the other equation

$$iD_0\eta^i = -mW^i + \frac{ie\lambda}{m^3}(ie\lambda W^j E^j + D^j W_{j0})E^i - \frac{ie\lambda}{m}W^j \epsilon_{ijk}B^k - \frac{1}{m}D_j(D_j W^i - D_i W^j) \quad (0.28)$$

$$iD_0\eta^i = -mW^i + \frac{ie\lambda}{m^3}(ie\lambda W^j E^j + D^j W_{j0})E^i - \frac{ie\lambda}{m}W^j \epsilon_{ijk}B^k - \frac{1}{m}D_j(D_j W^i - D_i W^j) \quad (0.29)$$

$$iD_0\boldsymbol{\eta} = -m\mathbf{W} - \frac{e^2\lambda^2}{m^3}\mathbf{E}(\mathbf{W} \cdot \mathbf{E}) - \frac{1}{m^2}\mathbf{E}(\mathbf{D} \cdot \boldsymbol{\eta}) - \frac{ie\lambda}{m}(\mathbf{W} \times \mathbf{B}) + \frac{1}{m}\mathbf{D} \times (\mathbf{D} \times \mathbf{W}) \quad (0.30)$$

$$iD_0\boldsymbol{\eta} = -m\mathbf{W} - \frac{e^2\lambda^2}{m^3}\{\mathbf{E}^2 - (\mathbf{S} \cdot \mathbf{E})^2\}\mathbf{W} - \frac{1}{m^2}\{\mathbf{E} \cdot \mathbf{D} - (\mathbf{S} \cdot \mathbf{E})(\mathbf{S} \cdot \mathbf{D})\}\boldsymbol{\eta} - \frac{e\lambda}{m}(\mathbf{S} \cdot \mathbf{B})\mathbf{W} - \frac{1}{m}(\mathbf{S} \cdot \mathbf{D})^2\mathbf{W} \quad (0.31)$$

$$iD_0\boldsymbol{\eta} = \left(-m - \frac{e^2\lambda^2}{m^3}\{\mathbf{E}^2 - (\mathbf{S} \cdot \mathbf{E})^2\} - \frac{e\lambda}{m}(\mathbf{S} \cdot \mathbf{B}) - \frac{1}{m}(\mathbf{S} \cdot \mathbf{D})^2\right)\mathbf{W} - \frac{1}{m^2}\{\mathbf{E} \cdot \mathbf{D} - (\mathbf{S} \cdot \mathbf{E})(\mathbf{S} \cdot \mathbf{D})\}\boldsymbol{\eta} \quad (0.32)$$

Now that the equations are in the correct form, we can write them as follows:

$$iD_0 \begin{pmatrix} W \\ \eta \end{pmatrix} = \begin{pmatrix} -\frac{1}{m^2}\{\mathbf{D} \cdot \mathbf{E} - (\mathbf{S} \cdot \mathbf{E})(\mathbf{S} \cdot \mathbf{D})\} & m + \frac{1}{m}\{\mathbf{D}^2 - (\mathbf{S} \cdot \mathbf{D})^2\} \\ -m - \frac{e^2\lambda^2}{m^3}\{\mathbf{E}^2 - (\mathbf{S} \cdot \mathbf{E})^2\} - \frac{e\lambda}{m}(\mathbf{S} \cdot \mathbf{B}) - \frac{1}{m}(\mathbf{S} \cdot \mathbf{D})^2 & -\frac{1}{m^2}\{\mathbf{E} \cdot \mathbf{D} - (\mathbf{S} \cdot \mathbf{E})(\mathbf{S} \cdot \mathbf{D})\} \end{pmatrix} \begin{pmatrix} W \\ \eta \end{pmatrix} \quad (0.33)$$