

MATH 1342: Elementary Statistics

Fall 2024 & Spring 2025

1 Unit 1: Exploring One-Variable Data

2

1 Unit 1: Exploring One-Variable Data

Basics of the AP Statistics Exam:

The AP Statistics Exam will test your understanding of the mathematical concepts covered in the course units, including your ability to use statistical methods and calculate the probability of an outcome. You'll need to bring a graphing calculator with statistics capabilities to the exam.

The exam is 3 hours long. 50% of the score will be 40 multiple choice questions in 90 minutes, and 50% of the score will be 6 free response questions in 90 minutes.

Unit 1 will be 15-23% of the exam score and contains variation in categorical and quantitative variables, representing data using tables or graphs, calculating and interpreting statistics, describing and comparing distributions of data, and the normal distribution.

1.1 - Representing Categorical and Quantitative Variables with Graphs

Data contains information about a group of individuals. The information is organized using variables.

Individuals are objects described by a set of data. Individuals may be people but may be animals or inanimate objects.

Variables are characteristics of individuals. A variable may take on different values of different variables. Variables can be split into two types: categorical or quantitative.

Categorical variables place individuals into specific groups.

Quantitative variables takes on numerical values for which it makes sense to do arithmetic operations like adding and averaging. Quantitative variables fall into two categories: discrete and continuous.

Be careful: just because it is a number doesn't make it quantitative.

Discrete variables are numerical values where counting makes sense; in other words, decimals would not be an appropriate way to record the data.

Continuous variables are numerical values where decimals are appropriate; it usually involves some form of measuring.

The difference between discrete and continuous isn't always clear. An example of this would be age.

One of the easiest ways to display categorical data is with a table.

Count is the amount of that category in a table and relative count is $\frac{\text{count}}{\text{total}}$.

If you wanted to display two categorical variables at a time, we could make a two way table.

To better visualize the data, there are graphs that we can make from the data. We want to visualize the graphs to get a better idea of the distribution.

Distribution of a variable tells us what values the variable takes and how often it takes these values.

Bar Graphs have the following characteristics:

- Label each axis clearly
- The x-axis will contain the categorical variable and the y-axis will display the counts
- Each category has its own bar and the bars cannot touch

- Order is not important when creating the x-axis

To make a histogram, we need to put the data into even intervals that capture our data. We will do this first by hand by counting how many data scores are in each bin.

To find the interval width, we can use the formula $\frac{\text{max-min}}{\# \text{ of wanted intervals}}$

To make the histogram:

- Draw rectangles for each interval with height representing the count
- Bars must touch
- Label the x-axis with the lower bound values of each interval