

STAT 88: Lecture 2

Contents

Section 1.3: Fundamental Rules

Last time Probability allows you to learn about a random sample from a known population

Probability: population \rightarrow sample

Statistics: sample \rightarrow population

Sec 1.1 Probability as Proportions

We call the set of all outcomes of an experiment Ω , the outcome space or the sample space. Let $A \subseteq \Omega$ be an event.

For equally likely outcomes, $P(A) = \frac{\#A}{\#\Omega}$.

Sec 1.2 Probability Bounds



When we don't know how much events overlap, we sometimes need to give upper and lower bounds for what a probability is.

Example: Let $A, B, C \subseteq \Omega$ with $P(A) = 0.1$, $P(B) = 0.05$, $P(C) = 0.01$. Then

$$\left(\begin{array}{l} \underline{P(A \cup B \cup C)} \leq \underline{P(A)} + P(B) + P(C) = 0.16 \text{ and} \\ \underline{P(A \cup B \cup C)} \geq \underline{P(A)} = 0.1. \end{array} \right.$$

Warm up: Based on historical averages,

- $P(A)$ = Chance that you catch bus to school = 70%.
- $P(B)$ = Chance that it rains = 50%.



- $P(C)$ = Chance you make it to class on time = 10%.


What is the chance that at least one of these events occurs (making no assumptions).
If it cannot be found exactly, find the best lower bound and upper bound that you can.

Exercise

Soln Find the "union" of the three events, i.e. $P(A \cup B \cup C)$
We can't find the answer exactly b/c we don't know if events A, B, C overlap.

① Upper bound : Extreme case.


$$P(A \cup B \cup C) \leq P(A) + P(B) + P(C) \\ = 30\%$$




Since $P(A \cup B \cup C)$ cannot be bigger than 100%,
the upper bound is 100%

② Lower bound : Another extreme case

$$P(A \cup B \cup C) \geq P(A) = \underline{70\%}$$



1.3. Fundamental Rules

In 1933, the Russian mathematician Andrey Kolmogorov established the axioms of the modern theory of probability. Intuitively, the axioms are generalizations of the natural properties of measures like length, area, or volume (think of Venn diagrams where $P(A)$ is the area of circle A). 



Formally, probability is a function on events, $P : A \subseteq \Omega \mapsto [0, 1]$, satisfying the following 3 axioms:

1. $P(A) \geq 0$ for all $A \subseteq \Omega$.

2. $P(\Omega) = 1$.

3. **Addition Rule.** If A and B are mutually exclusive, i.e. $A \cap B = \emptyset$, then $P(A \cup B) = P(A) + P(B)$.

Venn Diagram




Consequence of three axioms:

1. Complement rule: $P(A^c) = 1 - P(A)$.

$$B = A^c, P(\Omega) = 1$$



2. Difference rule: If $B \subseteq A$ then $P(A \setminus B) = P(A) - P(B)$.




3. Boole/Bonferroni's inequality: $P(\cup_{i=1}^n A_i) \leq \sum_{i=1}^n P(A_i)$ for all events A_i 's.

$$n=2 \Rightarrow \underline{P(A \cup B) \leq P(A) + P(B)} \quad (\text{Upper Bound})$$

Boole/Bonferroni's inequality is widely used in statistics and machine learning theory. In many cases, however, it gives rather a crude bound, e.g. if $P(A) = 0.8$ and $P(B) = 0.9$ then $P(A \cup B) \leq 1.7$ which is true but rather silly as we already know $P(A \cup B) \leq 1$.

$$\leq P(A) + P(B)$$

De Morgan's laws: $(A \cap B)^c = A^c \cup B^c$.



This generalizes to 3 or more events.

$$(A \cap B \cap C)^c = A^c \cup B^c \cup C^c$$

Another form of De Morgan's laws.

$$(A \cup B)^c = A^c \cap B^c$$


Example:

- $P(A)$ = Chance that you catch bus to school = 70%.
- $P(B)$ = Chance that it rains = 50%.

$P(A \cap B) = P(A)P(B)$ true
only when A and B independent
↑


Find the best lower bound for $P(A \cap B)$ you can without making any assumptions.

① Visual Soln



* We want the intersection to be as small as possible.

$70 - 50 = 20\%$: Smallest possible overlap.



② Using axioms and De Morgan's laws.

$$P(A \cap B) = 1 - P((A \cap B)^c)$$

De Morgan's laws

$$= 1 - P(A^c \cup B^c)$$

$$\geq 1 - P(A^c) - P(B^c)$$

$$= 1 - 0.3 - 0.5$$

$$= 0.2$$

* We know


$$P(A^c \cup B^c) \leq P(A^c) + P(B^c)$$

Example:

- $P(A)$ = Chance that you catch bus to school = 70%.
- $P(B)$ = Chance that it rains = 50%.
- $P(C)$ = Chance you make it to class on time = 10%.

Find the best lower bound for $P(A \cap B \cap C)$ you can without making any assumptions.

① Visual soln



It's possible for $B = A \cap B$ and C
not to have any overlaps, i.e. $A \cap B \cap C = \emptyset$
 $\Rightarrow P(A \cap B \cap C) \geq 0\%$

If $C > 80\%$, then
lower bound $> 0\%$.
 $A \cap B = 20\%$

② Using axioms and De Morgan's laws

$$\begin{aligned} P(A \cap B \cap C) &= 1 - P((A \cap B \cap C)^c) \\ &\xrightarrow{\text{De Morgan's laws}} = 1 - P(A^c \cup B^c \cup C^c) \\ &\geq 1 - P(A^c) - P(B^c) - P(C^c) \\ &= 1 - 0.3 - 0.5 - 0.9 \\ &= -0.7. \end{aligned}$$

$$\begin{aligned} P(A) &= 70\% \\ \Rightarrow P(A^c) &= 30\% \end{aligned}$$

We also know $P(A \cap B \cap C) \geq 0$.