

Sample Size (7.4), Confidence Intervals for Variance and Standard Deviation

Stat 400 - April 1, 2021

Today's topics

Review:

- Chi Squared Distribution, t distribution, CI for mean

New:

- Confidence Interval for variance (or sd)
- Required Sample Size

Chi-squared distribution (review) χ^2

The Chi-Squared distribution is a special case of the Gamma distribution where $\theta = 2$.

Also, sum of squares of normal distributions

<https://homepage.divms.uiowa.edu/~mbognar/applets/chisq.html>

notes

notes


t distribution:

If σ is unknown:

Use s instead of σ

$$T = \frac{\bar{X} - \mu}{s / \sqrt{n}} \sim t_{n-1}$$

↗
if I "standardize" the
sample mean using 's'
instead of 'σ'.



t distribution

Theorem 5.5-3

(Student's t distribution) Let


$$T = \frac{Z}{\sqrt{U/r}},$$

where Z is a random variable that is $N(0, 1)$, U is a random variable that is $\chi^2(r)$, and Z and U are independent. Then T has a t distribution with pdf

$$f(t) = \frac{\Gamma((r+1)/2)}{\sqrt{\pi r} \Gamma(r/2)} \frac{1}{(1 + t^2/r)^{(r+1)/2}}, \quad -\infty < t < \infty.$$

If interested, please refer to textbook for proof. (You are not expected to know how to do it).

Review: CI for mean



Given the following sample: {16, 12, 18, 13, 21, 15, 8, 17}

$$n=8$$

Construct a 92% confidence interval for the true mean.

$$\bar{x} = 15, \quad s^2 = \frac{\sum(x_i - \bar{x})^2}{n-1} = \frac{112}{7} = 16, \quad s=4$$

$$\alpha = 0.08, \alpha/2 = 0.04$$

$$df = n - 1 = 7, \quad t_{7, 0.04} = 2.046$$

$$\bar{x} \pm t_{\alpha/2} \cdot \frac{s}{\sqrt{n}}$$

$$CI: 15 \pm 2.046 \cdot \frac{4}{\sqrt{8}} = (12.107, 17.893)$$

x	$x - \bar{x}$	$(x - \bar{x})^2$
16	1	1
12	-3	9
18	3	9
13	-2	4
21	6	36
15	0	0
8	-7	49
17	2	4
	0	112

notes

General Form of Confidence Interval

Estimate \pm (Critical Value * SE of estimate)

e.g. if σ is known:

$$\bar{x} \pm z_{\alpha/2} * \frac{\sigma}{\sqrt{n}}$$

if σ is unknown:

$$\bar{x} \pm t_{n-1, \alpha/2} * \frac{s}{\sqrt{n}}$$

Required Sample Size

$$\bar{x} \pm z_{\alpha/2} * \frac{\sigma}{\sqrt{n}}$$

$$\varepsilon = z_{\alpha/2} * \frac{\sigma}{\sqrt{n}}$$


$$n = \left[\frac{z_{\alpha/2} * \sigma}{\varepsilon} \right]^2$$

Required Sample Size

How many test runs of an automobile are required for determining its average miles-per-gallon rating on the highway to within 0.5 miles per gallon with 99% confidence? Suppose that the variance of the population in mpg² is 6.25.

$$n = \left[\frac{z_{\alpha/2} \cdot \sigma}{\varepsilon} \right]^2$$

$$\alpha = 0.01 \quad z_{\alpha/2} = 2.576$$



$$n =$$

$$= 165.89$$

$$\boxed{n = 166}$$

Round up!

last example, found $s^2 = 16$ ← point estimate

Confidence Interval for σ^2

Now we want to make a confidence interval for σ^2 based on s^2 .

- The distribution of s^2 is not Normal.

- It can be shown that $\frac{(n-1)s^2}{\sigma^2} \sim \chi^2_{(n-1)}$
will not need to prove this

Proof of $\frac{(n-1)S^2}{\sigma^2} \sim \chi_{n-1}^2$

$$S^2 = \frac{1}{n-1} \sum (X_i - \bar{X})^2$$

Assume

1. \bar{X} (the sample mean) and S^2 are independent.
2. If $Z \sim N(0, 1)$ then $Z^2 \sim \chi^2(1)$.
3. If $X_i \sim \chi^2(1)$ and the X_i are independent then $\sum_{i=1}^n X_i \sim \chi^2(n)$.
4. A $\chi^2(n)$ random variable has the moment generating function $(1 - 2t)^{-n/2}$.

With some algebra, you can show, by adding $-\bar{X} + \bar{X}$ inside the parentheses and grouping appropriately, that $\sum_{i=1}^n (X_i - \mu)^2 = \sum_{i=1}^n (X_i - \bar{X})^2 + n(\bar{X} - \mu)^2$. Then, dividing through by σ^2 yields

$$\sum_{i=1}^n \left(\frac{X_i - \mu}{\sigma} \right)^2 = \sum_{i=1}^n \left(\frac{X_i - \bar{X}}{\sigma} \right)^2 + \left(\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \right)^2.$$

Denote these expressions by U , V , and W , respectively, so that the formula reads $U = V + W$. By facts (2) and (3) above, $U \sim \chi^2(n)$ and $W \sim \chi^2(1)$. Also, $V = \frac{(n-1)S^2}{\sigma^2}$.

Since \bar{X} and S^2 are independent, so are V and W . Thus $M_U(t) = M_V(t)M_W(t)$, where $M_X(t)$ denotes the moment generating function of the random variable X . By fact (4) above, this says that

$$\frac{1}{(1 - 2t)^{n/2}} = M_V(t) \frac{1}{(1 - 2t)^{1/2}}.$$

$$M_V(t) = \frac{1}{(1 - 2t)^{(n-1)/2}}, \quad \text{so} \quad V \sim \chi_{(n-1)}^2$$

notes

$$\frac{(n-1)S^2}{\sigma^2} \sim \chi^2_{(n-1)} \text{ e.g., } \chi^2_{(10)}$$

Derivation of CI for Variance

$$\frac{(n-1)S^2}{\sigma^2} \sim \chi^2_{(n-1)}$$

$f(x)$



$$P\left[\chi^2_{n-1, 1-\alpha/2} < \frac{(n-1)S^2}{\sigma^2} < \chi^2_{n-1, \alpha/2}\right] = 1-\alpha$$

on the next slide ↴

$$P\left[\chi^2_{n-1, 1-\alpha/2} < \frac{1}{\sigma^2} < \chi^2_{n-1, \alpha/2}\right] = 1-\alpha \Rightarrow P\left[\frac{\frac{(n-1)S^2}{\chi^2_{n-1, \alpha/2}}}{\chi^2_{n-1, 1-\alpha/2}} < \sigma^2 < \frac{(n-1)S^2}{\chi^2_{n-1, 1-\alpha/2}}\right] = 1-\alpha$$

Confidence Interval for σ^2

Confidence Interval for σ^2 :

$$\left(\frac{(n-1) \cdot s^2}{\chi_{\alpha/2}^2}, \frac{(n-1) \cdot s^2}{\chi_{1-\alpha/2}^2} \right)$$

CI for Variance

Confidence Interval for σ^2 :

$$\left(\frac{(n-1) \cdot s^2}{\chi^2_{\alpha/2}}, \frac{(n-1) \cdot s^2}{\chi^2_{1-\alpha/2}} \right)$$

Given the following sample: {16, 12, 18, 13, 21, 15, 8, 17}


Construct a 95% CI for the true variance.

$s^2=16$ $\alpha = 0.05$ $\alpha/2 = 0.025$ $1 - \alpha/2 = 0.975$ $df = 7$


$\chi^2_{(7, 0.025)} = 16.013$ $\chi^2_{(7, 0.975)} = 1.690$

$$\left(\frac{(8-1)16}{16.013}, \frac{(8-1)16}{1.690} \right) = (6.996, 66.272)$$

$df = 7$
 $x = 16.01276$ $P(X > x) = 0.025$



$df = 7$
 $x = 1.68987$ $P(X > x) = 0.975$



1-sided Confidence Intervals for σ^2

95% confidence lower bound:

$$\left(\frac{(n-1)s^2}{\chi^2_{(df, \alpha)}}, \infty \right)$$

95% confidence upper bound:

$$\left(-\infty, \frac{(n-1)s^2}{\chi^2_{(df, 1-\alpha)}} \right)$$

notes