

# Normal Mixture Revisited

5361 Homework 7

*Qinxiao Shi* \*

27 October 2018

## 1 Normal Mixture Revisited

Consider a normal mixture, prior for  $\mu_1, \mu_2 \sim N(0, 10^2)$ , and the prior for  $1/\sigma_1^2, 1/\sigma_2^2 \sim \Gamma(0.5, 10)$ . Further, all the priors are independent.

First, the mixture normal distribution is

$$f(x) = \delta N(\mu_1, \sigma_1^2) + (1 - \delta) N(\mu_2, \sigma_2^2)$$

Since  $\frac{1}{\sigma_1^2}, \frac{1}{\sigma_2^2} \sim \Gamma(0.5, 10)$ ,  $\sigma_1^2, \sigma_2^2 \sim \text{Inv-Gamma}(0.5, 10)$ . So the posterior density is

$$\pi(\theta|\mathbf{x}) \propto \prod_{i=1}^n f(x_i) \times \frac{1}{\Gamma(0.5)10^{0.5}} \left(\frac{1}{\sigma_1^2}\right)^{1.5} e^{-\frac{1}{10\sigma_1^2}} \times \frac{1}{\Gamma(0.5)10^{0.5}} \left(\frac{1}{\sigma_2^2}\right)^{1.5} e^{-\frac{1}{10\sigma_2^2}} \times \frac{1}{10\sqrt{2\pi}} e^{-\frac{\mu_1^2}{2 \times 10^2}} \times \frac{1}{10\sqrt{2\pi}} e^{-\frac{\mu_2^2}{2 \times 10^2}}$$

where  $\theta = (\delta, \mu_1, \mu_2, \sigma_1^2, \sigma_2^2)$ .

```
library('invgamma')
library('HI')
delta <- 0.7
n <- 100
set.seed(123)
u <- rbinom(n, prob = delta, size = 1)
x <- rnorm(n, ifelse(u == 1, 7, 10), 0.5)

logll <- function(theta, x){
  delta <- theta[1]
  mu1 <- theta[2]
  mu2 <- theta[3]
  sg1 <- theta[4]
  sg2 <- theta[5]

  sum(log(delta*dnorm(x, mu1, sg1^0.5)+(1-delta)*dnorm(x, mu2, sg2^0.5)))
}

loglr <- function(theta){
  delta <- theta[1]
  mu1 <- theta[2]
```

---

\*qinxiao.shi@uconn.edu

```

mu2 <- theta[3]
sg1 <- theta[4]
sg2 <- theta[5]

return(dnorm(mu1, 0, 10, log = T)+dnorm(mu2, 0, 10, log = T)+
       dinvgamma(sg1, shape = 0.5, scale = 10, log = T)+dinvgamma(sg2, shape = 0.5, scale = 10, log = T))
}

logpost <- function(theta, x) {
  delta <- theta[1]
  mu1 <- theta[2]
  mu2 <- theta[3]
  sg1 <- theta[4]
  sg2 <- theta[5]

  return(logll(theta, x)+loglr(theta))
}

mymcmc <- function(niter, thetaInit, x, nburn= 100) {
  p <- length(thetaInit)
  thetaCurrent <- thetaInit
  logFC <- function(th, idx) {
    theta <- thetaCurrent
    theta[idx] <- th
    logpost(theta, x)
  }
  out <- matrix(thetaInit, niter, p, byrow = TRUE)
  ## Gibbs sampling
  for (i in 2:niter) {
    for (j in 1:p) {
      if (j == 1 | j == 4 | j == 5){
        out[i, j] <- thetaCurrent[j] <-
          HI::arms(thetaCurrent[j], logFC,
                   function(x, idx) ((x > 0) * (x < 1)),
                   1, idx = j)
      } else if (j == 2 | j == 3) {
        out[i, j] <- thetaCurrent[j] <-
          HI::arms(thetaCurrent[j], logFC,
                   function(x, idx) ((x > -50) * (x < 50)),
                   1, idx = j)
      }
    }
  }
  out[-(1:nburn), ]
}


```


```


niter <- 3000
nburn <- 1000
thetaInit <- c(0.3, 10, 7, 0.25, 0.24)
sim <- mymcmc(niter, thetaInit, x)


par(mfrow = c(1,2))
for (i in 1:length(thetaInit)){
  plot(ts(sim[,i]))
  hist(sim[,i], main = expression(paste("Histogram of ith ", theta)),
       xlab = expression(paste("ith ", theta)))
}


```











So the estimated values of  $\theta$  are

$$\delta = 0.3, \mu_1 = 10, \mu_2 = 7, \sigma_1^2 = 0.25, \sigma_2^2 = 0.24$$