

Initial Assessments of Covariate-Adjusted Average Treatment Effect Estimation under Misspecified Multiple Imputation Models

Gwendolyn Jacobson

Jerome Reiter

Duke University

September 21, 2026

- Introduction and motivation
- Model and simulation specifications
- Preliminary results
- Future directions

Causal Inference Review

Let $i \in \{1, \dots, n\}$ be an experimental unit.

- $T_i \in \{0, 1\}$: treatment status (control, treated)
- $Y_i(0)$: potential outcome under control
- $Y_i(1)$: potential outcome under treatment

The observed outcome is given by

$$Y_i = T_i Y_i(1) + (1 - T_i) Y_i(0).$$

The estimand is the **average treatment effect (ATE)**:

$$\tau = \frac{1}{n} \sum_{i=1}^n (Y_i(1) - Y_i(0)).$$

Let n_1 be the number of units under treatment and $n_0 = n - n_1$ the number under control.

Let n_1 be the number of units under treatment and $n_0 = n - n_1$ the number under control.

Difference-in-Means:

$$\hat{\tau}^{\text{unadj}} = \frac{1}{n_1} \sum_{T_i=1} Y_i - \frac{1}{n_0} \sum_{T_i=0} Y_i$$

Let n_1 be the number of units under treatment and $n_0 = n - n_1$ the number under control.

Difference-in-Means:

$$\hat{\tau}^{\text{unadj}} = \frac{1}{n_1} \sum_{T_i=1} Y_i - \frac{1}{n_0} \sum_{T_i=0} Y_i$$

Outcome Regression:

$$\hat{\tau}^{\text{adj}} = \sum_{i=1}^n \{ \hat{\mu}_1(X_i) - \hat{\mu}_0(X_i) \},$$

where $\hat{\mu}_1$ and $\hat{\mu}_0$ are the fitted values of the outcome means.

Causal Inference Review

Let n_1 be the number of units under treatment and $n_0 = n - n_1$ the number under control.

Difference-in-Means:

$$\hat{\tau}^{\text{unadj}} = \frac{1}{n_1} \sum_{T_i=1} Y_i - \frac{1}{n_0} \sum_{T_i=0} Y_i$$

Outcome Regression:

$$\hat{\tau}^{\text{adj}} = \sum_{i=1}^n \{\hat{\mu}_1(X_i) - \hat{\mu}_0(X_i)\},$$

where $\hat{\mu}_1$ and $\hat{\mu}_0$ are the fitted values of the outcome means.

Note: $\hat{\tau}^{\text{adj}}$ frequently taken to be OLS estimate for T_i coefficient.

Setting: An analyst want to make inference about the effect of a treatment in a completely randomized experiment and has:

- Observed outcomes
- Treatment assignments
- *Pre-treatment* covariates

Randomization is “good” on average, but **regression adjustment** can help.

- Combat covariate imbalance [Ding, 2024]
- Improve estimator efficiency

Problem: Some units' pre-treatment covariates may be **missing**.

Problem: Some units' pre-treatment covariates may be **missing**.

- Missingness completely at random (MCAR): missingness *independent* of missing and observed values

Problem: Some units' pre-treatment covariates may be **missing**.

- Missingness completely at random (MCAR): missingness *independent* of missing and observed values
- Missingness at random (MAR): missingness independent of missing values *conditional* on observed

Problem: Some units' pre-treatment covariates may be **missing**.

- Missingness completely at random (MCAR): missingness *independent* of missing and observed values
- Missingness at random (MAR): missingness independent of missing values *conditional* on observed
- Missingness not at random (MNAR): missingness *dependent* on missing values

Problem: Some units' pre-treatment covariates may be **missing**.

- Missingness completely at random (MCAR): missingness *independent* of missing and observed values
- Missingness at random (MAR): missingness independent of missing values *conditional* on observed
- Missingness not at random (MNAR): missingness *dependent* on missing values

Suppose a binary covariate X_1 exhibits MCAR.


Problem: Some units' pre-treatment covariates may be **missing**.

- Missingness completely at random (MCAR): missingness *independent* of missing and observed values
- Missingness at random (MAR): missingness independent of missing values *conditional* on observed
- Missingness not at random (MNAR): missingness *dependent* on missing values


Suppose a binary covariate X_1 exhibits MCAR.

Goal: Overcome missingness with **multiple imputation (MI)** to perform regression adjustment.


Multiple Imputation



Multiple Imputation



Multiple Imputation



Multiple Imputation

Why multiply impute?

- Use complete data methods easily
- Quantify imputation uncertainty
- Excellent for sensitivity analysis

For $n = 10,000$ and $n_1 = 5,000$, we generate data as follows.

Treatment and Covariates:

$$\mathbb{P}(\mathbf{T} = \mathbf{t}) = \frac{1}{\binom{n}{n_1}} \quad t_i \in \{0, 1\}, \quad \|\mathbf{t}\|_1 = n_1$$

$$X_{1i} \stackrel{\text{i.i.d.}}{\sim} \text{Bernoulli}(0.5)$$

$$X_{2i} \stackrel{\text{i.i.d.}}{\sim} \text{Bernoulli}(0.5)$$

Outcomes:

$$Y_i(0) \mid X_{1i}, X_{2i} \stackrel{\text{ind.}}{\sim} \mathcal{N}(\beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \beta_{12} X_{1i} X_{2i}, \sigma^2)$$

$$Y_i(1) = Y_i(0) + \tau$$

$$\Downarrow$$

$$Y_i \mid T_i, X_{1i}, X_{2i} \stackrel{\text{ind.}}{\sim} \mathcal{N}(\beta_0 + \tau T_i + \beta_1 X_{1i} + \beta_2 X_{2i} + \beta_{12} X_{1i} X_{2i}, \sigma^2)$$

Missingness: Let $R_i = 1$ for observed X_{1i} and $R_i = 0$ for missing X_{1i} :

$$R_i \mid X_{1i} \stackrel{\text{i.i.d.}}{\sim} \text{Bernoulli}(0.3)$$

Data-Generating Model

Outcomes:

$$Y_i(0) \mid X_{1i}, X_{2i} \stackrel{\text{i.i.d.}}{\sim} N(\beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \beta_{12} X_{1i} X_{2i}, \sigma^2)$$

$$Y_i(1) = Y_i(0) + \tau$$

$$\Downarrow$$

$$Y_i \mid T_i, X_{1i}, X_{2i} \stackrel{\text{i.i.d.}}{\sim} N(\beta_0 + \tau T_i + \beta_1 X_{1i} + \beta_2 X_{2i} + \beta_{12} X_{1i} X_{2i}, \sigma^2)$$

Missingness: Let $R_i = 1$ for observed X_{1i} and $R_i = 0$ for missing X_{1i} :

$$R_i \mid X_{1i} \stackrel{\text{i.i.d.}}{\sim} \text{Bernoulli}(0.3)$$

Sensitivity analysis:

$$\tau, \beta_1, \beta_2 \in \{0.5, 7.5\} \quad (\text{"low" and "high" values})$$

$$\beta_{12} \in \{0, 0.5, 1, \dots, 9.5, 10\} \quad (\text{interaction strength})$$

Imputation Mechanism

First iteration:

- 1 Draw $p_X \sim \text{Beta}(1, 1)$.
- 2 Impute missing $X_{1i} \mid p_X \sim \text{Bernoulli}(p_X)$.
- 3 Calculate OLS coefficient and variance estimates.

Imputation Mechanism

First iteration:

- 1 Draw $p_X \sim \text{Beta}(1, 1)$.
- 2 Impute missing $X_{1i} \mid p_X \sim \text{Bernoulli}(p_X)$.
- 3 Calculate OLS coefficient and variance estimates.

Subsequent iterations:

- 1 Draw

$$p_X \sim \text{Beta} \left(1 + \sum_{R_i=0} \mathbb{I}[X_i = 1], 1 + \sum_{R_i=0} \mathbb{I}[X_i = 0] \right).$$

- 2 Impute missing X_{1i} based on p_X draw, imputation model, and previous OLS estimates.
- 3 Calculate new OLS coefficient and variance estimates.

Regression/MI Model Combinations

Recall: The regression model produces the ATE estimates, while the MI model completes the data set ahead of the regression.

Regression Model	MI Model	Imputation Contributors
Full	Full	$p_X, Y, T, X_1, X_2, X_1X_2$
	No interaction	p_X, Y, T, X_1, X_2
	X_1 only	p_X, Y, T, X_1
No interaction	No interaction	p_X, Y, T, X_1, X_2

Regression/MI Model Combinations

Recall: The regression model produces the ATE estimates, while the MI model completes the data set ahead of the regression.

Regression Model	MI Model	Imputation Contributors
Full	Full	$p_X, Y, T, X_1, X_2, X_1X_2$
	No interaction	p_X, Y, T, X_1, X_2
	X_1 only	p_X, Y, T, X_1
No interaction	No interaction	p_X, Y, T, X_1, X_2

For each missing X_{1i} , calculate

$$\mathbb{P}(X_{1i} = 1 \mid \cdots) = \frac{w_1}{w_1 + w_0},$$

where w_1 and w_0 are based on p_X and the OLS estimates.

Regression/ML Model Combinations

Full ML:

$$w_1 = p_X \cdot \text{dnorm} \left(Y_i; \hat{\beta}_0 + \hat{\tau}^{\text{adj}} T_i + \hat{\beta}_1 + \left(\hat{\beta}_2 + \hat{\beta}_{12} \right) X_{2i}, \hat{\sigma}^2 \right)$$

$$w_0 = (1 - p_X) \cdot \text{dnorm} \left(Y_i; \hat{\beta}_0 + \hat{\tau}^{\text{adj}} T_i + \hat{\beta}_2 X_{2i}, \hat{\sigma}^2 \right)$$

No-interaction ML:

$$w_1 = p_X \cdot \text{dnorm} \left(Y_i; \hat{\beta}_0 + \hat{\tau}^{\text{adj}} T_i + \hat{\beta}_1 + \hat{\beta}_2 X_{2i}, \hat{\sigma}^2 \right)$$

$$w_0 = (1 - p_X) \cdot \text{dnorm} \left(Y_i; \hat{\beta}_0 + \hat{\tau}^{\text{adj}} T_i + \hat{\beta}_2 X_{2i}, \hat{\sigma}^2 \right)$$

X_1 -only ML:

$$w_1 = p_X \cdot \text{dnorm} \left(Y_i; \hat{\beta}_0 + \hat{\tau}^{\text{adj}} T_i + \hat{\beta}_1, \hat{\sigma}^2 \right)$$


$$w_0 = (1 - p_X) \cdot \text{dnorm} \left(Y_i; \hat{\beta}_0 + \hat{\tau}^{\text{adj}} T_i, \hat{\sigma}^2 \right)$$

Simulation Set-up


- ① Generate $n = 10,000$ experimental units.
- ② Assign $n_1 = 5,000$ to treatment.
- ③ Induce missingness in X_1 at random ($p = 0.3$).
- ④ Generate outcomes for each combination of coefficients.
 - ① Calculate unadjusted ATEs.
 - ② Run imputation mechanism and calculate ATEs for all regression-MI models across several iterations.

Repeat entire process across several seeds and calculate MSEs.


High ATE Results



Low X_1 Coefficient Results



High X_1 Coefficient Results



Summary and Future Directions

- “Best” MI models by others
 - Need more simulations
- Examine broader range of coefficient values
- Consider alternative settings
 - Covariate dependence structure
 - Transformed data
- Compare to additional estimators
 - Centered covariates [Lin, 2013]
 - Missingness indicator method [Zhao and Ding, 2024], [Zhao et al., 2023]
- Identify diagnostics for guidance



Ding, P. (2024).

A First Course in Causal Inference.

CRC Press.



Lin, W. (2013).

Agnostic notes on regression adjustments to experimental data: reexamining freedman's critique.

The Annals of Applied Statistics, 7(1):295–318.



Zhao, A. and Ding, P. (2024).

To adjust or not to adjust? estimating the average treatment effect in randomized experiments with missing covariates.

Journal of the American Statistical Association,
119(545):450–460.




Zhao, A., Ding, P., and Li, F. (2023).


Covariate adjustment in randomized experiments with missing outcomes and covariates.

arXiv preprint arXiv:2311.10877v2.

Appendix: Low ATE Results



Appendix: Low X_2 Coefficient Results



Appendix: High X_2 Coefficient Results

