
# Recap general linear model

Lieven Clement


## Breast cancer example


- part of study https://doi.org/10.1093/jnci/djj052)
- Histologic grade in breast cancer clinically prognostic. Association of histologic grade on expression of KPNA2 gene that is known to be associated with poor BC prognosis.
- Population: all current and future breast cancer patients


---


POPULATION
Effect of hist. grade in population



POPULATION
Effect of hist. grade in population
EXP. DESIGN


POPULATION
Effect of hist. grade in population
EXP. DESIGN
SAMPLE
Effect of hist. grade in sample


POPULATION
Effect of hist. grade in population
EXP. DESIGN
SAMPLE
Effect of hist. grade in sample
DATA EXPLORATION


POPULATION
Effect of hist. grade in population
EXP. DESIGN
ESTIMATION &
INFERENCE
SAMPLE
Effect of hist. grade in sample
DATA EXPLORATION

---




## Data Exploration

### Import

```
library(tidyverse)
```

```
-- Attaching core tidyverse packages ------------------------ tidyverse 2.0.0 --
v dplyr     1.2.1     v readr     2.2.0
v forcats   1.0.1     v stringr   1.6.0
v ggplot2   4.0.3     v tibble    3.3.1
v lubridate 1.9.5     v tidyr     1.3.2
v purrr     1.2.2
-- Conflicts ------------------------------------------ tidyverse_conflicts() --
x dplyr::filter() masks stats::filter()
x dplyr::lag()    masks stats::lag()
i Use the conflicted package (<http://conflicted.r-lib.org/>) to force all conflicts to beco
```

```
gene <- read.table("https://raw.githubusercontent.com/statOmics/SGA21/master/data/kpna2.txt"
head(gene)
```

```
  grade node     gene
1     3    1 367.8179
2     3    1 590.3576
3     1    1 346.6583
4     1    1 258.4455
5     1    0 153.8416
6     3    0 643.6799
```

We will transform the variable grade and node to a factor

```
gene$grade <- as.factor(gene$grade)
gene$node <- as.factor(gene$node)
```


### Summary statistics

```
geneSum <- gene |>
  group_by(grade) |>
  summarize(mean = mean(gene),
            sd = sd(gene),
            n=length(gene)
            ) |>
  mutate(se = sd/sqrt(n))
geneSum
```

```
# A tibble: 2 x 5
  grade  mean    sd     n    se
  <fct> <dbl> <dbl> <int> <dbl>
1 1      233.  65.5    12  18.9
2 3      561. 144.     12  41.4
```

### Visualisation

```
gene |>
  ggplot(aes(x=grade,y=gene)) +
  geom_boxplot(outlier.shape=NA) +
  geom_jitter()
```





We can also save the plots as objects for later use!


```
p1 <- gene |>
  ggplot(aes(x=grade,y=gene)) +
  geom_boxplot(outlier.shape=NA) +
  geom_jitter()

p2 <- gene |>
  ggplot(aes(sample=gene)) +
  geom_qq() +
  geom_qq_line() +
  facet_wrap(~grade)

p1
```





```
p2
```



### Research questions

Researchers want to assess the association of the histological grade on KPNA2 gene expression




---

### Estimation of effect size and standard error

```
effectSize <- tibble(
  delta = geneSum$mean[2]- geneSum$mean[1],
  seDelta = sqrt(sum(geneSum$se^2))
  )
effectSize
```


```
# A tibble: 1 x 2
  delta seDelta
```

```
  <dbl>   <dbl>
1  329.    45.5
```

## Statistical Inference

- Researchers want to assess the association of histological grade on KPNA2 gene expression
- Inference?

---




---

- Researchers want to assess the association of histological grade on KPNA2 gene expression
- Inference?
- testing + CI $ \rightarrow$ Assumptions

---

- In general we start from **alternative hypothesis** $H_A$: we want to show an association
- Gene expression of grade 1 and grade 3 patients is on average different
- But, we will assess it by falsifying the opposite:
- The average KPNA2 gene expression of grade 1 and grade 3 patients is equal

---

- How likely is it to observe an equal or more extreme association than the one observed in the sample when the null hypothesis is true?
- When we make assumptions about the distribution of our test statistic we can quantify this probability: **p-value**.
- If the p-value is below a significance threshold $\alpha$ we reject the null hypothesis

*We control the probability of a false positive result at the $\alpha$-level (type I error)*

- The p-value will only be calculated correctly if the underlying assumptions hold!


```
library(gridExtra)
```

```
Attaching package: 'gridExtra'


The following object is masked from 'package:dplyr':

    combine
```

```
p1
```




```
p2
```



```r
t.test(gene~grade,data=gene)
```

```
    Welch Two Sample t-test

data:  gene by grade
t = -7.2132, df = 15.384, p-value = 2.598e-06
alternative hypothesis: true difference in means between group 1 and group 3 is not equal to
95 percent confidence interval:
 -425.4218 -231.6751
sample estimates:
mean in group 1 mean in group 3
       232.5003        561.0487
```


```r
effectSize <- effectSize |>
  mutate(t.stat=delta/seDelta) |>
  mutate(p.value= pt(-abs(t.stat),21.352)*2)

effectSize
```

```
# A tibble: 1 x 4
  delta seDelta t.stat     p.value
  <dbl>   <dbl>  <dbl>       <dbl>
1  329.    45.5   7.21 0.000000376
```

- Intensities are often not normally distributed and have a mean variance relation
- Commonly log2-transformed
- Differences on log scale:

$$\log_2(B) - \log_2(A) = \log_2 \frac{B}{A} = \log_2 FC_{\frac{B}{A}}$$




---

### Log transformation


```
gene <- gene |>
  mutate(lgene = log2(gene))

p1 <- gene |>
  ggplot(aes(x=grade,y=lgene)) +
  geom_boxplot(outlier.shape=NA) +
  geom_jitter()


p2 <- gene |>
  ggplot(aes(sample=lgene)) +
  geom_qq() +
  geom_qq_line() +
```

```
  facet_wrap(~grade)

p1
```




```
p2
```




```
logtest <- t.test(lgene~grade,data=gene,var.equal=TRUE)
logtest
```

```
    Two Sample t-test

data:  lgene by grade
t = -8.0455, df = 22, p-value = 5.372e-08
alternative hypothesis: true difference in means between group 1 and group 3 is not equal to
95 percent confidence interval:
 -1.610148 -0.950178
sample estimates:
mean in group 1 mean in group 3
       7.808478        9.088641
```

```
log2FC <- logtest$estimate[2]-logtest$estimate[1]
log2FC
```


```
mean in group 3
       1.280163
```


```r
names(log2FC) <- "g3-g1"
2^log2FC
```

```
   g3-g1
2.428664
```

## Conclusion

There is an extremely significant association of the histological grade on the gene expression in tumor tissue. On average, the gene expression for the grade 3 patients is 2.43 times higher than the gene expression in grade 1 patients (95% CI [1.93, 3.05], $p << 0.001$).







---

The patients also differ in their lymph node status. Hence, we have a two factorial design: grade x lymph node status!!!

Solution??




---

## General Linear Model

How can we integrate multiple factors and continuous covariates in a linear model.

$$y_i = \beta_0 + \beta_1 x_{i,1} + \beta_2 x_{i,2} + \beta_{12} x_{i,1} x_{i,2} + \epsilon_i,$$

with

- $x_{i,1}$ a dummy variable for histological grade: $x_{i,1} = \begin{cases} 0 & \text{grade 1} \\ 1 & \text{grade 3} \end{cases}$
- $x_{i,2}$ a dummy variable for lymph node status: $x_{i,2} = \begin{cases} 0 & \text{lymph nodes were not removed} \\ 1 & \text{lymph nodes were removed} \end{cases}$
- $\epsilon_i$?

---

### Implementation in R

```
lm1 <- lm(gene~grade*node,data=gene)
summary(lm1)
```

```
Call:
lm(formula = gene ~ grade * node, data = gene)

Residuals:
     Min       1Q   Median       3Q      Max
-201.748  -53.294   -6.308   46.216  277.601


Coefficients:
             Estimate Std. Error t value Pr(>|t|)
(Intercept)    180.51      44.37   4.068   0.0006 ***
grade3         401.33      62.75   6.396 3.07e-06 ***
node1          103.98      62.75   1.657   0.1131
grade3:node1  -145.57      88.74  -1.640   0.1166
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1


Residual standard error: 108.7 on 20 degrees of freedom
Multiple R-squared:  0.7437,    Adjusted R-squared:  0.7052
F-statistic: 19.34 on 3 and 20 DF,  p-value: 3.971e-06
```


---


### Assumptions

```
plot(lm1)
```


Residuals vs Fitted
Residuals
200
0
−200
7
1
10
200
300
400
500
Fitted values
lm(gene ~ grade * node)


Q−Q Residuals
Standardized residuals
3
2
1
0
−2
7
1
10
−2
−1
0
1
2
Theoretical Quantiles
lm(gene ~ grade * node)


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
200
300
400
500
7
1
10
Fitted values
lm(gene ~ grade * node)



Constant Leverage:
Residuals vs Factor Levels
Standardized residuals
−2
0
1
2
3
7
10
1
grade :
1
3
Factor Level Combinations

### Breast cancer example

- Paper: https://doi.org/10.1093/jnci/djj052
- Histologic grade in breast cancer provides clinically important prognostic information. Two factors have to be considered: Histologic grade (grade 1 and grade 3) and lymph node status (0 vs 1). The researchers assessed gene expression of the KPNA2 gene a protein-coding gene associated with breast cancer and are mainly interested in the association of histological grade. Note, that the gene variable consists of background corrected normalized intensities obtained with a microarray platform. Upon log-transformation, they are known to be a good proxy for the log transformed concentration of gene expression product of the KPNA2 gene.
- Research questions and translate them towards model parameters (contrasts)?
- Make an R markdown file to answer the research questions

```
library(ExploreModelMatrix)
explMx <- VisualizeDesign(gene,designFormula = ~grade*node)
explMx$plotlist
```

```
[[1]]
```




You can also explore the model matrix interactively:

```
ExploreModelMatrix(gene,designFormula = ~grade*node)
```

---

## Linear regression in matrix form

### Scalar form

- Consider a vector of predictors $\mathbf{x} = (x_1, \ldots, x_p)^T$ and
- a real-valued response $Y$
- then the linear regression model can be written as

$$Y = f(\mathbf{x}) + \epsilon = \beta_0 + \sum_{j=1}^{p} x_j \beta_j + \epsilon$$

with i.i.d. $\epsilon \sim N(0, \sigma^2)$

### Matrix form

- $n$ observations $(\mathbf{x}_1, y_1) \ldots (\mathbf{x}_n, y_n)$
- Regression in matrix notation

$$\mathbf{Y} = \mathbf{X} +$$

with $\mathbf{Y} = \begin{bmatrix} y_1 \\ \vdots \\ y_n \end{bmatrix}$, $\mathbf{X} = \begin{bmatrix} 1 & x_{11} & \ldots & x_{1p} \\ \vdots & \vdots & & \vdots \\ 1 & x_{n1} & \ldots & x_{np} \end{bmatrix}$, $= \begin{bmatrix} \beta_0 \\ \vdots \\ \beta_p \end{bmatrix}$ and $= \begin{bmatrix} \epsilon_1 \\ \vdots \\ \epsilon_n \end{bmatrix}$

### Least Squares (LS)

- Minimize the residual sum of squares

$$\begin{aligned} RSS() &= \sum_{i=1}^{n} e_i^2 \\ &= \sum_{i=1}^{n} \left( y_i - \beta_0 - \sum_{j=1}^{p} x_{ij} \beta_j \right)^2 \end{aligned}$$

- or in matrix notation

$$\begin{aligned} RSS() &= (\mathbf{Y}-\mathbf{X})^T(\mathbf{Y}-\mathbf{X}) \\ &= \|\mathbf{Y}-\mathbf{X}\|_2^2 \end{aligned}$$

with the $L_2$-norm of a $p$-dim. vector $v$ $\|\mathbf{v}\| = \sqrt{v_1^2 + ... + v_p^2}$ $\rightarrow \hat{} = \text{argmin}_\beta \|\mathbf{Y}-\mathbf{X}\|_2^2$

---

### Minimize RSS

$$\begin{aligned} \frac{\partial RSS}{\partial} &= \mathbf{0} \\ \frac{(\mathbf{Y}-\mathbf{X})^T(\mathbf{Y}-\mathbf{X})}{\partial} &= \mathbf{0} \\ -2\mathbf{X}^T(\mathbf{Y}-\mathbf{X}) &= \mathbf{0} \\ \mathbf{X}^T\mathbf{X} &= \mathbf{X}^T\mathbf{Y} \\ \hat{} &= (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{Y} \end{aligned}$$

---

### Geometric Interpretation

### Toy dataset

We will illustrate this on a toy dataset

```
library(tidyverse)
data <- data.frame(x=1:3,y=c(1,2,2))
data
```

```
  x y
1 1 1
2 2 2
3 3 2
```

### Matrix form for toy dataset

We can also write this in matrix form

$$\mathbf{Y} = \mathbf{X}\beta + \epsilon$$

with

$$\mathbf{Y} = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}, \quad \mathbf{X} = \begin{bmatrix} 1 & 1 \\ 1 & 2 \\ 1 & 3 \end{bmatrix}, \quad \beta = \begin{bmatrix} \beta_0 \\ \beta_1 \end{bmatrix} \quad \text{and} \quad \epsilon = \begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \epsilon_3 \end{bmatrix}$$

### Classical interpretation


Model fit and predictions based on the toy dataset

```
lm1 <- lm(y~x,data)
data$yhat <- lm1$fitted

data |>
  ggplot(aes(x,y)) +
  geom_point() +
  ylim(0,4) +
  xlim(0,4) +
  stat_smooth(method = "lm", color = "red", fullrange = TRUE) +
  geom_point(aes(x=x, y =yhat), pch = 2, size = 3, color = "red") +
  geom_segment(data = data, aes(x = x, xend = x, y = y, yend = yhat), lty = 2 )
```

```
`geom_smooth()` using formula = 'y ~ x'
```

```
Warning in max(ids, na.rm = TRUE): no non-missing arguments to max; returning
-Inf
```




### Projection


There is also another picture to regression:

- Instead of plotting each observation $i = 1 \dots n$ as a data-point in $\mathbb{R}^p$ with dimensions $1 \dots p$ for every variable/feature that is recorded for each observation
- We can also plot $\mathbf{Y}$, $\hat{\mathbf{Y}}$ and each column of $\mathbf{X}$: $\mathbf{X}_j$ with $j = 1 \dots p$ as a vector in $\mathbb{R}^n$ with dimensions $1 \dots n$ for every observation.
- In this representation linear regression can be interpreted as a projection of the vector $\mathbf{Y}$ onto the subspace of $\mathbb{R}^n$ that is spanned by the vectors for the predictors $\mathbf{X}_1 \dots \mathbf{X}_p$.
- The space $\mathbf{X}_1 \dots \mathbf{X}_p$ is also referred to as the column space of $\mathbf{X}$, the space that consists of all linear combinations of the vectors of the predictors or columns $\mathbf{X}_1 \dots \mathbf{X}_p$.

### Intermezzo: Projection of vector on X and Y axis

$$\mathbf{e} = \begin{bmatrix} e_1 \\ e_2 \end{bmatrix}, \mathbf{u}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

```
Warning: Using `size` aesthetic for lines was deprecated in ggplot2 3.4.0.
i Please use `linewidth` instead.
```




1. Projection of error on x-axis

$$\begin{aligned} \mathbf{u}_1^T\mathbf{e} &= \|\mathbf{u}_1\|_2\|\mathbf{e}_1\|_2 \cos < \mathbf{u}_1, \mathbf{e}_1 > \\ &= \begin{bmatrix} 1 & 0 \end{bmatrix} \begin{bmatrix} e_1 \\ e_2 \end{bmatrix} \\ &= 1 \times e_1 + 0 \times e_2 \\ &= e_1 \end{aligned}$$

2. Projection of error on y-axis

$$\begin{aligned} \mathbf{u}_2^T\mathbf{e} &= \begin{bmatrix} 0 & 1 \end{bmatrix} \begin{bmatrix} e_1 \\ e_2 \end{bmatrix} \\ &= 0 \times e_1 + 1 \times e_2 \\ &= e_2 \end{aligned}$$

3. Projection of error on itself

$$\begin{aligned}\mathbf{e}^T\mathbf{e} &= \begin{bmatrix} e_1 & e_2 \end{bmatrix}\begin{bmatrix} e_1 \\ e_2 \end{bmatrix} \\ &= e_1^2 + e_2^2 \\ &= \|e\|_2^2 \rightarrow \text{ Pythagorean theorem}\end{aligned}$$

---

**Interpretation of least squares as a projection**

Fitted values:

$$\begin{aligned}\hat{\mathbf{Y}} &= \mathbf{X}\hat{\beta} \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{Y} \\ &= \mathbf{H}\mathbf{Y}\end{aligned}$$

with $\mathbf{H}$ the projection matrix also referred to as the hat matrix.

```
X <- model.matrix(~x,data)
X
```

```
  (Intercept) x
1           1 1
2           1 2
3           1 3
attr(,"assign")
[1] 0 1
```

```
XtX <- t(X)%*%X
XtX
```

```
            (Intercept)  x
(Intercept)           3  6
x                     6 14
```

```
XtXinv <- solve(t(X)%*%X)
XtXinv
```

```
            (Intercept)    x
(Intercept)    2.333333 -1.0
x             -1.000000  0.5
```

```
H <- X %*% XtXinv %*% t(X)
H
```

```
           1         2          3
1  0.8333333 0.3333333 -0.1666667
2  0.3333333 0.3333333  0.3333333
3 -0.1666667 0.3333333  0.8333333
```

```
Y <- data$y
Yhat <- H%*%Y
Yhat
```

```
      [,1]
1 1.166667
2 1.666667
3 2.166667
```

- We can also interpret the fit as the projection of the $n \times 1$ vector $\mathbf{Y}$ on the column space of the matrix $\mathbf{X}$.
- So each column in $\mathbf{X}$ is also an $n \times 1$ vector.
- For the toy example n=3 and p=2. The other picture to linear regression is to consider $X_0$, $X_1$ and $Y$ as vectors in the space of the data $\mathbb{R}^n$, here $\mathbb{R}^3$ because we have three data points. So the column space of X is a plane in the three dimensional space.

$$\hat{\mathbf{Y}} = \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{Y}$$

1. Plane spanned by column space:

```
originRn <- data.frame(X1=0,X2=0,X3=0)
data$x0 <- 1
dataRn <- data.frame(t(data))

library(plotly)
```

```
Attaching package: 'plotly'

The following object is masked from 'package:ggplot2':

    last_plot
```

```
The following object is masked from 'package:stats':


    filter

The following object is masked from 'package:graphics':

    layout
```

```
p1 <- plot_ly(
    originRn,
    x = ~ X1,
    y = ~ X2,
    z= ~ X3, name="origin") |>
  add_markers(type="scatter3d") |>
  layout(
    scene = list(
      aspectmode="cube",
      xaxis = list(range=c(-4,4)), yaxis = list(range=c(-4,4)), zaxis = list(range=c(-4,4))
      )
    )
p1 <- p1 |>
  add_trace(
    x = c(0,1),
    y = c(0,0),
    z = c(0,0),
    mode = "lines",
    line = list(width = 5, color = "grey"),
    type="scatter3d",
    name = "obs1") |>
  add_trace(
    x = c(0,0),
    y = c(0,1),
    z = c(0,0),
    mode = "lines",
    line = list(width = 5, color = "grey"),
    type="scatter3d",
    name = "obs2") |>
  add_trace(
    x = c(0,0),
    y = c(0,0),
    z = c(0,1),
    mode = "lines",
```

```
    line = list(width = 5, color = "grey"),
    type="scatter3d",
    name = "obs3") |>
  add_trace(
    x = c(0,1),
    y = c(0,1),
    z = c(0,1),
    mode = "lines",
    line = list(width = 5, color = "black"),
    type="scatter3d",
    name = "X1") |>
    add_trace(
    x = c(0,1),
    y = c(0,2),
    z = c(0,3),
    mode = "lines",
    line = list(width = 5, color = "black"),
    type="scatter3d",
    name = "X2")
p1
```


X3
X2
X1

2. Vector of Y:


Actual values of $\mathbf{Y}$:

```
data$y
```

```
[1] 1 2 2
```

$$\mathbf{Y} = \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}$$

```
p2 <- p1 |>
  add_trace(
    x = c(0,Y[1]),
    y = c(0,Y[2]),
    z = c(0,Y[3]),
    mode = "lines",
    line = list(width = 5, color = "red"),
    type="scatter3d",
    name = "Y")
p2
```


4
3
2
1
0
−1
−2
−3
−4
X3
−3
−2
−1
0
1
2
3
X2
−1
0
1
2
3
X1

3. Projection of Y onto column space


Actual values of fitted values $\hat{\mathbf{Y}}$:

```
data$yhat
```

```
[1] 1.166667 1.666667 2.166667
```

$$\mathbf{Y} = \begin{bmatrix} 1.1666667 \\ 1.6666667 \\ 2.1666667 \end{bmatrix}$$

```
p2 <- p2 |>
  add_trace(
    x = c(0,Yhat[1]),
    y = c(0,Yhat[2]),
    z = c(0,Yhat[3]),
    mode = "lines",
    line = list(width = 5, color = "orange"),
    type="scatter3d",
    name="Yhat") |>
    add_trace(
    x = c(Y[1],Yhat[1]),
    y = c(Y[2],Yhat[2]),
    z = c(Y[3],Yhat[3]),
    mode = "lines",
    line = list(width = 5, color = "red", dash="dash"),
    type="scatter3d",
    name="Y -> Yhat"
    )
p2
```


X3
X2
X1

$\mathbf{Y}$ is projected in the column space of $\mathbf{X}$! spanned by the columns.

#### How does this projection works?

$$\begin{aligned}\hat{\mathbf{Y}} &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{Y} \\ &= \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1/2}(\mathbf{X}^T\mathbf{X})^{-1/2}\mathbf{X}^T\mathbf{Y} \\ &= \mathbf{U}\mathbf{U}^T\mathbf{Y}\end{aligned}$$

- $\mathbf{U}$ is a new orthonormal basis in $\mathbb{R}^2$, a subspace of $\mathbb{R}^3$
- The space spanned by U and X is the column space of X, e.g. it contains all possible linear combinantions of X. $\mathbf{U}^t\mathbf{Y}$ is the projection of Y on this new orthonormal basis

```
eigenXtX <- eigen(XtX)
XtXinvSqrt <- eigenXtX$vectors %*%diag(1/eigenXtX$values^.5)%*%t(eigenXtX$vectors)
U <- X %*% XtXinvSqrt
```

- $\mathbf{U}$ orthonormal basis

```
U
```

```
        [,1]        [,2]
1  0.9116067 -0.04802616
2  0.3881706  0.42738380
3 -0.1352655  0.90279376
```

```
t(U)%*%U
```

```
              [,1]          [,2]
[1,] 1.000000e+00 2.915205e-16
[2,] 2.915205e-16 1.000000e+00
```

- $\mathbf{U}\mathbf{U}^T$ equals projection matrix

```
U%*%t(U)
```


```
           1         2          3
1  0.8333333 0.3333333 -0.1666667
2  0.3333333 0.3333333  0.3333333
3 -0.1666667 0.3333333  0.8333333
```

```
H
```

```
           1         2          3
1  0.8333333 0.3333333 -0.1666667
2  0.3333333 0.3333333  0.3333333
3 -0.1666667 0.3333333  0.8333333
```


```
p3 <- p1 |>
  add_trace(
    x = c(0,U[1,1]),
    y = c(0,U[2,1]),
    z = c(0,U[3,1]),
    mode = "lines",
    line = list(width = 5, color = "blue"),
    type="scatter3d",
    name = "U1") |>
  add_trace(
    x = c(0,U[1,2]),
    y = c(0,U[2,2]),
    z = c(0,U[3,2]),
    mode = "lines",
    line = list(width = 5, color = "blue"),
    type="scatter3d",
    name = "U2")

p3
```


X3
X2
X1


- $\mathbf{U}^T\mathbf{Y}$ is the projection of $\mathbf{Y}$ in the space spanned by $\mathbf{U}$.
- Indeed $\mathbf{U}_1^T\mathbf{Y}$

```
p4 <- p3 |>
  add_trace(
    x = c(0,Y[1]),
    y = c(0,Y[2]),
    z = c(0,Y[3]),
    mode = "lines",
    line = list(width = 5, color = "red"),
    type="scatter3d",
    name = "Y") |>
  add_trace(
    x = c(0,U[1,1]*(U[,1]%*%Y)),
    y = c(0,U[2,1]*(U[,1]%*%Y)),
    z = c(0,U[3,1]*(U[,1]%*%Y)),
    mode = "lines",
    line = list(width = 5, color = "red",dash="dash"),
    type="scatter3d",
    name="Y -> U1") |> add_trace(
    x = c(Y[1],U[1,1]*(U[,1]%*%Y)),
    y = c(Y[2],U[2,1]*(U[,1]%*%Y)),
    z = c(Y[3],U[3,1]*(U[,1]%*%Y)),
    mode = "lines",
    line = list(width = 5, color = "red", dash="dash"),
    type="scatter3d",
    name="Y -> U1")
p4
```


X3
X2
X1


- and $\mathbf{U}_2^T\mathbf{Y}$

```
p5 <- p4 |>
  add_trace(
    x = c(0,U[1,2]*(U[,2]%*%Y)),
    y = c(0,U[2,2]*(U[,2]%*%Y)),
    z = c(0,U[3,2]*(U[,2]%*%Y)),
    mode = "lines",
    line = list(width = 5, color = "red",dash="dash"),
    type="scatter3d",
    name="Y -> U2") |> add_trace(
    x = c(Y[1],U[1,2]*(U[,2]%*%Y)),
    y = c(Y[2],U[2,2]*(U[,2]%*%Y)),
    z = c(Y[3],U[3,2]*(U[,2]%*%Y)),
    mode = "lines",
    line = list(width = 5, color = "red", dash="dash"),
    type="scatter3d",
    name="Y -> U2")
p5
```


X3
X2
X1

- $\hat{\mathbf{Y}}$ is the resulting vector that lies in the plane spanned by $\mathbf{U}_1$ and $\mathbf{U}_2$ and thus also in the column space of $\mathbf{X}$.

```
p6 <- p5 |>
  add_trace(
    x = c(0,Yhat[1]),
    y = c(0,Yhat[2]),
    z = c(0,Yhat[3]),
    mode = "lines",
    line = list(width = 5, color = "orange"),
    type="scatter3d",
    name = "Yhat") |>
  add_trace(
    x = c(Y[1],Yhat[1]),
    y = c(Y[2],Yhat[2]),
    z = c(Y[3],Yhat[3]),
    mode = "lines",
    line = list(width = 5, color = "maroon2"),
    type="scatter3d",
    name = "e") |>
  add_trace(
    x = c(U[1,1]*(U[,1]%*%Y),Yhat[1]),
    y = c(U[2,1]*(U[,1]%*%Y),Yhat[2]),
    z = c(U[3,1]*(U[,1]%*%Y),Yhat[3]),
    mode = "lines",
    line = list(width = 5, color = "orange", dash="dash"),
    type="scatter3d",
    name = "Y -> U")  |>
  add_trace(
    x = c(U[1,2]*(U[,2]%*%Y),Yhat[1]),
    y = c(U[2,2]*(U[,2]%*%Y),Yhat[2]),
    z = c(U[3,2]*(U[,2]%*%Y),Yhat[3]),
    mode = "lines",
    line = list(width = 5, color = "orange", dash="dash"),
    type="scatter3d",
    name = "Y -> U")
p6
```


X3
X2
X1

### Error

Note, that it is also clear from the equation in the derivation of the least squares solution that the residual is orthogonal on the column space:

$$-2\mathbf{X}^T(\mathbf{Y} - \mathbf{X}\beta) = 0$$

---

## Variance Estimator?

$$\begin{aligned}
\hat{\Sigma} &= \text{var}\left[(\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{Y}\right] \\
&= (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\text{var}\left[\mathbf{Y}\right]\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1} \\
&= (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T(\mathbf{I}\sigma^2)\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1} \\
&= (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{I} \quad \mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\sigma^2 \\
&= (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{X}(\mathbf{X}^T\mathbf{X})^{-1}\sigma^2 \\
&= (\mathbf{X}^T\mathbf{X})^{-1}\sigma^2
\end{aligned}$$

---

## Contrasts

Hypotheses often involve linear combinations of the model parameters!

e.g.

- $H_0 : \log_2 FC_{g3n1-g1n1} = \beta_{g3} + \hat{\beta}_{g3n1} = 0 \rightarrow$ "grade3+grade3:node1 = 0"
- Let

$$\beta = \begin{bmatrix} \beta_0 \\ \beta_{g3} \\ \beta_{n1} \\ \beta_{g3:n1} \end{bmatrix}$$

- we can write that contrast using a contrast matrix:

$$\mathbf{L} = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix} \rightarrow \mathbf{L}^T\beta$$

- Then the variance becomes:

$$\text{var}_{\mathbf{L}^T\hat{\beta}} = \mathbf{L}^T\Sigma_{\hat{\beta}}\mathbf{L}$$

---

## Homework: Adopt the gene analysis on log scale in matrix form!

1. Study the solution of the exercise to understand the analysis in R
2. Calculate

- model parameters and contrasts of interest
- standard errors, standard errors on contrasts
- t-test statistics on the model parameters and contrasts of interest

3. Compare your results with the output of the lm(.) function

---

### Inspiration

Tip: details on the implementation can be found in the book of Faraway (chapter 2). https://people.bath.ac.uk/jjf23/book/

- Design matrix

```
X <- model.matrix(~grade*node,data=gene)
```

- Transpose of a matrix: use function t(.)

```
t(X)
```

```
             1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24
(Intercept)  1 1 1 1 1 1 1 1 1  1  1  1  1  1  1  1  1  1  1  1  1  1  1  1
grade3       1 1 0 0 0 1 1 0 1  1  0  1  0  0  0  1  0  1  0  1  0  1  1  0
node1        1 1 1 1 0 0 1 1 0  0  0  0  0  0  0  1  1  0  1  1  1  0  1  0
grade3:node1 1 1 0 0 0 0 1 0 0  0  0  0  0  0  0  1  0  0  0  1  0  0  1  0
attr(,"assign")
[1] 0 1 2 3
attr(,"contrasts")
attr(,"contrasts")$grade
[1] "contr.treatment"

attr(,"contrasts")$node
[1] "contr.treatment"
```

- Matrix product %*% operator

```
t(X)%*%X
```

```
             (Intercept) grade3 node1 grade3:node1
(Intercept)           24     12    12            6
grade3                12     12     6            6
node1                 12      6    12            6
grade3:node1           6      6     6            6
```

- Degrees of freedom of a model?

$$df = n - p$$

```
summary(lm1)
```

```
Call:
lm(formula = y ~ x, data = data)

Residuals:
      1       2       3
-0.1667  0.3333 -0.1667

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept)   0.6667     0.6236   1.069    0.479
```

```
x             0.5000     0.2887   1.732    0.333

Residual standard error: 0.4082 on 1 degrees of freedom
Multiple R-squared:   0.75, Adjusted R-squared:    0.5
F-statistic:     3 on 1 and 1 DF,  p-value: 0.3333
```

```
dfRes <- (nrow(X)-ncol(X))
dfRes
```

```
[1] 20
```

- Variance estimator: MSE

$$\hat{\sigma}^2 = \frac{\sum_{i=1}^{n} \epsilon_i^2}{n-p}$$

- Invert matrix: use function solve(.)
- Diagonal elements of a matrix: use function diag(.)

```
t(X)%*%X
```

```
             (Intercept) grade3 node1 grade3:node1
(Intercept)           24     12    12            6
grade3                12     12     6            6
node1                 12      6    12            6
grade3:node1           6      6     6            6
```

```
diag(t(X)%*%X)
```

```
 (Intercept)       grade3        node1 grade3:node1
          24           12           12            6
```