

Sequencing: Bulk RNA-seq homework

Add your names here

8/6/2021

Contents

| | | |
|----------|--|----------|
| 1 | Default edgeR analysis | 1 |
| 2 | Impact of blocking | 1 |
| 3 | Analyze dataset using full-quantile normalization | 1 |
| 3.1 | Implement and apply full-quantile normalization | 1 |
| 3.2 | Visualize effect of FQ normalization | 2 |
| 3.3 | edgeR analysis using full-quantile normalized data | 2 |
| 3.4 | Compare DE genes at 5% FDR | 2 |

1 Default edgeR analysis

Analyze the data using `edgeR`, by using the code in the RNA-seq analysis intro lecture. **In all analyses, focus on the contrast comparing DPN treatment to control at 48h.**

2 Impact of blocking

Assess the difference in number of DE genes when not blocking on patient, i.e., removing the `patient` effect of the model. Compare the p-value distributions between these two models (i.e., with and without blocking).

3 Analyze dataset using full-quantile normalization

3.1 Implement and apply full-quantile normalization

```
### implement FQ normalization
FQnorm <- function(counts){
  ...
}

### normalize the data using FQ
```

3.2 Visualize effect of FQ normalization

Visualize the distributions of `log1p`-transformed counts (use the `density` function) to compare sample-specific count distributions before and after FQ normalization. What's the impact of FQ normalization on the differences in distribution between samples?

3.3 edgeR analysis using full-quantile normalized data

Don't forget to remove the `calcNormFactors` step in the `edgeR` analysis as data have already been normalized when using FQ-normalized counts as input!

```
### use FQ-normalized data as input to the edgeR analysis
```

3.4 Compare DE genes at 5% FDR

```
### Get list of DE genes using TMM and FQ normalization
```

```
### Compare list using, e.g., a Venn diagram
```