






MODEL WITH BLOCK EFFECTS




Outcome:  $L_i = \text{pulled left}$

$$L_i \sim \textit{Bernoulli}(p_i)$$

$$\textit{logit}(p_i) = \alpha_{actor[i]} + \gamma_{block[i]} + \beta_{treatment[i]}$$







Accounts systematic differences across blocks.

Next step is to add priors that introduce dependencies across coefficients.

# MODEL WITH BLOCK EFFECTS

Outcome:  $L_i$  = pulled left

$$L_i \sim \text{Bernoulli}(p_i)$$

$$\text{logit}(p_i) = \alpha_{\text{actor}[i]} + \gamma_{\text{block}[i]} + \beta_{\text{treatment}[i]}$$

Accounts systematic differences across blocks.

Next step is to add priors that introduce dependencies across coefficients.

# ADDING REGULARIZING PRIORS FOR BLOCK

Outcome:  $L_i$  = pulled left

$$L_i \sim \text{Bernoulli}(p_i)$$

$$\text{logit}(p_i) = \alpha_{\text{actor}[i]} + \gamma_{\text{block}[i]} + \beta_{\text{treatment}[i]}$$

$$\gamma_j \sim \text{Normal}(0, \sigma_\gamma), \text{ for } j = 1..6$$