


RANDOM SLOPES MODEL



$$y_i \sim \textit{Normal}(\mu_i, \sigma)$$

$$\mu_i = \alpha_0 + \alpha_{block[i]} + (\beta_0 + \beta_{block[i]})x_i$$

$$\alpha_k \sim \textit{Normal}(0, \sigma_\alpha), \text{ for } k \text{ in } \{1, \cdots, N_{blocks}\}$$

$$\beta_k \sim \textit{Normal}(0, \sigma_\beta), \text{ for } k \text{ in } \{1, \cdots, N_{blocks}\}$$


$$\alpha_0, \beta_0 \sim \textit{Normal}(0, 1)$$

$$\beta \sim \textit{Normal}(0, 0.3)$$

$$\sigma, \sigma_{block} \sim \textit{Exponential}(1)$$



Allows the slope
associated with the
x predictor to vary
across blocks, but
maintains a
dependency across
blocks



RANDOM SLOPES MODEL

$$y_i \sim \text{Normal}(\mu_i, \sigma)$$

$$\mu_i = \alpha_0 + \alpha_{\text{block}[i]} + (\beta_0 + \beta_{\text{block}[i]})x_i$$

$$\alpha_k \sim \text{Normal}(0, \sigma_\alpha), \text{ for } k \text{ in } \{1, \dots, N_{\text{blocks}}\}$$

$$\beta_k \sim \text{Normal}(0, \sigma_\beta), \text{ for } k \text{ in } \{1, \dots, N_{\text{blocks}}\}$$

$$\alpha_0, \beta_0 \sim \text{Normal}(0, 1)$$

$$\beta \sim \text{Normal}(0, 0.3)$$

$$\sigma, \sigma_{\text{block}} \sim \text{Exponential}(1)$$

Allows the slope associated with the x predictor to vary across blocks, but maintains a **dependency** across blocks

MORE COMPLEX STRUCTURING OF THE DEPENDENCY ACROSS COEFFICIENTS


Phylogenetic models

Evolutionary relatedness



Animal models

Genetic relatedness



Spacial auto-correlation

Spacial proximity

