

ADDING REGULARIZING PRIORS FOR BLOCK



Outcome: L_i = pulled left

$$L_i \sim \textit{Bernoulli}(p_i)$$

$$\textit{logit}(p_i) = \alpha_{actor[i]} + \gamma_{block[i]} + \beta_{treatment[i]}$$

$$\gamma_j \sim \textit{Normal}(0, \sigma_\gamma), \text{ for } j = 1..6$$





This estimates how similar the γ_j coefficients are to each other and changes the estimates accordingly

ADDING REGULARIZING PRIORS FOR BLOCK

Outcome: L_i = pulled left

$$L_i \sim \text{Bernoulli}(p_i)$$

$$\text{logit}(p_i) = \alpha_{\text{actor}[i]} + \gamma_{\text{block}[i]} + \beta_{\text{treatment}[i]}$$

$$\gamma_j \sim \text{Normal}(0, \sigma_\gamma), \text{ for } j = 1..6$$



This estimates how similar the γ_j coefficients are to each other and changes the estimates accordingly

REGULARIZING PRIOR FOR THE ACTOR COEFFICIENT

Outcome: L_i = pulled left

$$L_i \sim \text{Bernoulli}(p_i)$$

$$\text{logit}(p_i) = \alpha_{\text{actor}[i]} + \gamma_{\text{block}[i]} + \beta_{\text{treatment}[i]}$$

$$\gamma_j \sim \text{Normal}(0, \sigma_\gamma), \text{ for } j = 1..6$$

$$\alpha_j \sim \text{Normal}(\alpha_0, \sigma_\alpha), \text{ for } j = 1..7$$