

ch1_phase1

October 1, 2020

```
[1]: from sympy import Matrix, Symbol, derive_by_array, Lambda, □
      ↪ symbols, Derivative, diff
      from sympy.abc import x, y, i, j, a, b
```

Defining variable-element matrices $X \in \mathbb{R}^{n \times m}$ and $W \in \mathbb{R}^{m \times p}$:

```
[2]: def var(letter: str, i: int, j: int) -> Symbol:
      letter_ij = Symbol('{}_{-}{}'.format(letter, i+1, j+1), □
      ↪ is_commutative=True)
      return letter_ij

n,m,p = 3,3,2

X = Matrix(n, m, lambda i,j : var('x', i, j)); X
```

```
[2]: 
$$\begin{bmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ x_{31} & x_{32} & x_{33} \end{bmatrix}$$

```

```
[3]: W = Matrix(m, p, lambda i,j : var('w', i, j)); W
```

```
[3]: 
$$\begin{bmatrix} w_{11} & w_{12} \\ w_{21} & w_{22} \\ w_{31} & w_{32} \end{bmatrix}$$

```

Defining $N = \nu(X, W) = X \times W$

- $\nu: \mathbb{R}^{(n \times m) \times (m \times p)} \rightarrow \mathbb{R}^{n \times p}$
- $N \in \mathbb{R}^{n \times p}$

```
[4]: v = Lambda((a,b), a*b); v
```

[4]: $((a, b) \mapsto ab)$

[5]: $N = v(X, W); N$

[5]:
$$\begin{bmatrix} w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13} & w_{12}x_{11} + w_{22}x_{12} + w_{32}x_{13} \\ w_{11}x_{21} + w_{21}x_{22} + w_{31}x_{23} & w_{12}x_{21} + w_{22}x_{22} + w_{32}x_{23} \\ w_{11}x_{31} + w_{21}x_{32} + w_{31}x_{33} & w_{12}x_{31} + w_{22}x_{32} + w_{32}x_{33} \end{bmatrix}$$

Defining $S = \sigma_{\text{apply}}(N) = \sigma_{\text{apply}}(v(X, W)) = \sigma_{\text{apply}}(X \times W) = \{\sigma(XW_{ij})\}$.

Assume that $\sigma_{\text{apply}} : \mathbb{R}^{n \times p} \rightarrow \mathbb{R}^{n \times p}$ while $\sigma : \mathbb{R} \rightarrow \mathbb{R}$, so the function σ_{apply} takes in a matrix and returns a matrix while the simple σ acts on the individual elements $N_{ij} = XW_{ij}$ in the matrix argument N of σ_{apply} .

- $\sigma : \mathbb{R} \rightarrow \mathbb{R}$
- $\sigma_{\text{apply}} : \mathbb{R}^{n \times p} \rightarrow \mathbb{R}^{n \times p}$
- $S \in \mathbb{R}^{n \times p}$

```
[6]: from sympy import Function

# Nvec = Symbol('N', commutative=False)

sigma = Function('sigma')
sigma(N[0,0])
```

[6]: $\sigma(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13})$

```
[7]: # way 1 of declaring S
S = N.applyfunc(sigma); S
#type(S)
#Matrix(3, 2, lambda i, j: sigma(N[i,j]))
```

[7]:
$$\begin{bmatrix} \sigma(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13}) & \sigma(w_{12}x_{11} + w_{22}x_{12} + w_{32}x_{13}) \\ \sigma(w_{11}x_{21} + w_{21}x_{22} + w_{31}x_{23}) & \sigma(w_{12}x_{21} + w_{22}x_{22} + w_{32}x_{23}) \\ \sigma(w_{11}x_{31} + w_{21}x_{32} + w_{31}x_{33}) & \sigma(w_{12}x_{31} + w_{22}x_{32} + w_{32}x_{33}) \end{bmatrix}$$

```
[8]: # way 2 of declaring S (better way)
sigmaApply = lambda matrix: matrix.applyfunc(sigma)

sigmaApply(N)
```

[8]:

$$\begin{bmatrix} \sigma(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13}) & \sigma(w_{12}x_{11} + w_{22}x_{12} + w_{32}x_{13}) \\ \sigma(w_{11}x_{21} + w_{21}x_{22} + w_{31}x_{23}) & \sigma(w_{12}x_{21} + w_{22}x_{22} + w_{32}x_{23}) \\ \sigma(w_{11}x_{31} + w_{21}x_{32} + w_{31}x_{33}) & \sigma(w_{12}x_{31} + w_{22}x_{32} + w_{32}x_{33}) \end{bmatrix}$$

[9]: `sigmaApply(X**2)` # can apply this function to any matrix
↪ argument.

[9]:
$$\begin{bmatrix} \sigma(x_{11}^2 + x_{12}x_{21} + x_{13}x_{31}) & \sigma(x_{11}x_{12} + x_{12}x_{22} + x_{13}x_{32}) & \sigma(x_{11}x_{13} + x_{12}x_{23} + x_{13}x_{33}) \\ \sigma(x_{11}x_{21} + x_{21}x_{22} + x_{23}x_{31}) & \sigma(x_{12}x_{21} + x_{22}^2 + x_{23}x_{32}) & \sigma(x_{13}x_{21} + x_{22}x_{23} + x_{23}x_{33}) \\ \sigma(x_{11}x_{31} + x_{21}x_{32} + x_{31}x_{33}) & \sigma(x_{12}x_{31} + x_{22}x_{32} + x_{32}x_{33}) & \sigma(x_{13}x_{31} + x_{23}x_{32} + x_{33}^2) \end{bmatrix}$$

[10]: `S = sigmaApply(v(X,W))` # composing
`S`

[10]:
$$\begin{bmatrix} \sigma(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13}) & \sigma(w_{12}x_{11} + w_{22}x_{12} + w_{32}x_{13}) \\ \sigma(w_{11}x_{21} + w_{21}x_{22} + w_{31}x_{23}) & \sigma(w_{12}x_{21} + w_{22}x_{22} + w_{32}x_{23}) \\ \sigma(w_{11}x_{31} + w_{21}x_{32} + w_{31}x_{33}) & \sigma(w_{12}x_{31} + w_{22}x_{32} + w_{32}x_{33}) \end{bmatrix}$$

Defining $L = \Lambda(S) = \Lambda(\sigma_{\text{apply}}(\nu(X, W))) = \Lambda(\{\sigma(XW_{ij})\})$. In general, let the function be defined as:

$$\begin{aligned} L &= \Lambda \begin{pmatrix} \sigma(XW_{11}) & \sigma(XW_{12}) & \dots & \sigma(XW_{1p}) \\ \sigma(XW_{21}) & \sigma(XW_{22}) & \dots & \sigma(XW_{2p}) \\ \vdots & \vdots & & \vdots \\ \sigma(XW_{n1}) & \sigma(XW_{n2}) & \dots & \sigma(XW_{np}) \end{pmatrix} \\ &= \sum_{i=1}^p \sum_{j=1}^n \sigma(XW_{ij}) \\ &= \sigma(XW_{11}) + \sigma(XW_{12}) + \dots + \sigma(XW_{np}) \end{aligned}$$

NOTE HERE: * $\Lambda: \mathbb{R}^{n \times p} \rightarrow \mathbb{R}$ * $L \in \mathbb{R}$

[11]: `lambdaF = lambda matrix : sum(matrix)`
`lambdaF(S)`

[11]:
$$\begin{aligned} &\sigma(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13}) && + && \sigma(w_{11}x_{21} + w_{21}x_{22} + w_{31}x_{23}) && + \\ &\sigma(w_{11}x_{31} + w_{21}x_{32} + w_{31}x_{33}) && + && \sigma(w_{12}x_{11} + w_{22}x_{12} + w_{32}x_{13}) && + \\ &\sigma(w_{12}x_{21} + w_{22}x_{22} + w_{32}x_{23}) + \sigma(w_{12}x_{31} + w_{22}x_{32} + w_{32}x_{33}) \end{aligned}$$

```
[12]: L = lambdaF(sigmaApply(v(X, W)))
      L
      #L = lambda mat1, mat2: lambdaF(sigmaApply(v(mat1, mat2)))
      #L(X, W)
```

```
[12]: 
$$\begin{aligned} &\sigma(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13}) + \sigma(w_{11}x_{21} + w_{21}x_{22} + w_{31}x_{23}) + \\ &\sigma(w_{11}x_{31} + w_{21}x_{32} + w_{31}x_{33}) + \sigma(w_{12}x_{11} + w_{22}x_{12} + w_{32}x_{13}) + \\ &\sigma(w_{12}x_{21} + w_{22}x_{22} + w_{32}x_{23}) + \sigma(w_{12}x_{31} + w_{22}x_{32} + w_{32}x_{33}) \end{aligned}$$

```

```
[13]: #derive_by_array(L, X)
```

```
[14]: derive_by_array(L, S)
```

```
[14]: 
$$\begin{bmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{bmatrix}$$

```

```
[15]: from sympy import sympify, lambdify
      n = lambdify((X[0,0],X[0,1],X[0,2],W[0,0],W[1,0],W[2,0]), □
      ↪ N[0,0])
      n(1,2,3,4,3,2)
```

```
[15]: 16
```

```
[16]: f = Function('f') #(sympify(N[0,0]))
      f(N[0,0])
```

```
[16]:  $f(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13})$ 
```

```
[17]: f(N[0,0]).diff(X[0,0])
```

```
[17]: 
$$w_{11} \frac{d}{d\xi_1} f(\xi_1) \Big|_{\xi_1 = w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13}}$$

```

```
[18]: n = v(X,W); n
      n11 = Function('{}'.format(n[0,0]))
      n11
```

```
[18]: w_11*x_11 + w_21*x_12 + w_31*x_13
```

```
[19]: s_ij = Function('s_ij')
      sig = Function('sig')(x)
```

```
[20]: # KEY: got not expecting UndefinedFunction error again here too
      #S_ij = Matrix(3, 2, lambda i,j: Function('s_{}'.format(i+1,j+1))(Function('{}'.format(N[i,j]))))
```

```
[21]: #S_ij[0,0](sympify(N[0,0])).diff(sympify(N[0,0]))
      F = 3*x*y

      xy = Symbol('{}'.format(F))
      xy.subs({x:3})
      sympify(xy).subs({x:3})
```

```
[21]: 3 * x * y
```

Sympy Example of trying to differentiate with respect to an expression not just a variable.

```
[22]: from sympy.abc import t

      F = Function('F')
      f = Function('f')
      U = f(t)
      V = U.diff(t)

      direct = F(t, U, V).diff(U); direct
```

```
[22]: 
$$\frac{\partial}{\partial \xi_2} F\left(t, \xi_2, \frac{d}{dt} f(t)\right) \Big|_{\xi_2=f(t)}$$

```

```
[23]: F(t,U,V)
```

```
[23]: 
$$F\left(t, f(t), \frac{d}{dt} f(t)\right)$$

```

```
[24]: F(t,U,V).subs(U,x)
```

```
[24]: 
$$F\left(t, x, \frac{d}{dt} x\right)$$

```

```
[25]: F(t,U,V).subs(U,x).diff(x)
```

```
[25]: 
$$\frac{\partial}{\partial \xi_2} F\left(t, \xi_2, \frac{d}{dt} x\right) \Big|_{\xi_2=x}$$

```

```
[26]: F(t,U,V).subs(U,x).diff(x).subs(x, U)
```

```
[26]: 
$$\frac{\partial}{\partial \xi_2} F\left(t, \xi_2, \frac{d}{dt} f(t)\right) \Big|_{\xi_2=f(t)}$$

```

```
[27]: indirect = F(t,U,V).subs(U, x).diff(x).subs(x,U); indirect
```

```
[27]: 
$$\frac{\partial}{\partial \xi_2} F\left(t, \xi_2, \frac{d}{dt} f(t)\right) \Big|_{\xi_2=f(t)}$$

```

```
[28]: F = Lambda((x,y), 3*x* y)
      F(1,2)
```

```
[28]: 6
```

```
[29]: U = x*y
      G = 3*x*y
      xy
```

```
[29]: 3 * x * y
```

```
[30]: F.diff(xy)
```

```
[30]: 0
```

```
[31]: # derive_by_array(S, N) # ERROR
```

```
[32]: s11 = S[0,0]
      s11
```

```
[32]: 
$$\sigma(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13})$$

```

```
[33]: #s11.diff(n11)
```

```
[34]: derive_by_array(L, S)
```

```
[34]:
```

$$\begin{bmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{bmatrix}$$

```
[35]: x, y, r, t = symbols('x y r t') # r (radius), t (angle theta)
      f, g, h = symbols('f g h', cls=Function)
      h = g(f(x))
      Derivative(h, f(x)).doit()
```

```
[35]: 
$$\frac{d}{df(x)} g(f(x))$$

```

```
[36]: h.args[0]
      h.diff(h.args[0])
```

```
[36]: 
$$\frac{d}{df(x)} g(f(x))$$

```

```
[37]: S = sigmaApply(v(X,W)); S
```

```
[37]: 
$$\begin{bmatrix} \sigma(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13}) & \sigma(w_{12}x_{11} + w_{22}x_{12} + w_{32}x_{13}) \\ \sigma(w_{11}x_{21} + w_{21}x_{22} + w_{31}x_{23}) & \sigma(w_{12}x_{21} + w_{22}x_{22} + w_{32}x_{23}) \\ \sigma(w_{11}x_{31} + w_{21}x_{32} + w_{31}x_{33}) & \sigma(w_{12}x_{31} + w_{22}x_{32} + w_{32}x_{33}) \end{bmatrix}$$

```

```
[38]: from sympy.abc import n

      n11 = (X*W)[0,0]
      m = lambda mat1, mat2: sympify(Symbol('{}'.format((mat1 *
      ↪mat2)[0,0] )))
      s = sigma(m(X,W)); s
```

```
[38]: 
$$\sigma(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13})$$

```

```
[39]: s.subs({W[0,0]: 14}) # doesn't work to substitute into an
      ↪undefined function
```

```
[39]: 
$$\sigma(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13})$$

```

```
[40]: Derivative(s, m(X,W)).doit()
```

```
[40]: 
$$\frac{d}{dw_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13}} \sigma(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13})$$

```

```
[41]: #s11 = Function('s_{11}')(n11); s11
      #sigma(n11).diff(n11)

      #s11.diff(n11)
      sigma(n11)
```

```
[41]:  $\sigma(w_{11}x_{11} + w_{21}x_{12} + w_{31}x_{13})$ 
```

```
[42]: # ERROR HERE TOO
      type(sigma(n11).args[0])
```

```
[42]: sympy.core.add.Add
```

```
[43]: type(n11)
```

```
[43]: sympy.core.add.Add
```

```
[44]: #sigma(n11).diff(sigma(n11).args[0]) ## ERROR
```

```
[45]:
```

```
[45]: b = Symbol('{}'.format(n11))
      ns_11 = Function(b, real=True)
      ns_11

      # ERROR cannot diff wi.r. to undefinedfunction
      # sigma(n11).diff(ns_11)

      #
      #sigma(b).diff(b).subs({b:1})
```

```
[45]: w_11*x_11 + w_21*x_12 + w_31*x_13
```

```
[46]: f, g = symbols('f g', cls=Function)
      xy = Symbol('x*y'); xy
      #sympify(xy).subs({x:2, y:4})
      f(g(x,y)).diff(xy)
```


[46]: 0

```
[47]: # TODO SEEM to have got the expression but it is not working
      ↪since can't substitute anything .... ???
      f(xy).diff(xy).subs({x:2})
```

[47]: $\frac{d}{dx * y} f(x * y)$

```
[48]: Function("x*y")(x,y)
      xyf = lambdify([x,y],xy)
      xyf(3,4)
      f(g(xy)).diff(xy)
      #
```

[48]: $\frac{d}{dg(x * y)} f(g(x * y)) \frac{d}{dx * y} g(x * y)$

```
[49]: xyd = Derivative(x*y, x*y,0).doit();xyd

      #Derivative(3*xyd, xyd, 1).doit() ### ERROR can't calc deriv w.
      ↪r.t to x*y
```

[49]: xy

```
[50]: #derive_by_array(S, N)
```

```
[51]:
```

```
[51]:
```