

Worksheet 10 (Solutions)

1. **(Ratio Test)** Let $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Poisson}(\lambda)$. What is the test statistic Λ for the corresponding likelihood ratio test for the null hypothesis $H_0 : \lambda = 1$.

Solution: TODO

2. **(Ratio Test)** Let $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Exp}(\lambda)$. What is the test statistic Λ for the corresponding likelihood ratio test for the null hypothesis $H_0 : \lambda = 1$.

Solution: TODO

3. **(Ratio Test)** Let $X \sim \text{Bin}(n, p_1)$ and $Y \sim \text{Bin}(n, p_2)$ be independent random variables. We want to test the hypothesis that $H_0 : p_1 = p_2$. What are the corresponding Θ and Θ_0 in our updated formulation of hypothesis testing?¹ If we use a Likelihood Ratio Test for this hypothesis, how many degrees of freedom should Λ have?

¹ We will derive the actual test itself in a more general form next class.

Solution: TODO

4. **(Ratio Test)** Recall that we used the one-sample ANOVA test with the null-hypothesis that the means of K samples are all the same. Write down and describe the values of Θ and Θ_0 that correspond to this test. If we use a Likelihood Ratio Test for this hypothesis, how many degrees of freedom should Λ have?

Solution: TODO

5. **(MLE Practice)** Let $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Uniform}(0, a)$. Find the MLE estimator for a . Note: You cannot do this using the derivative. Just think about it!

Solution: TODO