Project 1

STAT 801A, Fall 2026

Due on September 18, 2026 at 5:00 pm

Overview

The American Community Survey is a large national survey that collects information on many aspects of life in the United States. You are tasked with analyzing a subset of the 2022 survey for people who live and work in Lancaster County, Nebraska. In this particular subset, respondents were asked how long it takes for them to commute to work and only responses for people who drive alone are included. The data is available in the file `travel-time-31109.csv`.

Variable Descriptions:

- `RESPONDENT`: Unique identifier for each respondent
- `TRANTIME`: Travel time to work in minutes
- `DEPARTS`: Time of day the respondent leaves for work (e.g., 7:30 AM = 730)
- `ARRIVES`: Time of day the respondent arrives at work (e.g., 7:30 AM = 730)

Problems

Complete each of the following problems using the provided Quarto template. Your code must be reproducible, but you are free to use any R packages that you would like (i.e., I can get the same output file by rendering your Quarto file). For each problem, include your code, output, and a brief written interpretation of your results. Submit your completed Quarto file and the rendered PDF file to Canvas.

1. Read in the data into an object called `ne_work_commute` and use the `head()` function and its correct arguments to print out the first 10 observations.
2. Use the `dim()` function to get the dimensions of the dataset. In writing (1 sentence), state the number of rows and columns of `ne_work_commute`.
3. Compute the mean of `TRANTIME`.
4. Compute the median of `TRANTIME`.

5. Compute the variance of `TRANTIME`.
6. Compute the standard deviation of `TRANTIME`.
7. Use the `summary()` function on `TRANTIME` and interpret what each value represents.
8. Create a histogram of `TRANTIME` using interpretable text for the title and axis text (Note: You may use either base R or `ggplot2`.)
9. Would the mean or median be a better description of travel time? Briefly explain your decision.
10. Your supervisor wants to remove outliers when reporting the summary statistics using the interquartile range (IQR) method. Create a vector called `filtered_travel_time` that contains values that are not outliers. A value is considered an outlier if it is less than $Q_1 - 1.5 \times IQR$ or greater than $Q_3 + 1.5 \times IQR$. (Hint: the `IQR()` function would be helpful here.)
11. Create a histogram of `filtered_travel_time`.
12. Repeat Problems 3–6 using `filtered_travel_time` and arrange your results into a nicely formatted table. Round all values for this table to two decimal places. (Hint: create a data frame with the summary statistics and use the `knitr::kable()` function.)
13. Briefly describe what changed with your summary statistics after removing the outliers.
14. Let X be normally distributed with the mean and variance you computed for `filtered_travel_time` (do not round these values). What is the probability that a person who lives and works in Lancaster County has a travel time of:

    a. less than 10 minutes
    b. between 10 and 20 minutes
    c. greater than 20 minutes