

Thermal optima and heat stress reveal vulnerabilities in a common butterfly

Grace Jeschke, Leslie Ries, Naresh Neupane, and Elise A Larsen  Ries Lab, Biology Department, Georgetown University

GJ: PhD student; grj14@georgetown.edu ORCID: 0000-0001-6271-7750

Background

Silver-spotted skipper (*Epargyreus clarus*)¹

- Larvae eat many legume spp. (Fabaceae)
- Winters as pupa
- Occurs across North America (Fig. 2)
- Number of generations per year (voltinism) varies
 - more generations at lower latitudes
 - May interact with climate change¹⁰



Fig. 1. Clockwise from top left: new silver-spotted skipper egg; egg later in development; hatchlings; adult butterflies

Climate change & generations

- ↑ Higher accumulation of growing degree days (GDD)
- ↓ Extreme weather and phenological mismatch^{11,12}




Fig. 2 Range map of silver-spotted skipper from iNaturalist² observations with MA and NC noted. Numbers on the right indicate number of generations, with 1.5 indicating a partial 2nd generation. Sources^{1, 3, 4, 5, 6, 7, 8, 9}

Adaptation Potential

- Thermal limits¹⁰
- Preserved obligate species interactions
 - e.g., larval host-plants, symbionts¹²
- Ability to add partial generations (bet-hedging)
- Longer growing season may increase voltinism

Methods

Silver-spotted skippers have been extensively studied in the DC region. To expand existing data⁷, we collected eggs from populations (circled in map):

- south (NC)
- north (MA)



- We measured
- growth and development
 - the effect of diet on immune function (data not shown)
 - survival and fitness,
 - thermal performance.

To evaluate thermal performance, we reared individuals at five oscillating temperatures: 16C, 21C, 26C, 30C, 34C, all +/- 5C.




Fig. 3 Above: sample record of one individual's rearing temperature, source and development; left: final-instar silver-spotted skipper larva.

Results

Survival and Development

- At 16 +/- 5C, none survived to adulthood. Most died prior to the 2nd of five instars.
- Growth rate increased with temperature until 34 +/- 5C, at which, growth rate and survival fell.
- Growth at 30C was slightly higher vs 26 in MA cohort, and no different in NC.
- MA larvae grew more slowly and had higher mortality.

Diapause

- Most (90%) of MA pupae entered diapause. Those with direct development (n = 7) had faster growth.

Estimated thermal performance curves (TPC) Fig. 4

- Solid lines indicate relative growth rate at given temp (x); dashed lines indicate relative survival (y)
- Optimal performance in the temperature change where both curves >= 50%¹³
- T_{opt} is the peak growth rate; T_{max} is the temp where growth rate decreases





Figure 5: Forecasting

Top: total Growing Degree Days (GDD) calculated from thermal performance results accumulated between May 1-Sep 30 calculated from past temperature (1990-2020) and predicted future models (2050-2070). MA has 1-1.5 generations according to the model, with caveats; these may stay the same or increase in future. 1996 (the dashed line) is the first year that the last skipper observation in MA was later than August.⁵ NC has had 2-3 generations, with 3-4 predicted in the future.

Bottom: total LDD (+1 for log scale) calculated as for GDD: LDD indicates heat stress accumulated above the GDD T_{max}. MA experienced almost no LDD between 1990-2020, and is predicted to remain low. In contrast, NC has had cyclical years of high LDD, and the most extreme climate model predicts an increase to >100 LDD.

Conclusions & Ongoing Work

- In the future, developmental rate & # of generations could increase, or be blunted by LDD
- More work is needed to understand the effect of temperatures above the linear range. We do not yet understand the extent to which a species can tolerate LDD at any given developmental stage, and/or to what extent the sequence and intensity of GDD/LDD matter.

NEXT:

- Validate the phenology and generations predicted by GDD models using citizen science data, including decades of structured citizen science survey data
- Expand model to other areas of range and validate
- With the same citizen science data, investigate the role of other factors threatening butterfly populations




Fig. 6: Example of intended output. Thermal tolerance of *Speyeria idalia*. Color ramp indicates seasonal LDD averaged over the years indicated, for north-central USA. LDD are predicted to increase, especially in mid-summer. Data visualization by Naresh Neupane.



MIDWEST
Climate Adaptation
Science Center

- SOURCES**
1. Scott, J. A. 1986. Butterflies of North America. Stanford University Press, Stanford, California.
 2. iNaturalist <https://www.inaturalist.org/taxa/81559-Epargyreus-clarus> retrieved 6 Aug 2024
 3. Ebner, J. A. 1970. The butterflies of Wisconsin. Board of Trustees, Milwaukee Public Museum, Milwaukee, Wisconsin.
 4. Shapiro, A. 2007. Field guide to butterflies of the San Francisco Bay and Sacramento Valley Regions. Berkeley: University of California Press.
 5. Stichter, S. 2014. The butterflies of Massachusetts: silver-spotted skipper. <<http://www.butterfliesofmassachusetts.net/silverspot-sk.htm>>. Downloaded on 2 March 2022.
 6. Schlicht, D.W., Downey, J.C. and Nekola, J.C. 2007. The butterflies of Iowa. University of Iowa Press, Iowa City.
 7. Abarca, M., E. A. Larsen, J. T. Lili, M. Weiss, E. Lind, and L. Ries. 2018. Inclusion of host quality data improves predictions of herbivore phenology. Entomologia Experimentalis et Applicata 166:648–660.
 8. Ittner, D.C., Shuey, J. A., and J. V. Calhoun. 1992. Butterflies and skippers of Ohio. Ohio Biological Survey. Bulletin. Series Vol. 9 No. 1.

9. Irwin, R. R. and J. C. Downey. 1973. Annotated checklist of the butterflies of Illinois. Illinois Natural History Survey, Urbana.
10. Abarca, M., and R. Spahn. 2021. Direct and indirect effects of altered temperature regimes and phenological mismatches on insect populations. Current Opinion in Insect Science 47:67-74.
11. Perkins-Kirkpatrick, S. E., and P. B. Gibson. 2018. Changes in regional heatwave characteristics as a function of increasing global temperature. Scientific Reports:1–12.
12. Schieuning, M., E. L. Neuschulz, J. Albrecht, I. M.A. Bender, D. E. Bowler, D. M. Dehling, S. A. Fritz, C. Hof, T. Mueller, L. Nowak, M. C. Sorensen, K. Böhning-Gaese, and W. D. Kissling. 2020. Trait-based assessments of climate-change impacts on interacting species. Trends in Ecology & Evolution, 35:319-28.
13. Abarca, M., Parker, A. L., Larsen, E. A., Umbanhowar, J., Earl, C., Guralnick R, et al. (2024) How development and survival combine to determine the thermal sensitivity of insects. PLoS ONE 19(1): e0291393. <https://doi.org/10.1371/journal.pone.0291393>