

Intro to AI and ML

Matrix Project

Christian Stavan Nilesch,
Uttam Choudhary

February 14, 2019

Geometry Question-

A circle passes through the points $\begin{pmatrix} 2 \\ 3 \end{pmatrix}$ and $\begin{pmatrix} 4 \\ 5 \end{pmatrix}$. If its centre lies on the line

$$(-1 \ 4) \mathbf{x} + 3 = 0 \quad (1)$$

find its radius.

Solution-

Assume centre of required circle is given by \mathbf{c} and radius r . Then equation of circle will be:

$$\mathbf{x}^T \mathbf{x} - 2\mathbf{c}^T \mathbf{x} = r^2 - \mathbf{c}^T \mathbf{c} \quad (2)$$

Let $\mathbf{A} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$ and $\mathbf{B} = \begin{pmatrix} 4 \\ 5 \end{pmatrix}$ be the given points. Substituting the two points \mathbf{A} and \mathbf{B} in the equation of circle:

$$\begin{pmatrix} 2 & 3 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \end{pmatrix} - 2\mathbf{c}^T \begin{pmatrix} 2 \\ 3 \end{pmatrix} = r^2 - \mathbf{c}^T \mathbf{c} \quad (3)$$

and

$$\begin{pmatrix} 4 & 5 \end{pmatrix} \begin{pmatrix} 4 \\ 5 \end{pmatrix} - 2\mathbf{c}^T \begin{pmatrix} 4 \\ 5 \end{pmatrix} = r^2 - \mathbf{c}^T \mathbf{c} \quad (4)$$

Solution(contd)-

On taking difference of above two equations ((4)-(3)) we get:

$$41 - 13 - 2\mathbf{c}^T \begin{pmatrix} 2 \\ 2 \end{pmatrix} = 0 \quad (5)$$

or

$$2\mathbf{c}^T \begin{pmatrix} 2 \\ 2 \end{pmatrix} = 28 \quad (6)$$

or

$$\mathbf{c}^T \begin{pmatrix} 1 \\ 1 \end{pmatrix} = 7 \quad (7)$$

on taking transpose both sides

$$\begin{pmatrix} 1 & 1 \end{pmatrix} \mathbf{c} = 7 \quad (8)$$

Solution(contd)-

It is also given that centre lies on line:

$$(-1 \ 4) \mathbf{x} = -3 \quad (9)$$

Using equations (8) and (9) we have the following linear eqn in two variables:

$$\begin{pmatrix} -1 & 4 \\ 1 & 1 \end{pmatrix} \mathbf{c} = \begin{pmatrix} -3 \\ 7 \end{pmatrix} \quad (10)$$

or

$$\mathbf{c} = \begin{pmatrix} -1 & 4 \\ 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} -3 \\ 7 \end{pmatrix} \quad (11)$$

or

$$\mathbf{c} = \frac{1}{5} \begin{pmatrix} -1 & 4 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} -3 \\ 7 \end{pmatrix} \quad (12)$$

Solution(contd)-

or

$$\mathbf{c} = \frac{1}{5} \begin{pmatrix} 31 \\ 4 \end{pmatrix} \quad (13)$$

or

$$\mathbf{c} = \begin{pmatrix} \frac{31}{5} \\ \frac{4}{5} \end{pmatrix} \quad (14)$$

Now radius can be found by calculating distance between **C** and **A**.
Therefore,

$$radius^2 = (\mathbf{c} - \mathbf{A})^T (\mathbf{c} - \mathbf{A}) \quad (15)$$

Substituting values of **c** and **A** we have:

$$radius^2 = \begin{pmatrix} \frac{21}{5} & \frac{-11}{5} \end{pmatrix} \begin{pmatrix} \frac{21}{5} \\ \frac{-11}{5} \end{pmatrix} \quad (16)$$

Solution(contd)-

or

$$radius^2 = \frac{441}{25} + \frac{121}{25} \quad (17)$$

or

$$radius^2 = \frac{562}{25} \quad (18)$$

or

$$radius = 4.74 \quad (19)$$

Solution Figure-

