Calculus 2, Assignment 4

BM Corser

April 13, 2017

1. (a) In this context, k cannot be 0, since $k = 0$ implies there is no relationship between $\frac{\mathrm{d}T}{\mathrm{d}t}$ and $T_s - T$. This gives us $|k| > 0$. However, I don't see a colloquial reason for $k > 0$, since either or both of T_s and T can be negative. I can see it's in some sense meaningless to take a negative factor, since if a is a factor of b then $-a$ is also a factor of b.

   ...

   After having found a value for T, on the other hand, I can see that a negative value of k would lead to a situation where instead of T approaching T_s as t approaches infinity, T would also go to minus infinity (which breaks the model of reality).

   (b) $\frac{\mathrm{d}T}{\mathrm{d}t} = k(T_s - T)$ is a first order variables separable ordinary differential equation and as such we can write

$$\begin{aligned}
\int \frac{1}{T_s - T}\mathrm{d}T &= k\int \mathrm{d}t \\
k(t+c) &= -\ln(T_s - T) \\
e^{-k(t+c)} &= T_s - T \\
T &= T_s - e^{-k(t+c)}.
\end{aligned}$$

   If $T = T_0$ when $t = 0$ we can write

$$\begin{aligned}
T_0 &= T_s - e^{-kc} \\
e^{-kc} &= T_s - T_0 \\
c &= -\frac{\ln(T_s - T_0)}{k}
\end{aligned}$$

and

$$\begin{aligned}
T &= T_s - e^{-k\left(t - \frac{\ln(T_s - T_0)}{k}\right)} \\
&= T_s - e^{\ln(T_s - T_0) - kt} \\
&= T_s - \frac{e^{\ln(T_s - T_0)}}{e^{kt}} \\
&= T_s - \frac{T_s - T_0}{e^{kt}}.
\end{aligned}$$

(c) We are given $T(0) = 37$ and $T_s = 24$. Let the amount of time between death and discovery be A, now $T(A) = 34$, $T(A + 30) = 32$ and

$$\begin{aligned}
e^{kA} &= \frac{13}{10} \\
kA &= \ln\left(\tfrac{13}{10}\right), \\
e^{k(A+30)} &= \frac{13}{8} \\
kA + k30 &= \ln\left(\tfrac{13}{8}\right).
\end{aligned}$$

Substituting our value for kA into the second equation

$$\begin{aligned}
k &= \frac{\ln\left(\frac{13}{8}\right) - \ln\left(\frac{13}{10}\right)}{30} \\
&= \frac{\ln\left(\frac{5}{4}\right)}{30}.
\end{aligned}$$

(d) With a value for k, our formula for T will feature the structure $e^{\ln\left(\frac{5}{4}\right)\cdot\frac{t}{30}}$ which can be written $\left(e^{\ln\left(\frac{5}{4}\right)}\right)^{\frac{t}{30}}$ and more simply as $\left(\frac{5}{4}\right)^{\frac{t}{30}}$, hence

$$T = T_s - \frac{(T_s - T_0)}{\left(\frac{5}{4}\right)^{\frac{t}{30}}}$$

and and

$$34 = 24 - (-13) \cdot \exp\left(-\frac{A\left(\ln\left(\frac{13}{8}\right) - \ln\left(\frac{13}{10}\right)\right)}{30}\right)$$